


A NEW METHOD FOR α -SUBSTITUTION IN NITROGEN-
CONTAINING HETEROCYCLES


Norihiro Tokitoh, Renji Okazaki, and Naoki Inamoto

Department of Chemistry, Faculty of Science, The University
of Tokyo, Hongo, Bunkyo-ku, Tokyo 113, Japan

α -Siloxamines obtained from tertiary amines via their N-oxides react with some nucleophiles to afford new tertiary amines having an α -substituent derived from the nucleophiles. For example, siloxyammonium salts (2) prepared from N-methyl-1,2,3,4-tetrahydroisoquinoline N-oxide (1) and trialkylsilyl trifluoromethanesulfonate undergo rearrangement in the presence of strong base to give α -siloxamine (3), which can be subjected to nucleophilic substitution reaction with organometallics such as Grignard reagents to afford 1-substituted N-methyl-1,2,3,4-tetrahydroisoquinoline (4).



Similarly, N-methylpiperidine N-oxide (5) can be converted into α -substituted derivatives (6), which has a new α -substituent in exocyclic α -position.



Besides Grignard reagents, trialkylaluminums, silyl enol ethers, silyl cyanides, and trialkylphosphites can also be used as nucleophiles. Of particular note is the accomplishment of the α -substitution of amines by phenyl and vinyl groups with sp^2 carbon as a reacting center.

In view of ready availability of amine N-oxides, a wide variety of nucleophiles, and its complementarity to the electrophilic α -substitution method, the present one-pot sequence provides a new convenient synthetic method for nitrogen-containing heterocycles.

Fuluroide-induced electrophilic transalkylation reaction using the above-mentioned α -siloxamines and alkyl halides will also be described.