

REACTION OF 3-EXOCYCLIC UNSATURATED 2,5-PIPERAZINEDIONE
WITH ALCOHOLIC AND PHENOLIC HYDROXY GROUP

Chung-gi Shin, Yoshiharu Nakajima, and Yasuchika Yonezawa
Faculty of Technology, Kanagawa University, Kanagawa-ku,
Yokohama 221, Japan

In connection with the synthetic study on aspirochlorin, we wish to report the facile synthesis of the framework of didethioaspirochlorin and new syntheses of 3-aminocoumarin and -chroman derivatives.

1,4-Diacetyl-2,5-piperazinedione (PDO = 2,5-piperazinedione) was condensed with salicylaldehyde in the presence of *t*-BuOK in DMF to give (Z)-3-salicylidene-PDO (1) and another unknown product. Subsequently, the latter was treated with 1 M HCl to give known 3-hydroxycoumarin, which was consistent with the product derived by the reaction of 3-(N-acetyl)-aminocoumarin with 1 M HCl.

On the other hand, when a solution of 1 in MeOH was irradiated by UV, the expected coumarin derivative was also obtained quantitatively. From the result, it was supposed that the condensation of PDO with salicylaldehyde gave a mixture of (Z)- and (E)-isomers and that the latter was immediately converted into aminocoumarins.

In order to synthesize the desired spiro-PDO derivative, when (Z)-1 was treated with *t*-BuOCl in CHCl_3 , spiro[3H-3-chlorobenzofuran-2,3'-(1'-acetyl-2',5'-dioxo)-piperazine] (2) was obtained as a mixture of two diastereomers in an almost quantitative yield. Furthermore, in a similar manner, so many spiro derivatives were synthesized and converted into 3-aminocoumarin derivatives.

Interestingly, it was found that further treatment of 2 with NaOMe or ZnCO_3 gave 3-(N-acetylglucyl)amino-3,4-dimethoxy- and 4-chloro-3-methoxychroman in good yields.

