

ARYL MIGRATION OF HETEROAROMATICS HAVING AROYL GROUP


Takeo Higashino, Masumi Takemoto, Akira Miyashita, and

Eisaku Hayashi


Shizuoka College of Pharmacy, 2-2-1 Oshika, Shizuoka 422,

Japan

The reaction of 4-arylquinazolines (1) with sodium hydroxide in dimethyl sulfoxide (DMSO) was found to proceed in two ways. One is the aryl migration to lead to 4-aryl-3,4-dihydro-4-quinazolinecarboxylic acids (2), and the other is the fission of the C⁴-CO bond to yield quinazoline (3) and aroic acids (4).



When the reaction mixture after stirring 7-aryl-3-phenyl-3H-1,2,3-triazolo[4,5-d]pyrimidines (5) with sodium hydroxide in DMSO was subjected to potassium ferricyanide oxidation, 5 was easily converted into the corresponding 7-aryl-3-phenyl-3H-1,2,3-triazolo[4,5-d]pyrimidines (6) by way of 7-aryl-6,7-dihydro-3-phenyl-3H-1,2,3-triazolo[4,5-d]pyrimidine-7-carboxylic acids (7).



In the case of the reaction of 2-arylquinoxalines (8), 2-arylquinoxalines (9), quinoxaline (10), aroic acids (4), and α -aryl- α -(methylsulfinylmethyl)-2-quinoxalinemethanols (11) were obtained.




Chart 3

Mechanism for the aryl migration, similar to that for benzylic acid rearrangement, is proposed.