

NOVEL SYNTHESIS OF HETEROCYCLIC COMPOUNDS HAVING ANGULAR
NITROGEN ATOM BY INTRAMOLECULAR DIELS-ALDER REACTION OF 1-AZADIENES


Masataka Ihara,^a Akihiro Kawaguchi,^a Tomoko Kirihaara,^a Keiichi Fukumoto,^a
and Tetsuji Kametani^b

(a)Pharmaceutical Institute, Tohoku University, Aobayama, Sendai 980,
Japan: (b)Institute of Medicinal Chemistry, Hoshi University, Ebara
2-4-41, Shinagawa-ku, Tokyo 142, Japan

Intramolecular Diels-Alder reaction is one of the most powerful methods for synthesis of polycyclic ring system with stereocontrol. It is expected that the intramolecular cycloaddition of 1-azadienes provides a useful tool for the construction of heterocyclic compounds having angular nitrogen atom, common frameworks of many alkaloids. We have investigated the formation of such system from α,β -unsaturated amides and the subsequent cycloaddition.

(A) Synthesis of Benzo[a]quinolizidines

The 3,4-dihydroisoquinoline(1) was converted into the α,β -unsaturated amides (2) which were heated in the presence of trimethylchlorosilane, triethylamine, and zinc chloride to give benzo[a]quinolizidines(3).



(B) Synthesis of Indolo[a]quinolizidines

According to the similar procedure, 3,4-dihydro- β -carboline was transformed into indolo[a]quinolizidines(4), one of which (4b) is a potential synthetic intermediate of eburnamonine.

(C) Synthesis of Indolizidines

The α,β -unsaturated esters, derived from the amino-acetal(5), was submitted to the intramolecular cycloaddition to afford indolizidines(6).

(D) Synthesis of Quinolizidines: Stereoselective Synthesis of ($\pm$)-Epilupinine

Reduction of the lactam(8), synthesized from the alcohol(7), gave ($\pm$)-epilupinine(9).

