

TOTAL SYNTHESIS OF TRYPTOQUIVALINES

Masako Nakagawa, Manabu Ito, Yuko Hasegawa, Satoko Akashi,

Mikio Taniguchi, and Tohru Hino


Faculty of Pharmaceutical Sciences, Chiba University

1-33 Yayoi-cho, Chiba-shi, 260, Japan

We report a short synthesis of (+) and (-)-tryptoquivaline G, **1b** and (-) and (+)-tryptoquivaline L, **1c** by oxidative double cyclization. We also report the first total synthesis of (+)-tryptoquivaline, **1a** which is a tremorgic mycotoxin.

1. Synthesis of tryptoquivaline G AND L


Oxidation of **2** with NBS in TFA gave **3** and



4 which were deprotected followed by m-CPBA oxidation to give (+)-**1b**, (+)-**1c**, respectively. Analogous reactions starting from L-tryptophan gave (-)-**1c** and (-)-**1b**.

2. Synthesis of tryptoquivaline 1a

The condensation of (S)- α -acetoxyisovaleraldehyde **7** with **6** gave **8**. Acylation



i, **7**, molecular sieves 4A, TsOH, CH_2Cl_2 , r.t.; ii, $\text{CCl}_3\text{CH}_2\text{O}_2\text{CNHCHMe}_2\text{CO}_2\text{C}_6\text{H}_4\text{-p-NO}_2$, **5**, KF, MeCN, 18-crown-6, EtN(i-Pr)₂, 35°C, 4 h; iii, DDQ, CHCl_3 , 30°C, 3 h; iv, H_2 , Pd/C; v, N-iodosuccinimide (3 equiv), $\text{CF}_3\text{CO}_2\text{H}$, 50°C; vi, Zn, AcOH; vii, m- $\text{ClC}_6\text{H}_4\text{CO}_2\text{H}$.

of **8** with **5** followed by DDQ oxidation and debenzylation gave **10b**. The reaction of **10b** with NIS provided **11a** which was treated with Zn-AcOH followed by m-CPBA to give tryptoquivaline **1a**, mp 214-217°C, $[\alpha]_D^{+160}$ (c 0.10, CHCl_3) which was identical with the specimen obtained from *Aspergillus fumigatus*.