

SYNTHESIS OF 3-SUBSTITUTED 4H-1-BENOTHIOPYRANS AND RELATED
HETEROCYCLIC COMPOUNDS

Hiroyuki Nakazumi, • Takashi Endo, Hikaru Sonoda, Toshiyuki Nakaue and
Teijiro Kitao

Department of Applied Chemistry, Faculty of Engineering, University of
Osaka Prefecture, Sakai, Osaka 591, Japan

The introduction of specific substituent group into the 3-position of 2-phenyl 4H-1-benzothiopyran-4-ones (thioflavones) 1 has been examined, because the introduction of a chloromethyl and acetoxymethyl group into the corresponding position of 1,4-naphthoquinones and 4H-1-benzothiopyran-4-one 1,1-dioxides is required for bioactivation. 3-(Formyl)thioflavone 3 was prepared by the Sarett oxidation of 3-(hydroxymethyl)thioflavone 2a with chrominium trioxide. A formyl group of 3 was also converted into cyano, acryl and carboxyl group in good yields. The Meerwein reaction of thiochromone as well as chromone with p-nitrobenzenediazonium ion gave only 3-(4'-nitrophenyl)thiochromone (isothioflavone). It was found that 3-(amino)-thiochromone was easily prepared by reaction of 3-(bromo)thiochromen-4-one with sodium azide.

Condensation of 3-(amino)thiochromone with diethyl ethoxymethylenemalonate and with dimethyl acetylenedicarboxylate gave intermediates, which were thermally cyclized to give 4,10-dihydro-4,10-dioxo-1H-[1]-benzothiopyrano[3,2-b]pyridinecarboxylates. 3-(Formyl)thiochromone was condensed with o-phenylenediamine to give 7-oxo-7,13-dihydro-[1]-benzothiopyrano[2,3-b]-1,5-benzodiazepine. The *in vitro* antimicrobial activity and *in vivo* antitumor activity (lymphocytic leukemia P-388) of 3-substituted thioflavones have been also examined.