

SYNTHESIS OF SULFUR-CONTAINING BICYCLIC β -LACTAMS
USING 1-AZADIENES AS A SYNTHETIC BLOCK

Mitsuo Komatsu, Hirosumi Ogawa, Masa-aki Mohri, Shoichi Kume,

Yoshiki Ohshiro, and Toshio Agawa

Department of Applied Chemistry, Faculty of Engineering, Osaka University

Yamada-oka 2-1, Suita, Osaka 565

Development of synthetic routes to functionalized heterocycles is of great importance and we found azadienes, nitrogen analogs of butadienes, are effective heterocyclic synthons for such purpose. For example, we have reported utilization of azadienes in syntheses of 3- to 5-membered rings having an alkenyl or imidoyl group, pyridines, pyridones, thiazoles, and isothiazoles. In this report, we studied syntheses of sulfur-containing bicyclic β -lactams (thiaalkanams and thiaisoalkanams) using 1-azadienes as a synthetic block.

Reaction of SCl_2 with the bis(alkenyl)- β -lactam 2, prepared from the 1-azadiene 1a, using high dilution method gave the thiaisoheptanam 3a in a good yield along with the hydrolyzed product 3b. Similarly the α -methylene- β -lactam 4 afforded the thiaisoheptanams 5 and 6. Furthermore, the *N*-allyl- β -styryl- β -lactam 7, prepared from the *N*-silylazadiene 1b, reacted with SCl_2 to give the thianonanams 8 and 9.

