

TOTAL SYNTHESIS OF (+)- α -CYCLOPIAZONIC ACID





Hideaki Muratake and Mitsutaka Natsume*

Research Foundation Itsuu Laboratory

2-28-10 Tamagawa, Setagaya-ku, Tokyo 158, Japan

Abstract — (+)- α -Cyclopiazonic acid (2) was synthesized from 1-methoxycarbonyl-4-(3-oxo-1-butyl)indole (1) by way of 17, 18, and 19, using a reaction of MeLi with a pyrroline derivative 33 in the presence of $\text{BF}_3 \cdot \text{Et}_2\text{O}$.

Some time ago, we reported a novel reaction for the preparation of 4-alkylindoles from 1-methoxycarbonylpyrrole.¹ 1-Methoxycarbonyl-4-(3-oxo-1-butyl)indole (1) obtained by this procedure has served as a good starting material in synthesizing a number of ergot alkaloids.² Here we describe a further utilization of 1 for a total synthesis in the racemic form of α -cyclopiazonic acid (2), a mycotoxin isolated first from *Penicillium cyclopium* Westling,^{3,4} accompanied by the related alkaloids, cyclopiazonic acid imine (3) and β -cyclopiazonic acid (bissecodehydro-cyclopiazonic acid) (4).⁵ This study confirms the proposed structures of 2 and 3,⁶ since the latter has been derived from 2 by treatment with aqueous ammonia.⁵



3-Formylindole derivative 7, readily prepared from 1 by way of 5 and 6 using conventional reaction steps depicted in Chart 1, was tosylated to afford 8 in order to make the formyl function sufficiently reactive to a carbanion generated from ethyl isocyanoacetate⁷ and *t*-BuOK in THF. An oxazolidine derivative 9⁸ was a reaction product at first but further addition of *t*-BuOK cleaved the oxazolidine ring to form the desired compounds 10 and 11 in 61% and 5% yields, respectively,

a: HOCH₂CH₂OH, *p*-TsOH, C₆H₆, reflux, 1 h. *b*: 2% KOH in MeOH-DME-H₂O (3:1:1), r.t., 1.5 h. *c*: POCl₃-DMF in Et₂O-DMF, 0°C → r.t., 30 min. *d*: *p*-TsCl, Et₃N, CH₂Cl₂, r.t., 1 h. *e*: CNCH₂COOEt, *t*-BuOK, THF, -81 → -20°C, 1 h; and then *t*-BuOK, -20°C, 30 min. *f*: 50% NaH, DMF, 0°C, 20 min; and then ClCH₂OMe in THF, -83 → -20°C, 30 min. *g*: acetone, *p*-TsOH, r.t., 14 h. *h*: DBU, C₆H₆, reflux, 6 h from 15, 16 or 21, 3.5 h from 17. *i*: (i) Ph₃P=CH₂, THF, -80°C, 20 min; (ii) 50% NaH, DMF-THF (2:3), 0°C → r.t., 1 h. *j*: 1 N HCl in DMSO-EtOH-H₂O (1:1:1), reflux, 2 h. *j'*: 1 N HCl in DMSO-EtOH-H₂O (2:2:1), reflux, 5 h. *k*: ClCOOMe, Et₃N, CH₂Cl₂, -20°C → r.t., 1 h 15 min. *l*: ClCOOCH₂Ph, Et₃N, CH₂Cl₂, -20°C → r.t. *m*: PhSeCl, SiO₂, CH₂Cl₂, -82°C, 15 min and then r.t., 1.5 h. *n*: Raney Ni (W-2), EtOH, reflux, 4 h.

Chart 1

accompanied by the formation of 12 in 15% yield. The geometrical stereostructures of 10 and 11 remain undetermined. The formamide function in 10 and 11 was protected by the methoxymethyl group to give 13 and 14, and the ethylene ketal group was removed to afford 15 and 16, which were ready for the cyclization induced by the intramolecular Michael reaction. This was achieved by heating either 15 or 16 with DBU in refluxing benzene, and tricyclic compounds 17 and 18 were obtained in 50% and 28% or 52% and 27% yields, accompanied by the formation of the third product 19 with inseparable contaminants in a small amount. The respective yields of the three products reflected the ratio of the equilibrium during the Michael addition reaction and this was verified by the observation that the same treatment of 17 as above produced 18 and 19 in 27% and 6% yields with the recovery of 17 in 52% yield.

Stereochemistry of the tricyclic compounds 17 and 18 is determined as follows. The reaction of Ph₃P=CH₂ with 17 or 18, followed by treatment with NaH, furnished a mixture of 20 and 21 with a partial isomerization of the COOEt group in either case, 20 being thermodynamically more favored. This fact was supported by the predominant formation of 20, when 21 was refluxed with DBU. Acid hydrolysis of the N_B-protecting groups of 20 afforded the expected primary amine 22 in 58% yield, together with a by-product 23 in 23% yield. The amino group in 22 was protected again by the benzyloxycarbonyl group to give 24 and this was treated with PhSeCl in the presence of SiO₂⁹ to produce tetracyclic compound 25 having the phenylselenenyl function, which was removed reductively with Raney Ni in refluxing EtOH to yield 26 and 27. The ¹H NMR spectrum¹⁰ of 26 exhibits a big coupling value (*J*=12.5 Hz) between H-4 and H-8, suggesting that 26 possesses an undesired trans stereochemistry of the C/D ring juncture.

The structure of the by-product obtained at the acid hydrolysis of 20 is considered

to be 23 according to the precedent reported by Goto *et al.*,¹¹ as a cyclization product of an $-N=CH_2$ moiety of an intermediate to the C-2 position of the N-tosyl-indole function, which seems to be still nucleophilic probably due to the styrene reactivity. By-product 23 was converted to a methoxycarbonyl derivative 28, whose 1H NMR spectrum¹² revealed the stereochemistry at the C-5, C-6, and C-7 positions as shown by analysis of the coupling pattern of H-5 (d, $J = 6$ Hz) and H-6 (dddd, $J = 12, 6, 3, 3$ Hz), the latter being involved in the long-range coupling with each of the methylene protons at C-3 position. This fact together with the 1H NMR spectrum of 26 confirms the stereostructures of 20 and its COOEt epimer 21, and therefore, strongly suggests those of 17 and 18. The similar acid treatment of 21 afforded another set of compounds 29 and 30, which were separated and characterized after converting 30 into 31. This indicates that the stereocenter of the amino acid ester part remained unaffected by the acid treatment.


The acid hydrolysis of the ketone derivatives 17, 18, and 19 provides key compounds for the total synthesis (Chart 2); *i.e.*, a ketimine derivative 32 was a product from 17, whereas another ketimine 33 was the sole product from either 18 or 19. The structures of 32 and 33 are supported by the following facts, keeping in mind the above argument about the stable nature of the COOEt group with an acid.

(i) The relationship between the C and D rings of 32 is demonstrated to be *cis* by its 1H NMR spectrum,¹³ in which a relatively small coupling value ($J_{4,8} = 7.5$ Hz) compared to that of 26 was deduced from the proton signal at the C-4 position.

(ii) Compound 33 is an epimer of 32 with respect to the configuration of the COOEt group, because 32 was isomerized in part to 33 by heating with DBU. The predominant recovery of 32 implies the thermodynamically stable character of its COOEt function.

(iii) A high field shift of the methyl proton signal (0.78 ppm) of the COOEt group was observed in the 1H NMR spectrum¹⁴ of 33 and this can be explained only by the ring current effect of the indole ring, offering strong evidence for the structure 33, where the methyl group is closely located above the aromatic ring. Therefore, in summary, epimerization at the C-8 position has taken place during the acid treatment of 17 and 18 to yield C/D *cis* ketimines 32 and 33 corresponding to the stereochemistry of α -cyclopiazonic acid (2) and iso- α -cyclopiazonic acid (38).³ At the same time, the structure of 19 is established as shown.

Introduction of the methyl group into the ketimine function requires a quite novel



a: 1 N HCl in EtOH-H₂O (4:1), reflux, 3 h from 17, 4 h from 18, 6 h from 19. b: DBU, C₆H₆, reflux, 3 h. c: (i) BF₃·Et₂O, THF, 0°C, 5 min; (ii) MeLi in Et₂O, -80 - -65°C, 45 min. d: (i) 10% KOH in EtOH-H₂O (3:1), reflux, 3 h; (ii) *p*-TsOEt, EtOH, reflux, 14 h. e: (i) diketene, CH₂Cl₂, reflux, 1 h for 38, 2 h for 2; (ii) *t*-BuOK, THF-EtOH (5:1), 0°C → r.t., 19.5 h for 38, 15 h for 2. f: Et₃N, C₆H₆, reflux, 15 h.

Chart 2

method, since the addition to the ketimine system of an alkylmetallic reagent is generally unattainable, especially with the neighboring hydrogen atoms such as C₇-Me and C₈-H.¹⁵ In order to overcome this difficulty, a special device consisting of an initial formation of a complex with BF₃·Et₂O, followed by the addition of MeLi was applied to 33,¹⁶ utilizing the knowledge about a possible steric hindrance of the COOEt group, which was suggested by the above-mentioned NMR phenomenon. Requisite compounds 34 and 35 were obtained in 40% and 4% yields, respectively, accompanied by the formation of 32 in 9% yield. The tosyl group was removed at this stage by warming 34 and 35 with caustic alkali. Reesterifying the resulting amino acids with ethyl *p*-toluenesulfonate¹⁷ afforded a mixture of 36 and 37 from 34 but only 37 from 35. Compound 36 was treated successively with diketene¹⁸ in CH₂Cl₂ and *t*-BuOK in a mixture of THF and EtOH to produce (±)-iso-α-cyclopiiazonic acid (38), which was identical with the authentic sample (UV, IR, MS, and ¹H NMR) prepared from natural α-cyclopiiazonic acid according to the literature.³ Treating 37 the same as above completed the synthesis of (±)-α-cyclopiiazonic acid (2), identified with the natural specimen (TLC, UV, IR, MS, and ¹H

NMR), confirming its proposed structure. Treatment of (+)-38 with Et₃N in refluxing benzene afforded a 5:2 mixture of (+)-2 and (+)-38, whose recrystallization from MeOH yielded a further crop of (+)-2 in 40% yield.¹⁹

ACKNOWLEDGMENT

We wish to thank Dr. S. Ohmomo of the University of Tsukuba for a generous gift of α -cyclopiazonic acid. A part of this work was supported by a Grant-in-Aid for Scientific Research from the Ministry of Education, Science and Culture, which is gratefully acknowledged.

REFERENCES AND NOTES

1. M. Natsume and H. Muratake, Tetrahedron Lett., 1979, 3477.
2. M. Natsume and H. Muratake, Heterocycles, 14, 1101 (1980); 16, 375, 1481 (1981); H. Muratake, T. Takahashi, and M. Natsume, Heterocycles, 20, 1963 (1983).
3. C. W. Holzapfel, Tetrahedron, 24, 2101 (1968).
4. α -Cyclopiazonic acid has also been isolated from Aspergillus versicolor (Vuill.) Tiraboshi, Aspergillus flavus Link, Aspergillus oryzae Wehmer and Penicillium camemberti. S. Ohmomo, M. Sugita, and M. Abe, Nippon Nôgei-kagaku Kaishi, 47, 57 (1973); K. C. Luk, B. Kobbe, and J. M. Townsend, Appl. Environ. Microbiol., 33, 211 (1977); R. Orth, Ann. Nutr. Aliment., 31, 617 (1977); T. Yokota, A. Sakurai, S. Iriuchijima, and N. Takahashi, Agric. Biol. Chem., 45, 53 (1981); P. Still, C. Eckardt, and L. Leistner, Fleischwirtschaft, 58, 876 (1978).
5. C. W. Holzapfel, R. D. Hutchison, and D. C. Wilkins, Tetrahedron, 26, 5239 (1970).
6. The structure of dioxypyrrolidine part has been already determined. M. J. Nolte, P. S. Steyn, and P. L. Wessels, J. Chem. Soc., Perkin Trans. 1, 1980 1057.
7. U. Schöllkopf, F. Gerhart, and R. Schröder, Angew. Chem., 81, 701 (1969).
8. A single compound 9 with unidentified stereostructure was obtained in 70% yield, when the reaction was stopped at the low temperature stage.
9. D. L. J. Clive, V. Farina, A. Singh, C. K. Wong, W. A. Kiel, and S. M. Menchen, J. Org. Chem., 45, 2120 (1980).
10. ¹H NMR of 26 (CDCl₃, 90 MHz) δ : 1.17 (s, Me), 1.28 (s, Me), 1.41 (t, J = 7.5

- Hz, $\text{COOCH}_2\text{CH}_3$), 1.98 (ddd, $J = 12.5, 10.5, 5$ Hz, H-8), 2.24 (br s, NH) 2.32 (s, $\text{SO}_2\text{C}_6\text{H}_4\text{CH}_3$), 2.68 (dd, $J = 15.5, 10.5$ Hz, H-9), 2.91 (dd, $J = 15.5, 5$ Hz, H'-9), 3.11 (ddd, $J = 12.5, 10, 2$ Hz, H-4), 3.88 (d, $J = 10$ Hz, H-5), 4.35 (br q, $J = 7.5$ Hz, COOCH_2Me), 7.39 (d, $J = 2$ Hz, H-2).
11. S. Nakatsuka, H. Miyazaki, and T. Goto, Chem. Lett., 1981, 407.
12. ^1H NMR of 28 (CDCl_3 , 60°C , 90 MHz) δ : 1.07 (t, $J = 7.5$ Hz, $\text{COOCH}_2\text{CH}_3$), 1.87 (br s, $=\text{C}-\text{CH}_3$), ca. 2.21-2.59 (m, H-7), 2.31 (s, $\text{SO}_2\text{C}_6\text{H}_4\text{CH}_3$), 2.77 (dd, $J = 18.5, 6$ Hz, H-8), 2.97 (dd, $J = 18.5, 9.5$ Hz, H'-8), 3.38 (dddd, $J = 12, 6, 3, 3$ Hz, H-6), 3.74 (s, COOMe), ca. 3.74-4.20 (m, COOCH_2Me), 4.74 (dd, $J = 18, 3$ Hz, H-3), 4.96 and 5.00 (br s each, $=\text{CH}_2$), 5.33 (dd, $J = 18, 3$ Hz, H'-3), 5.43 (d, $J = 6$ Hz, H-5).
13. ^1H NMR of 32 (CDCl_3 , 90 MHz) δ : 1.36 (t, $J = 7.5$ Hz, $\text{COOCH}_2\text{CH}_3$), 2.13 (d, $J = 1.5$ Hz, $\text{C}_7\text{-Me}$), 2.31 (s, $\text{SO}_2\text{C}_6\text{H}_4\text{CH}_3$), 2.66 (dd, $J = 18, 11$ Hz, H-9), 3.01-3.44 (m, H-8 and H'-9), 3.76 (dd, $J = 7.5, 7.5$ Hz, H-4), 4.33 (q, $J = 7.5$ Hz, COOCH_2Me), 4.44 (br d, $J = 7.5$ Hz, H-5).
14. ^1H NMR of 33 (CDCl_3 , 90 MHz) δ : 0.78 (t, $J = 7.5$ Hz, $\text{COOCH}_2\text{CH}_3$), 2.12 (d, $J = 1.5$ Hz, $\text{C}_7\text{-Me}$), 2.31 (s, $\text{SO}_2\text{C}_6\text{H}_4\text{CH}_3$), 2.89-3.38 (m, H-8, H-9, and H'-9), 3.79 (q, $J = 7.5$ Hz, COOCH_2Me), 3.93 (br dd, $J = 8, 7$ Hz, H-4), 4.94 (br d, $J = 8$ Hz, H-5).
15. M. Wada, Y. Sakurai and K. Akiba, Tetrahedron Lett., 25, 1079 (1984), and literatures cited therein.
16. The ready addition of an organometallic reagent to oxiranes and oxetanes with the help of $\text{BF}_3 \cdot \text{Et}_2\text{O}$ has been reported. M. J. Eis, J. E. Wrobel, and B. Ganem, J. Am. Chem. Soc., 106, 3693 (1984).
17. K. Ueda, Bull. Chem. Soc. Jpn., 52, 1879 (1979).
18. C. W. Holzapfel and P. J. Gildenhuys, S. Afr. J. Chem., 30, 125 (1977); Chem. Abstr., 87, 151941e (1977).
19. At the final stage of our work, we received private information from Prof. Kozikowski about his successful total synthesis of ($\pm$)-2. A. P. Kozikowski, M. N. Greco, and J. P. Springer, J. Am. Chem. Soc., 106, 6873 (1984).

Received, 6th February, 1985