

SYNTHESIS OF A NOVEL RING SYSTEM:

PYRIDO[3',2':4,5]ISOTHIAZOLO[2,3-a]BENZIMIDAZOLE

Ricardo Bossio, Stefano Marcaccini,* Valerio Parrini,
and Roberto Pepino



C N R, Centro di studio sulla chimica e la struttura dei composti eterociclici e loro applicazioni, Dipartimento di Chimica Organica, Università di Firenze, via G.Capponi 9, 50121 Firenze, Italy

Abstract- Reaction between 2-mercaptonicotinic acid and *o*-phenylenediamine afforded 2-(2-mercapto-3-pyridyl)benzimidazole which underwent oxidative cyclization with bromine to give pyrido[3',2':4,5]isothiazolo[2,3-a]benzimidazole, a novel ring system.

Among the various synthetic methods leading to the formation of the benzimidazole ring, that described by Phillips¹, involving the reaction between *o*-phenylenediamine and carboxylic acids or anhydrides in acidic medium, is especially noteworthy. This method appears to be generally applicable to the synthesis of 2-alkylbenzimidazoles which are obtained in good yields whereas the yields of 2-arylbenzimidazoles are frequently poor².

Porai-Koshits, Ginzburg and Efros³ stated that good yields of 2-arylbenzimidazoles are achieved on heating *o*-phenylenediamine and aromatic acids in acidic medium in sealed tubes.

A careful examination of the literature showed that no data are available concerning the synthesis of 2-(2-mercaptoaryl or heteroaryl)benzimidazoles, we therefore decided to attempt the synthesis of 2-(2-mercaptophenyl)benzimidazole (Ia) and 2-(2-mercapto-3-pyridyl)benzimidazole (Ib).




This choice was made considering our interest in the synthesis of novel heterocyclic systems having fused imidazole or benzimidazole rings⁴⁻⁸. In fact the position of the mercapto group and the imino one in Ia-b allows oxidative cyclization

leading to the formation of the isothiazole ring. (scheme 1 below)

Scheme 1



The synthesis of Ia-b was accomplished by reacting *o*-phenylenediamine with 2-mercaptobenzoic acid and 2-mercaptopyridonic acid respectively:



The reactions were performed in aqueous hydrochloric acid at 180 °C in sealed tubes giving poor yields of Ia-b. Attempts to improve the yields of Ia-b were unsuccessful. As expected attempts to obtain Ia-b without employing sealed tubes also failed. By treating Ia-b with bromine in acetic acid an oxidative cyclization took place giving benzimidazo[1,2-b][1,2]benzisothiazole (IIa) and pyrido[3',2':4,5]isothiazolo[2,3-a]benzimidazole (IIb) respectively. Compound IIa was previously obtained by Giannola, Bajardi and Palazzo⁹ as a by-product in the synthesis of 12*H*-benzimidazo[2,1-b][1,3]benzothiazin-12-one. The same authors⁹ prepared this compound by reacting *o*-phenylenediamine with 3*H*-1,2-benzodithiole-3-thione.

The structure of pyrido[3',2':4,5]isothiazolo[2,3-a]benzimidazole (IIb) a hitherto unknown ring system, was confirmed by means of mass spectrometry. In fact in the mass spectrum of IIb the ions [M]⁺ *m/z* 225 and [benzene-NS]⁺ *m/z* 122 are detectable and this agrees with the assigned structure.

EXPERIMENTAL

Melting points were obtained in open capillary tubes and are uncorrected. The ir spectra were measured on a Perkin-Elmer 283 spectrophotometer for potassium bromide discs. The mass spectrum of IIb was recorded with a Kratos MS 80 instrument.

2-(2-Mercaptophenyl)benzimidazole (Ia)

A suspension of 2-mercaptobenzoic acid (3g, 19.4mmoles) and *o*-phenylenediamine (2.1 g, 19.4mmoles) in 20% aq. HCl (40 ml) was heated at 180 °C for 45 min in a sealed tube. The dark green reaction mixture was treated with hot water and the resulting solution decolourized with charcoal, and then filtered. Compound Ia (0.52g, 12% yield) was separated out by treating the filtrate with dil. NaOH until the pH was 7.

Mp 241-242 °C from DMF/EtOH; ir: 3420 (cm^{-1}).

Anal. Calcd. for $\text{C}_{13}\text{H}_{10}\text{N}_2\text{S}$: C, 69.00; H, 4.45; N, 12.38. Found: C, 69.10; H, 4.39; N, 12.31.

2-(2-Mercapto-3-pyridyl)benzimidazole (Ib)

This compound (Ib) was prepared starting from *o*-phenylenediamine and 2-mercaptopyridine-3-carboxylic acid following the method described for Ia except that the reaction time was 1.5 h and 8% aq. HCl was employed. Mp 296-297 °C from DMF (14% yield); ir: 3450 (cm^{-1}).

Anal. Calcd. for $\text{C}_{12}\text{H}_9\text{N}_3\text{S}$: C, 63.41; H, 3.99; N, 18.49. Found: C, 63.49; H, 3.88; N, 18.46.

Oxidative Cyclization of Ia-b

General procedure - The theoretical amount of bromine in AcOH was slowly dropped into a well-stirred suspension of I in AcOH maintaining the temperature at 20 °C. The resulting mixture was allowed to react at room temperature for 2 h, then treated with ether and filtered. The solid product was suspended in cold 5% NaHCO_3 solution and then collected by filtration, washed with water, and dried.

Compound IIa was obtained in 80% yield. Mp 283-284 °C from DMF/AcOEt (reported⁹ 284-286 °C). This compound was identical to an authentic specimen.

Compound IIb was obtained in 78% yield. Mp 246-247 °C from DMF/AcOEt; ir: 1420 (cm^{-1}), 735 (cm^{-1}); ms (70 eV): $[\text{M}]^+$ m/z 225, $[\text{benzene-NS}]^+$ m/z 122, $[\text{benzene-NC}]^+$ m/z 102, $[\text{benzene-N}]^+$ m/z 90.

Anal. Calcd. for $\text{C}_{12}\text{H}_7\text{N}_3\text{S}$: C, 63.98; H, 3.13; N, 18.65. Found: C, 63.87; H, 3.06; N, 18.60.

ACKNOWLEDGEMENT

The authors are grateful to Mrs. Brunella Innocenti for elemental analyses.

REFERENCES AND NOTES

* Author to whom correspondence should be addressed.

1. M.A. Phillips, *J. Chem. Soc.*, 1931, 1143.
2. K. Hofmann, "The Chemistry of Heterocyclic Compounds", A. Weissberger, Ed., Interscience Publisher, Inc., New York, 1953, p. 261.
3. B.A. Porai-Koshits, O.F. Ginzburg, and L.S. Efros, *J. Gen. Chem. (U.S.S.R.)*, 1947, 17, 1768; *Chem. Abstr.*, 1948, 42, 5903.
4. E. Belgodere, R. Bossio, R. Cencioni, S. Marcaccini, and R. Pepino, *Heterocycles*, 1984, 22, 807.
5. E. Belgodere, R. Bossio, R. Cencioni, S. Marcaccini, and R. Pepino, *J. Heterocyclic*

- Chem.*, 1984, 21, 1241.
6. E.Belgodere, R.Bossio, S.Marcaccini, S.Parti, and R.Pepino, *Heterocycles*, 1985, 23, 349.
 7. R.Bossio, S.Marcaccini, V.Parrini, and R.Pepino, *Heterocycles*, 1985, 23, 391.
 8. R.Bossio, S.Marcaccini, V.Parrini, and R.Pepino, *J.Heterocyclic Chem.*, in press.
 9. L.I.Giannola, M.L.Bajardi, and S.Palazzo, *J.Heterocyclic Chem.*, 1977, 14, 507.

Received, 17th July, 1985