


REACTIVITY OF 5,6-UNSUBSTITUTED 1,3-DIOXIN-4-ONE AS
AN EQUIVALENT OF FORMYLKETENE OR FORMYLACETIC ACID

Masayuki Sato, Naoki Yoneda, Keiko Sekiguchi, and Chikara Kaneko

Pharmaceutical Institute, Tohoku University, Aobayama, Sendai 980, Japan

We have established recently a general and efficient synthetic method of 5,6-unsubstituted 1,3-dioxin-4-ones (**1**)¹ and demonstrated their successful use as a viable alternative reagent for formyl acetic ester (**2**) in the so-called de Mayo reaction. The method provided a novel way for the introduction of carboxaldehyde and acetic acid appendages at the vicinal position of alkanes starting from corresponding alkenes² and has been applied to a one-pot synthesis of *cis*-2-formyl-5-hydroxy-2-cyclopentene-1-acetic acid γ -lactone, a key intermediate for the synthesis of prostaglandin derivatives.³ It has also become obvious that **1** generates formylketene (**3**) under a quite mild condition (heating at about 100-120°C) in an aprotic solvent. Thus, **1** can be regarded as a synthetic equivalent of formylketene (**3**). **3** reacts *in situ* either with polarized unsaturated functions (1,2-dipoles: $X=Y \leftrightarrow X^--Y^+$) in a 4+2 manner to give a variety of six-membered heterocycles or with hetero- or C-nucleophiles (RXH) to give formylacetylation products.



1. M. Sato, K. Sekiguchi, H. Ogasawara, and C. Kaneko, *Synthesis*, **1985**, 224.
2. M. Sato, H. Ogasawara, K. Sekiguchi, and C. Kaneko, *Heterocycles*, **22**, 2563 (1984).
3. M. Sato, K. Sekiguchi, and C. Kaneko, *Chemistry Lett.*, **1985**, in press.
4. M. Sato, N. Yoneda, and C. Kaneko, *Chem. Pharm. Bull.*, submitted.