

A NOVEL 1,2-ALKYL MIGRATION OF TRIALKYL(4-PYRIDYL)BORATES

Minoru Ishikura, Tsukasa Ohta, and Masanao Terashima*

Faculty of Pharmaceutical Sciences, Higashi-Nippon-Gakuen University,
Ishikari-Tobetsu, Hokkaido 061-02, Japan

Abstract—The reaction of lithium trialkyl(4-pyridyl)borates with Lewis acids or acylating agents produced 4-alkylpyridines via 1,2-alkyl migration from boron to pyridine ring.

Tetracoordinated organoborates which are adjacent to an electron deficient carbon atom are known to be readily accompanied by 1,2-migration of an organyl group from boron to carbon.¹ To our knowledge, the only one example of this type of 1,2-alkyl migration reaction from boron to pyridine ring was reported by Nozaki, i.e., trialkyl(6-bromo-2-pyridyl)borates underwent the 1,2-alkyl shift from boron to 2-position of pyridine with concerted cleavage of pyridine ring to give 5-alkyl-5-dialkylboryl-2,4-pentadienenitriles.² In the course of our studies on pyridylborane compounds,³ we found that 4-alkylpyridines (2) could be produced from trialkyl(4-pyridyl)borates (1) via the 1,2-alkyl migration sequence in the presence of Lewis acids or acylating agents, as shown in Chart 1.




Chart 1

Reaction of 4-lithiopyridine, derived from 4-bromopyridine (1 mol eq) and BuLi (1 mol eq) in Et₂O,⁴ with tributylborane (1 mol eq) under nitrogen atmosphere at -70°C produced lithium tributyl(4-pyridyl)borate (1, R=Bu) in situ, which decomposed on being allowed to stand at room temperature overnight to give a resinous substance.

An attempted reaction of 1 (R=Bu) with alkyl halides (MeI, PhCH₂Br) (-70°C → room temperature) overnight gave no isolable product. Alternatively, substantial amount of 4-butylpyridine was obtained on treatment of 1 (R=Bu) with 2 mol eq of BF₃OEt₂ or (CF₃CO)₂O in Et₂O under nitrogen atmosphere at -70°C, and then room temperature overnight, followed by the oxidation with alkaline hydrogen peroxide. Table summarizes the results of brief studies of the 1,2-alkyl migration reaction of 1 in the presence of Lewis acids or acylating agents.

Table 1,2-Alkyl migration reaction of 1

BR ₃ of <u>1</u>	Reagent	Yield(%) ^{a)} of <u>2</u>	BR ₃ of <u>1</u>	Reagent	Yield(%) ^{a)} of <u>2</u>
BBu ₃	BF ₃ OEt ₂	58 ^{b)}	B(sec-Bu) ₃	BF ₃ OEt ₂	48 ^{c)}
	B(OMe) ₃	38		TiCl ₄	42
	AlCl ₃	20		(CF ₃ CO) ₂ O	44
	TiCl ₄	45		(CH ₃ CO) ₂ O	15
	(CF ₃ CO) ₂ O	56		PhCOCl	34
	(CH ₃ CO) ₂ O	33	B(Hex) ₃	BF ₃ OEt ₂	59 ^{d)}
	PhCOCl	40		(CF ₃ CO) ₂ O	52
Bu-9-BBN	BF ₃ OEt ₂	23	B(cyc-Hex) ₃	BF ₃ OEt ₂	26 ^{e)}
	(CF ₃ CO) ₂ O	20		(CF ₃ CO) ₂ O	23

a) Isolated yield by flash chromatography (SiO₂/hexane:AcOEt=4:1) based on 4-bromopyridine b) 4-Butylpyridine [bp 91°C/18 mmHg (lit.⁵ 84°C/8 mmHg)] c) 4-sec-Butylpyridine [bp 85°C/18 mmHg (lit.⁵ 197°C)] d) 4-Hexylpyridine [bp 140°C/18 mmHg (lit.⁵ 110°C/5 mmHg)] e) 4-Cyclohexylpyridine [bp 135°C/15 mmHg; picrate, mp 148-151°C (lit.⁶ mp 150-151°C)]

Next, the reaction of 1 with 3-indolylacetyl chloride was undertaken. Borate (1; R=Bu, sec-Bu) was reacted with 3-indolylacetyl chloride (1 mol eq) in Et₂O under nitrogen atmosphere (-70°C → room temperature, overnight). After concentration of the reaction mixture in vacuo, the residue was directly subjected to flash chromatography (SiO₂/hexane:AcOEt=4:1) to afford small amount of fairly unstable dieneamide (5a, 10%; 5b, 10%)⁷ together with substantial amount of 4-alkylpyridine (2; R=Bu, 44%; R=sec-Bu, 37%). The formation of dieneamide (5) may be explained by sequences (3 ⇌ 4 ⇌ 5) depicted in Chart 2. The structure of 5 was further confirmed by the immediate conversion to amide (6)⁸ by catalytic hydrogenation over PtO₂ in EtOH under atmospheric pressure.




Chart 2

Typical procedure: 4-Butylpyridine—Tributylborane (1M solution in hexane, 9 ml) was added slowly to an ethereal solution (50 ml) of 4-lithiopyridine, derived from 4-bromopyridine (1.2 g, 9 mmol) and BuLi (1.5M solution in hexane, 6 ml) in Et₂O at -70°C under nitrogen atmosphere, the mixture was gradually warmed to -10°C and then cooled to -30°C. Boron trifluoride etherate (2.3 ml, 18 mmol) was added dropwise, the whole was slowly warmed to room temperature, and then stirred overnight. After treatment with 10% NaOH (20 ml) and 30% H₂O₂ (5 ml) solutions under ice-cooling, the mixture was extracted with AcOEt, the extract was dried over MgSO₄. The solvent was removed and the residue was purified by flash chromatography (SiO₂/hexane:AcOEt=4:1) to give 620 mg (58%) of 4-butylpyridine.

ACKNOWLEDGEMENT

This work was supported in part by a Grant-in-Aid for Scientific Research (No. 59570902) from the Ministry of Education, Science and Culture of Japan.

REFERENCES AND NOTES

1. A. Pelter and K. Smith, "Comprehensive Organic Chemistry," ed. by D. H. R. Barton and W. D. Ollis, Pergamon, Oxford, 1979, vol. 3, p 883.
2. K. Utimoto, N. Sasaki, M. Obayashi and H. Nozaki, Tetrahedron, 1976, 32, 769.
3. M. Ishikura, M. Kamada, I. Oda and M. Terashima, Heterocycles, 1985, 23, 117.
4. F. C. Fischer and E. Havinga, Recueil, 1965, 84, 439.
5. C. T. Kyte, G. H. Jeffery and A. I. Vogel, J. Chem. Soc., 1960, 4454.
6. A. T. Vompe and N. N. Manich, Dokl. Akad. Nauk. SSSR, 1966, 170, 89.
7. 5a; viscous oil. IR(neat) cm^{-1} : 3300, 3108, 1638, 1594. NMR(CDCl_3) δ : 0.50-1.80(m, 9H), 2.50-3.00(m, 1H), 3.87(s, 2H), 4.10-4.60(m, 1H), 5.05 (d, 1H, J=9Hz), 5.20-5.50(m, 1H), 6.72(d, 1H, J=9Hz), 6.80-7.80(m, 5H), 8.50 (br s, 1H). High-MS(m/z) : Calcd for $\text{C}_{19}\text{H}_{22}\text{N}_2\text{O}$: 294.17319. Found: 294.17599.
5b; viscous oil. IR(neat) cm^{-1} : 3300, 2960, 1636, 1596. NMR(CDCl_3) δ : 0.60-1.80(m, 9H), 2.90-3.10(m, 1H), 3.60-4.00(m, 1H), 3.87(s, 2H), 4.60-5.10(m, 2H), 6.73(d, 1H, J=9Hz), 7.00-7.70(m, 5H), 8.20(br s, 1H). High-MS(m/z): Calcd for $\text{C}_{19}\text{H}_{22}\text{N}_2\text{O}$: 294.17319. Found: 294.17338.
8. 6a; syrup. IR(CHCl_3) cm^{-1} : 3488, 3204, 1624, 1620. NMR(CDCl_3) δ : 0.60-1.90(m, 14H), 2.30-3.20(m, 2H), 3.83(s, 2H), 3.50-3.90(m, 1H), 4.50-4.80 (m, 1H), 6.90-7.70(m, 5H), 8.20(br s, 1H). High-MS(m/z): Calcd for $\text{C}_{19}\text{H}_{26}\text{N}_2\text{O}$: 298.20447. Found: 298.20458. 6b; mp 115-116°C (AcOEt-ether). IR(CHCl_3) cm^{-1} : 3488, 3272, 1624. NMR(CDCl_3) δ : 0.60-1.90(m, 14H), 2.30-3.10(m, 2H), 3.84(s, 2H), 3.90-4.20(m, 1H), 4.60-4.90(m, 1H), 7.00-7.80(m, 5H), 8.15(br s, 1H). High-MS(m/z): Calcd for $\text{C}_{19}\text{H}_{26}\text{N}_2\text{O}$: 298.20447. Found: 298.20287.

Received, 30th June, 1986