

SYNTHESIS OF 2H-PYRAZOLO[3,4-b]PYRIDINES FROM
1,1,2,2-TETRACYANOCYCLOPROPANES

Masahiko Takahashi,* Toshiaki Orihara, Takanori Sasaki, and
Tetsuya Yamatera

Department of Industrial Chemistry, Faculty of Engineering,
Ibaraki University, Hitachi, Ibaraki 316, Japan


Ken'ichi Yamazaki and Atomi Yoshida

Research Institute, Daiichi Seiyaku Co., Ltd., Kita-kasai 1-16-13,
Edogawaku, Tokyo 134, Japan

Abstract—Reaction of 3-substituted 1,1,2,2-tetracyanocyclopropanes (1) with phenylhydrazine proceeded through the formation of pyrazole-4-spirocyclopropanes (4) followed by the ring opening of the cyclopropane ring to give 4-vinylpyrazoles (5), from which 2H-pyrazolo[3,4-b]pyridines (6) were obtained.

1,1,2,2-Tetracyanocyclopropanes (1) are expected to show characteristic chemical behaviors because of the presence of two sets of geminal electron-withdrawing cyano groups. The reactivities of 1, however, have attracted less attention in spite of their ready availability,¹ as compared with those of cyclopropanes activated by geminal two electron-withdrawing groups.² Hydrolysis of 3,3-dialkyl-1,1,2,2-tetracyanocyclopropanes followed by thermal ring opening of cyclopropanes was reported to give alkylidenesuccinimides.³ Metallocyclobutane complexes were prepared from 1 on treatment with Pt(Ph₃P)₄ or Pd(Ph₃P)₄.⁴ Recently, 2-azacycloheptatrienylidene was generated thermally from azacycloheptatriene-2-spiro-3'-tetracyanocyclopropanes.⁵ In connection with the recent development in heterocyclic synthesis using cyclopropanes⁶ and cyclopropenes⁷ and with our interest in this field,⁸ we examined the reactivities of 1 and have found a novel route to 2H-pyrazolo[3,4-b]pyridines (6).

3-Alkyl or aryl-1,1,2,2-tetracyanocyclopropanes (1a-e)⁹ were treated with phenylhydrazine in MeOH at room temperature to give 1:1 addition products in



Scheme 1

43-57% yields (Table 1).

In the NMR spectra of the products, no absorptions of methine protons corresponding to the C₃-H of 1 were observed, and therefore cyclopropa[d]pyridazines (2) and pyrazole-4-spirocyclopropanes (4) were excluded as the initial 1:1 addition products (Scheme 1). Alternative possible structures for these products would be 1,2-diazepines (3), produced by the ring expansion of 2 or 4-vinylpyrazoles (5), formed by the ring opening of 4. However, 5 were finally chosen as the 1:1 addition products, since they were easily cyclized, as expected,^{10, 11} to form 2H-pyrazolo[3,4-b]pyridines (6) in 46-81% yields (Table 2). The cyclization conditions depended upon the substituents as follows: a) a mixture of 5a or b and Et₃N in n-BuOH was refluxed, b) a mixture of 5c or d in 90% aq. MeOH containing 1% NaOH was stirred at room temperature, and c) a mixture of 5e in acetic anhydride was refluxed. Although another regioisomers, 1H-pyrazolo[3,4-b]pyridines (7), are possible for these cyclized products, it was difficult to distinguish between 6 and 7 on the basis of the spectral data and some attempted reactions. The structures were determined ultimately to be 6 by X-ray crystallography as

Table 1. Physical and Spectral Properties of Compounds 5

<u>5</u>	Yield (%)	Mp (°C)	MS (M ⁺)	IR (KBr, cm ⁻¹)	¹ H-NMR ^a (δ , ppm)
<u>a</u>	50	266-268	264	3400, 3250, 3150, 2200, 1625, 1600	2.51 (s, 3H), 7.36-8.14 (m, 5H), 9.11 (broad)
<u>b</u>	48	247-249	278	3450, 3250, 3100 2200, 1620, 1560	1.20 (t, J=8 Hz, 3H), 2.89 (q, J=8 Hz, 2H), 7.34-8.16 (m, 5H), 9.21 (s)
<u>c</u>	57	211-213	326	3430, 3275, 3160, 2200, 1630, 1595	6.82-7.25 (m, 10H), 8.65 (broad)
<u>d</u>	43	215-217	360	3450, 3320, 3210, 2200, 1620, 1595	6.63-7.84 (m, 9H), 9.35 (s)
<u>e</u>	45	>300	-	3450, 3300, 3200, 2200, 1650, 1615	7.19-8.25 (m, 9H) 9.05 (broad)

^aThe solvents used were pyridine-d₅ for 5a and 5b, acetone-d₆ for 5c and 5d, and trifluoroacetic acid for 5e.

Table 2. Physical and Spectral Properties of Compounds 6

<u>6</u>	Yield (%)	Mp (°C)	MS (M ⁺)	IR (KBr, cm ⁻¹)
<u>a</u>	73	>300	-	3540, 3410, 3300, 3100, 2200, 1640
<u>b</u>	71	279-281	278	3540, 3260, 3080, 2200, 1640, 1580
<u>c</u>	72	290-292	326	3540, 3300, 3100, 2200, 1635, 1580
<u>d</u>	46	276-278	360	3600, 3420, 3255, 3140, 2200, 1635
<u>e</u>	81	>300	-	3450, 3340, 3220, 2200, 1625, 1605




Fig. 1. Molecular structure of 6c.

shown in the Figure 1. This result is in accord with the observation¹² that acetylation of 3,5-diamino-4-phenylpyrazole occurred on the 3-amino group rather than the 5-amino one.

The present work revealed a new ring opening reaction of tetracyanocyclopropanes, which permits the ready access to multifunctionalized 2H-pyrazolo[3,4-b]pyridines.

EXPERIMENTAL

Melting points were determined on a Yanagimoto micromelting apparatus and are uncorrected. ¹H-NMR, IR, and mass spectra were measured with a JEOL JMX-60, a JASCO A-102, and a JEOL JMS DX-300 spectrometer, respectively. Microanalysis was performed with a Shimadzu UM-3B microanalyzer. X-ray diffraction data were obtained with a Phillips four-circle X-ray autodiffractometer by using Cu K α ($\lambda=1.54178$ Å) radiation.

General Procedure for 3,5-Diamino-1-phenyl-4-(1'-substituted 2',2'-dicyanovinyl)-pyrazoles (5a-e).

To a stirred solution of 1 (4.0 mmol) in MeOH (12 ml), phenylhydrazine (6.0 mmol) was added. The reaction mixture was stirred at room temperature for 12-24 h. The resulting precipitates were collected by filtration and recrystallized from CHCl₃-DMF (5a, b) or MeOH-DMF (5c-e).

Table 3. Elemental Analysis for Compounds 5 and 6

Compounds	Formula	Calculated		Found	
		C	H	C	H
<u>5a</u>	$C_{14}H_{12}N_6$	63.62	4.58	63.49	3.91
<u>5b</u>	$C_{15}H_{14}N_6$	64.73	5.07	65.13	4.75
<u>5c</u>	$C_{19}H_{14}N_6$	69.92	4.32	70.08	4.31
<u>5d</u>	$C_{19}H_{13}N_6Cl$	63.25	3.63	63.25	3.72
<u>5e</u>	$C_{19}H_{13}N_7O_2$	61.45	3.52	61.72	3.75
<u>6a</u>	$C_{14}H_{12}N_6$	63.62	4.58	63.46	4.63
<u>6b</u>	$C_{15}H_{14}N_6$	64.73	5.07	64.76	5.03
<u>6c</u>	$C_{19}H_{14}N_6 \cdot CH_3OH$	67.03	5.06	66.81	5.07
<u>6d</u>	$C_{19}H_{13}N_6Cl$	63.25	3.63	63.03	3.88
<u>6e</u>	$C_{19}H_{13}N_7O_2$	61.45	3.53	61.33	3.73

Cyclization of 5a-e to 4-Substituted 3,6-Diamino-5-cyano-2-phenylpyrazolo[3,4-b]-pyridines (6a-e).

a) Compounds 6a and b: A mixture of 5a or b (0.5 mmol) and Et_3N (1 ml) in *n*-BuOH (20 ml) was refluxed for 8 h. After cooling, the precipitates were collected by filtration and recrystallized from MeOH (6a) or *n*-BuOH (6b).

b) Compounds 6c and d: A mixture of 5c or d (0.5 mmol) in 90% aq. MeOH (20 ml) containing 1% NaOH was stirred at room temperature for 2 h. The precipitates formed were collected by filtration and recrystallized from MeOH-DMF.

c) Compound 6e: A mixture of 5e (0.3 mmol) in acetic anhydride (10 ml) was refluxed for 2 h. After cooling, it was diluted with water (50 ml) and was allowed to stand overnight. The precipitates formed were collected by filtration and recrystallized from water-DMF.

Crystal Data for 6c.

$C_{19}H_{14}N_6 \cdot CH_3OH$ (recrystallized from DMF-MeOH), FW=358.40, triclinic, space group $P\bar{1}$, $a=10.838$, $b=12.409$, $c=7.566$ Å, $\alpha=110.37^\circ$, $\beta=84.52^\circ$, $\gamma=106.80^\circ$, $V=913.18$ Å³, $Z=2$, $D_c=1.304$, $D_m=1.297$ g cm⁻³, $R=5.85\%$, 3257 independent reflections.

REFERENCES

1. A review for malononitrile derivatives: F. Freeman, Synthesis, 1981, 925.
2. S. Danishefsky, Acc. Chem. Res., 1979, 12, 66.
3. H. Hart and F. Freeman, J. Amer. Chem. Soc., 1963, 85, 1161; H. Hart and F. Freeman, J. Org. Chem., 1963, 28, 2063.
4. M. Lenarda, R. Ros, M. Graziani, and U. Belluco, J. Organometal. Chem., 1972, 46, C29; D. J. Yarrow, J. A. Ibers, M. Lenarda, and M. Graziani, J. Organometal. Chem., 1974, 70, 133.
5. S. Murata, T. Sugawara, and H. Iwamura, J. Chem. Soc., Chem. Commun., 1984, 1198.
6. Reviews: R. V. Stevens, Acc. Chem. Res., 1977, 10, 193; E. Wenkert, Heterocycles, 1980, 14, 1703.
7. M. L. Deem, Synthesis, 1982, 701.
8. M. Takahashi, T. Nogami, and K. Nidaira, Heterocycles, 1984, 22, 581; M. Takahashi, N. Takeshi, K. Myojoh, H. Sano, and T. Morisawa, J. Heterocycl. Chem., 1983, 20, 209.
9. H. Hart and F. Freeman, J. Org. Chem., 1963, 28, 1220; Y. C. Kim and H. Hart, Tetrahedron, 1969, 25, 3869.
10. A review for α,β -unsaturated nitriles: M. H. Elnagdi, H. A. Elfahham, and G. E. H. Elgemeie, Heterocycles, 1983, 20, 519.
11. A review for 3(5)-aminopyrazoles: M. H. Elnagdi, F. M. Abdel-Galil, and B. Y. Riad, Heterocycles, 1983, 20, 2437.
12. G. Zvilichovsky and M. David, J. Chem. Soc. Perkin Trans. I, 1983, 11.

Received, 16th June, 1986