



SYNTHESIS OF SUBSTITUTED 4,5-DIHYDRO-4-OXO-2-[(2-TRANS-PHENYLCYCLOPROPYL)AMINO]-3-FURANCARBOXYLIC ACIDS AND ETHYL ESTERS

Vassil St. Georgiev*, Robert A. Mack, and C. Richard Kinsolving


Department of Organic Chemistry, Pennwalt Corporation,
Pharmaceutical Division, Rochester, New York 14623, U.S.A.

Abstract - The synthesis and biological activity of a series of novel 4,5-dihydro-4-oxo-2-[(2-trans-phenylcyclopropyl)amino]-3-furancarboxylic acids and their ethyl esters, is described. When tested in broth and agar dilution tests, some of the title compounds exerted in vitro antimicrobial activity against a number of gram-positive and gram-negative bacteria, as well as in vitro antifungal activity against dermatophyte and yeast fungi.

A number of 2-[(N-substituted)amino]-4,5-dihydro-4-oxo-3-furancarboxylic acids (1) and their esters have been previously prepared as intermediates in the synthesis of various naturally occurring furoquinolines (2)¹⁻⁶, or as products of a heterocyclization reaction that involved a cyclic C-O - insertion of an isocyanate synthon⁷.




In addition, the synthesis of ethyl 2-ethoxy-4,5-dihydro-4-oxo-3-furancarboxylate (3) was described by Mulholland et al.⁸ as part of the preparation of tetronic acid [2,4(3H,5H)-furan-2-one] (4).



In the present study, we wish to report the synthesis and biological activity of a series of novel substituted 4,5-dihydro-4-oxo-2-[(2-trans-phenylcyclopropyl)amino]-3-furancarboxylic acids and their ethyl esters. The preparation of compounds 8 and 9 involved a base-catalyzed cyclocondensation of ethyl 4-chloroacetoacetate (5) with an appropriately substituted trans-phenylcyclopropyl isocyanate (6). The reaction most likely proceeded through the intermediacy of the ketene N-acetal 7 and a C-O -insertion into the ring⁷, to give the desired ethyl 4,5-dihydro-4-oxo-2-[(2-trans-phenylcyc-

1opropyl)amino]-3-furancarboxylate (8). Basic hydrolysis of the latter provided the corresponding free acid 9. Exposure of compound 9 to aqueous solution of sodium carbonate led to the preparation of the sodium salt 10 (Scheme 1).

Scheme 1



When tested for biological activity, the unsubstituted analogs 8a (R= H) and 9a (R= H) demonstrated antimicrobial and antifungal activities. Thus, in broth and agar dilution tests, compounds 8a and 9a exerted in vitro antimicrobial activity against a variety of gram-positive and gram-negative bacteria (Table 1), as well as antifungal activity against Candida albicans and two dermatophytes, Trichophyton mentagrophytes and Microsporum audouinii (Table 2).

Table 1. In vitro Antimicrobial Activity of 4,5-Dihydro-4-oxo-2-[(2-trans-phenylcyclopropyl)amino]-3-furancarboxylic Acids and Ethyl Esters

Broth and Agar Dilution Tests (minimal inhibitory concentration, $\mu\text{g/ml}$)						
Compd	<u>Streptococcus</u> <u>pneumoniae</u>	<u>Streptococcus</u> <u>pyogenes</u>	<u>Neisseria</u> <u>gonorrhoeae</u>	<u>Haemophilus</u> <u>influenzae</u>	<u>Proteus</u> <u>vulgaris</u>	<u>Clostridium</u> <u>histolyticum</u>
8a	318.0 ^{a,b}	318.0 ^a	31.8 ^b	318.0 ^b	-	159.0 ^a
9a	69.0 ^a 460.0 ^b	460.0 ^{a,b}	46.0 ^b	230.0 ^a 69.0 ^b	690.0 ^b	-

^a broth dilution test; ^b agar dilution test.

Table 2. In vitro Antifungal Activity of 4,5-Dihydro-4-oxo-2-[(2-trans-phenylcyclopropyl)amino]-3-furancarboxylic Acids and Ethyl Esters

Broth and Agar Dilution Tests (minimal inhibitory concentration, $\mu\text{g/ml}$)			
Compd	<u>Trichophyton mentagrophytes</u>	<u>Microsporum audouinii</u>	<u>Candida albicans</u>
8a	47.8 ^a	47.8 ^a	478.0 ^{a,b}
9a	-	230.0 ^a	690.0 ^b 460.0 ^a

^a agar dilution test; ^b broth dilution test.

EXPERIMENTAL

Melting points were determined on a Thomas-Hoover capillary melting point apparatus and are uncorrected. The IR spectra were obtained on a Nicolet MX-1 FT spectrometer as KBr discs. The ¹H-NMR spectra were taken on a Varian EM-360A (60 MHz) spectrometer using tetramethylsilane (Me₄Si) as an internal standard. All spectra were consistent with the assigned structures.

Ethyl 4,5-Dihydro-4-oxo-2-[(2-trans-phenylcyclopropyl)amino]-3-furancarboxylate (8a: R= H)

Under a nitrogen atmosphere, triethylamine (4.5 ml) was added dropwise over 15 min to a well-stirred solution of trans-phenylcyclopropyl isocyanate (5.45 g) and ethyl 4-chloroacetoacetate (4.50 g, 27.34 mmol) in petroleum ether-ethyl acetate (10:1), at 5-10°C (ice-water bath). During the addition a heavy precipitate formed. The reaction mixture was stirred at 0-5°C for 1 h, and the precipitate was filtered off under reduced pressure. The residual solid was stirred for 1 h in petroleum ether, then filtered off and stirred with 1 N hydrochloric acid for 1 h. The resulting cake was

rinsed with water and then sucked dry on the filter. Recrystallisation from 2-propanol supplied 5.54 g of the title compound as white crystalline product melting at 140-142°C. Anal. Calcd for $C_{16}H_{17}NO_4$: C, 66.89; H, 5.96; N, 4.88. Found: C, 67.08; H, 6.12; N, 4.95.

Ethyl 4,5-Dihydro-4-oxo-2-[(2-trans-(4-chlorophenyl)cyclopropyl)amino]-3-furancarboxylate (8b):
R = Cl-4)

The title derivative was prepared by a procedure similar to that described for the synthesis of compound 8a, mp 146-148°C (2-propanol). Anal. Calcd for $C_{16}H_{16}ClNO_4$: C, 59.77; H, 5.01; Cl, 11.02; N, 4.35. Found: C, 59.78; H, 5.12; Cl, 11.38; N, 4.17.

Ethyl 4,5-Dihydro-4-oxo-2-[(2-trans-(4-methylphenyl)cyclopropyl)amino]-3-furancarboxylate (8c):
R = CH₃-4)

Compound 8c was obtained by a procedure similar to that described for the synthesis of derivative 8a. It was hydrolyzed to the corresponding free acid 9c without further purification.

4,5-Dihydro-4-oxo-2-[(2-trans-phenylcyclopropyl)amino]-3-furancarboxylic Acid (9a: R = H)

Ethyl 4,5-dihydro-4-oxo-2-[(2-trans-phenylcyclopropyl)amino]-3-furancarboxylate (8a) (15.50 g, 54 mmol) was added to a solution of sodium carbonate (11.45 g, 108 mmol) in 170 ml of water.

Then, steam was passed through the reaction mixture for 2 hours. After refluxing for an additional 60 min, the reaction mixture was cooled to 5-10°C and filtered in order to remove some solid impurities. Acidification to pH 1 with concentrated hydrochloric acid precipitated the free acid 9a. The latter was recrystallized from ethanol to give 7.15 g of pure product, melting at 155-157°C. Anal. Calcd for $C_{14}H_{13}NO_4$: C, 64.89; H, 5.05; N, 5.40. Found: C, 64.63; H, 4.98; N, 5.33.

4,5-Dihydro-4-oxo-2-[(2-trans-(4-chlorophenyl)cyclopropyl)amino]-3-furancarboxylic Acid (9b: R = Cl-4)

The title derivative was prepared by a procedure similar to that described for the synthesis of compound 9a, mp 173-175°C (ethanol). Anal. Calcd for $C_{14}H_{12}ClNO_4$: C, 57.25; H, 4.12; Cl, 12.07; N, 4.77. Found: C, 57.03; H, 4.26; Cl, 12.07; N, 4.75.

4,5-Dihydro-4-oxo-2-[(2-trans-(4-methylphenyl)cyclopropyl)amino]-3-furancarboxylic Acid (9c: R = CH₃-4)

Compound 9c was obtained by a procedure similar to that described for the preparation of derivative 9a, mp 155-157°C. Anal. Calcd for $C_{15}H_{15}NO_4$: C, 65.93; H, 5.53; N, 5.13. Found: C, 66.08; H, 5.60; N, 5.05.

4,5-Dihydro-4-oxo-2-[(2-trans-phenylcyclopropyl)amino]-3-furancarboxylic Acid Sodium Salt (10a: R = H)

The acid 9a (5.05 g, 19.5 mmol) was neutralized with sodium carbonate (2.56 g, 24.2 mmol) in 150 ml of water. The solution was warmed to 50°C, then filtered and concentrated to 50 ml under reduced pressure. The addition of a small amount of ethanol caused a rapid crystallization of the sodium salt 10a. Recrystallization from water provided 4.66 g of pure sodium salt as the monohydrate.

Anal. Calcd for $C_{14}H_{14}NO_5Na$: C, 56.19; H, 4.72; N, 4.68; Na, 7.68. Found: C, 56.42; H, 4.76; N, 4.74; Na, 7.38.

REFERENCES

1. S. Prabhakar, B. R. Pai, and V. N. Ramachandran, Indian J. Chem., 1971, 9, 191.
2. T. R. Govindachari, S. Prabhakar, V. N. Ramachandran, and B. R. Pai, Indian J. Chem., 1971, 9, 1031.
3. I. Mester and M. Ionescu, Phytochemistry, 1971, 10, 2205.
4. V. N. Ramachandran, B. R. Pai, N. Somasundaram, and C. S. Swaminathan, Indian J. Chem., 1973, 11, 1088.
5. H. J. Gais and J. Hafner, Heterocycles, 1976, 4, 1921.
6. T. Yazima and K. Munakata, Agric. Biol. Chem., 1980, 44, 235.
7. L. Capuano and W. Fischer, Chem. Ber., 1976, 109, 212.
8. T. P. C. Mulholland, R. Foster, and D. B. Haydock, J. Chem. Soc., Perkin Trans. 1, 1972, 1225.

Received, 21st July, 1986