


THE CONVERGENT AB + CD ROUTE TO 9-ACETYL-7,8,9,10-TETRAHYDRONAPHTHACENEDIONE

Krishnan Ravichandran, Donald J. Gosciniak, and Michael P. Cava*
Department of Chemistry, University of Alabama, Box H, Tuscaloosa,
AL 35487-9671, U.S.A.

Abstract - The annelated sulfolene derivative (9) has been used as an "AB" synthon in a convergent AB + CD route to 9-acetyl-7,8,9,10-tetrahydro-5,12-naphthacenedione (6), a key precursor to the highly deoxygenated anthracyclinone 4-demethoxy-6,11-dideoxydaunomycinone (5).

Synthetic anthracycline antibiotics may be an exception to the general concept¹ that a synthetic analog of a natural antibiotic is unlikely to be more active than the parent structure. Indeed, the totally synthetic 4-demethoxydaunomycin (2) has been found to be more active² than daunomycin (1); and the less oxygenated analog 4-demethoxy-11-deoxydaunomycin (4) is more effective³ than 3.



1R₁=OMe, R₂=R₃=OH, R₄=daunosaminyl



2R₁=H, R₂=R₃=OH, R₄=daunosaminyl

3R₁=OMe, R₂=H, R₃=OH, R₄=daunosaminyl



4R₁=R₂=H, R₃=OH, R₄=daunosaminyl

5R₁=R₂=R₃=R₄=H

In this regard, we recently reported the first synthesis of 4-demethoxy-6,11-dideoxydaunomycinone (5), the least oxygenated anthracyclinone structure which still contains an unchanged A ring.⁴ In this synthesis the key intermediate 9-acetyl-7,8,9,10-tetrahydro-5,12-naphthacenedione (6) was obtained by a "DCB + A" construction from an anthraquinone precursor.



OR



We now report the synthesis of 6 by a convergent DC + BA route utilizing the annelated sulfolene 9 as the AB source. This approach has been previously unexplored in anthracycline chemistry.⁵ We reasoned that sulfone 9 should be an ideal AB precursor since the cis diene 10 resulting from its thermally induced extrusion of SO₂ should react in a [4 + 2] cycloaddition with an appropriate naphthoquinone derivative [Eq. 2].

The requisite 3,4-bis(bromomethyl)-2,5-dihydrothiophene-1,1-dioxide (8) was easily prepared from the commercially available 2,3-dimethyl-1,3-butadiene by the known procedure.⁷ Treatment of 8 with NaI followed by trapping the resulting diene in situ with methyl vinyl ketone afforded the thermally labile 9 in 60% yield. The structure of 9 was confirmed by its ¹H NMR and IR spectra. Immediate treatment of 9 with an excess of 2-bromo-1,4-naphthoquinone⁸ provided the desired ketone 6 (50%), identical in all respects (NMR, MS, mp) with an authentic sample.⁴

For comparison purposes, we also investigated the assembly of 6 by the conventional AB + CD route, (Eq. 1), a strategy analogous to Wong's⁶ original approach to 4-demethoxydaunomycinone. Surprisingly the requisite starting material 2-acetyl-1,2,3,4-tetrahydronaphthalene (7) was unknown in the literature; it could be obtained in only poor yield (25%) by the treatment of α,α -dibromo-*o*-xylene with methyl vinyl ketone and activated zinc dust. Fusion of 7 with an intimate mixture of phthalic anhydride and AlCl₃-NaCl at 180°C for 10 min, and treatment of the resultant mass with a saturated solution of oxalic acid directly provided 6, albeit in only 30% yield.

EXPERIMENTAL

General Methods. Melting points (mp) were determined on a Thomas-Hoover apparatus and are uncorrected. Mass spectra were determined on a Hewlett-Packard 5985 A system. ¹H NMR spectra were recorded on a Nicolet NT-200 (200 MHz) spectrometer using Me₄Si as an internal standard and are reported in δ units. Elemental analyses were performed by Atlantic Microlab Inc. All organic extracts were washed and dried over anhydrous Na₂SO₄ prior to filtration and evaporation.

2,3,4,5,6,7-Hexahydro-4-acetylthiophene-1,1-dioxide (9). Compound 8 (0.4 g, 1.32 mmol) dissolved in acetone (15 ml) was added in a dropwise manner to a refluxing solution of methyl vinyl ketone (2.0 ml), sodium iodide (0.4 g) and acetone (10 ml) under N₂ atm. After refluxing for 2 h, the solvent was removed under reduced

pressure and the residue was dried under high vacuum (approx. 0.5 torr) to remove all traces of methyl vinyl ketone. Chromatography (SiO_2 , 50:50 hexane-EtOAc) afforded 9 (170 mg, 60%) as an unstable solid which was immediately used in the next step.

^1H NMR 1.71 (m, 2H, CH_2), 2.10 (s, 3H, COCH_3), 2.20 (m, 4H, $2\times\text{CH}_2$), 2.80 (m, 1H, CH), 3.80 (m, 4H, $2\times\text{CH}_2$). IR_{max} 1150, 1295, 1710 cm^{-1} .

2-Acetyl-1,2,3,4-tetrahydronaphthalene (7). Zinc was activated by stirring with a saturated solution of NH_4Cl for 30 min under N_2 atm.

To a stirred suspension of methyl vinyl ketone (3.0 ml) and activated Zn dust (0.4 g) in DMF (25 ml) under N_2 atm, was added a solution of α,α -dibromo-*o*-xylene (2.0 g, 7.57 mmol) in DMF (25 ml). After stirring at room temperature for 6 h, the Zn dust was filtered and the filtrate diluted with H_2O (250 ml). The resulting white polymeric material was filtered and the clear filtrate was extracted with EtOAc, washed with H_2O and then evaporated under reduced pressure. Chromatography (SiO_2 , 80:20 hexane-EtOAc) provided 7 as a colorless oil (0.33 g, 25%), ^1H NMR 1.69 (m, 2H, CH_2), 2.15 (m, 1H, CH), 2.24 (s, 3H, COCH_3), 2.90 (m, 4H, benzylic) 7.10 (s, 4H, aromatic) MS = m/z (relative intensity) 174 (M^+ , 76.3), 159 (84.1), 131 (100). 2,4-DNP derivative: mp 164°C. Anal. calcd. for $\text{C}_{18}\text{H}_{18}\text{N}_4\text{O}_4$: C, 61.02; H, 5.08. Found: C, 60.95, H, 5.15.

9-Acetyl-7,8,9,10-tetrahydronaphthacenedione (6). a) A mixture of 9 (0.15 g, 0.7 mmol), 2-bromo-1,4-naphthoquinone (0.33 g, 1.4 mmol) and NaHCO_3 (25 mg) in DMA (10 ml) was heated for 6 h at 70°C under N_2 atm. After cooling to room temperature, the contents were diluted with H_2O (100 ml) and air bubbled in for 20 min. The solid was filtered and chromatographed (SiO_2 , 70:30 hexane-EtOAc) to afford 6 (60 mg, 50%) identical in all respects with an authentic sample.⁴

b) An intimate mixture of 7 (0.15 g, 0.86 mmol), phthalic anhydride (0.25 g, 1.68 mmol), AlCl_3 (1.2 g, 9.0 mmol) and NaCl (0.32 g, 5.47 mmol) was heated at 180°C for 5 min. The melt was cooled, a saturated solution of oxalic acid (20 ml) was added and the mixture was warmed on a steam bath for 10 min. The product was extracted into ethyl acetate, which was washed with H_2O and evaporated under reduced pressure. Chromatography (SiO_2 , 50:50 hexane- CH_2Cl_2) provided 6 as a light yellow solid (80 mg, 30%), identical in all respects with an authentic sample.⁴

ACKNOWLEDGEMENT

Technical assistance from Ms. V. A. Farr is gratefully acknowledged. This work was supported by NIH-CA 30377.

REFERENCES AND NOTES

1. S. Neidel, Nature (London), 1977, 268, 195.
2. F. Arcamone, L. Bernardi, P. Giardino, B. Patelli, A. DiMarco, A. M. Casazza, G. Pratessi and P. Reggiani, Cancer Treat. Rep., 1976, 60, 829.
3. H. Umezawa, Y. Takahashi, M. Kinoshita, H. Naganawa, K. Tatsuta, and T. Takeuchi, J. Antibiot., 1980, 33, 1581.
4. K. Ravichandran, F. A. J. Kerdesky and M. P. Cava, J. Org. Chem., 1986, 51, 2044.
5. F. Arcamone, "Doxorubicin", Academic Press, New York, 1981.
6. C. M. Wong, D. Popien, R. Schwen and J. T. Raa, Can. J. Chem., 1971, 49, 2712.
7. G. B. Butler and R. M. Ottenbrite, Tet. Lett., 1967, 48, 4873.
8. It was interesting to note that replacing 2-bromo-1,4-naphthoquinone with 2,3-dichloronaphthoquinone and employing identical experimental conditions did not provide the required 6.

Received, 24th July, 1986