

TOTAL SYNTHESSES OF (±)-AGROCLAVINE-I, (±)-6-NOR-
CHANOCLAVINE-II, AND (±)-CHANOCLAVINE-II¹

Masanori Somei,* Fumio Yamada, and Yoshihiko Makita
Faculty of Pharmaceutical Sciences, Kanazawa University
13-1 Takara-machi, Kanazawa 920, Japan

Abstract ——— The first total syntheses of (±)-6-nor-
chanoclavine-II and (±)-chanoclavine-II are achieved. Total
synthesis of (±)-agroclavine-I is also reported.

In the previous papers,² we have established a practical and common synthetic method for ergot alkaloids. The community of the synthetic method is now successfully extended to total syntheses of (±)-agroclavine-I (8),³ (±)-6-nor-chanoclavine-II (11),⁴ and (±)-chanoclavine-II (13).⁵ In this communication, we describe these results in detail.


First, 4,5-trans-4-nitro-5-(2-methyl-1-propen-1-yl)-1,3,4,5-tetrahydrobenz[cd]indole (2) was prepared in four steps in 42% overall yield starting from 3-indolecarbaldehyde (1) as reported previously.² Subsequent treatment of 2 with sodium methoxide in methanol and protonation with 10% aqueous acetic acid readily inverted the stereochemistry of the carbon-4 bound to the nitro group to afford 4,5-cis-4-nitro-5-(2-methyl-1-propen-1-yl)-1,3,4,5-tetrahydrobenz[cd]indole (3)^{6a} in 85% yield together with the recovery of 2 in 8% yield.

With the compound (3) in hand, the syntheses of 8, 11, and 13 were carried out by employing our general synthetic operations.² Thus, reduction of 3 with zinc amalgam in aqueous hydrochloric acid (HCl) gave the corresponding amine (4)^{6b} in 98% yield. Treatment of 4 with methyl chloroformate (ClCOOMe) in methylene chloride (CH₂Cl₂) in the presence of triethylamine (NEt₃) afforded 66% yield of the desired carbamate (5).^{6c} Reduction of 5 with lithium aluminum hydride (LiAlH₄) in absolute tetrahydrofuran (THF) gave (±)-5-epi-6,7-secoagroclavine (6)^{6d} in 91% yield. Subsequent regioselective oxidation of Z-methyl⁷ in the isobutenyl group at the 5-position was attained by the treatment of 6 with selenium dioxide (SeO₂) in dioxane and water to

produce Z-4,5-cis-5-(2-hydroxymethyl-1-propen-1-yl)-4-methylamino-1,3,4,5-tetrahydrobenz[cd]indole [7, ($\pm$)-isochanoclavine-II]^{6e} in 34% yield. Cyclization of 7 with phosphorus oxychloride in the presence of potassium carbonate⁸ produced ($\pm$)-agroclavine-I (8)^{3,6f} in 87% yield. Proton nuclear magnetic resonance (¹H-NMR) spectrum of 8 was identical with that of the authentic ($\pm$)-agroclavine-I, which was synthesized by Ninomiya and co-workers.^{3b}

On the other hand, oxidation of 3 with SeO₂ in dioxane and water gave E-4,5-cis-5-(2-hydroxymethyl-1-propen-1-yl)-4-nitro-1,3,4,5-tetrahydrobenz[cd]indole (9)^{6g} and E-4,5-cis-3-(4-nitro-1,3,4,5-tetrahydrobenz[cd]indol-5-yl)-2-methylacrylic aldehyde (10)^{6h} in 31% and 4% yields,⁷ respectively, together with the recovery of 3 in 62% yield. Reduction of 9 with zinc amalgam in aqueous HCl afforded ($\pm$)-6-nor-chanoclavine-II (11)^{4,6i} in 94% yield. Treatment of 11 with ClCOOMe in CH₂Cl₂ in the presence of NEt₃ afforded 92% yield of the corresponding carbamate (12).^{6j} Finally, reduction of 13 with LiAlH₄ in absolute THF afforded ($\pm$)-chanoclavine-II (13)^{6k} in 69% yield. ¹H-NMR spectrum of 13 was identical with that of the natural alkaloid, reported in the literature.⁵

Thus, the first and simple total syntheses of 11 and 13 were accomplished.



ACKNOWLEDGEMENT

The authors express their cordial gratitude to Prof. I. Ninomiya, Kobe Women's College of Pharmacy, for giving us the ^1H -NMR spectrum of (+)-agroclavine-I.

REFERENCES AND NOTES

1. This report is part XXXVII of a series entitled "The Chemistry of Indoles". Part XXXVI: M. Somei, E. Iwasa, and F. Yamada, *Heterocycles*, **24**, 3065 (1986).
2. M. Somei, Y. Makita, and F. Yamada, *Chem. Pharm. Bull.*, **34**, 948 (1986); M. Somei and F. Yamada, *ibid.*, **32**, 504 (1984).
3. As far as we know, two groups succeeded in the total synthesis of (+)-agroclavine-I: a) A.P. Kozikowski and P.D. Stein, *J. Am. Chem. Soc.*, **107**, 2569 (1985); b) T. Kiguchi, C. Hashimoto, and I. Ninomiya, *Heterocycles*, **23**, 2891 (1985). Formal total synthesis: W.J. Wheeler, *Tetrahedron Lett.*, **27**, 3469 (1986). Isolation: V.G. Sakharovsky and A.G. Kozlovsky, *Tetrahedron Lett.*, **25**, 109 (1984).
4. J.M. Cassady, C.I. Abou-Chaar, and H.G. Floss, *Lloydia*, **36**, 390 (1973).
5. D. Stauffacher and H. Tschertter, *Helv. Chim. Acta*, **47**, 2186 (1964).
6. a) mp 147.0-148.0°C. IR (KBr): 3381, 1620, 1606, 1526, 1444, 1378 cm^{-1} . ^1H -NMR (CDCl_3) δ : 1.67 (3H, d, $J=1.2$ Hz), 1.83 (3H, d, $J=1.5$ Hz), 3.33 (1H, dd, $J=15.4$ and 5.3 Hz), 3.59 (1H, ddd, $J=15.4$, 10.0, and 1.5 Hz), 4.73 (1H, dd, $J=10.3$ and 4.4 Hz), 4.96 (1H, ddd, $J=10.0$, 5.3, and 4.4 Hz), 5.13 (1H, br d, $J=10.3$ Hz), 6.75-6.98 (1H, m), 6.92 (1H, br s), 7.00-7.25 (2H, m), 7.98 (1H, br s). b) mp 113.0-114.0°C. IR (KBr): 2920, 1598, 1572, 756 cm^{-1} . ^1H -NMR (CDCl_3) δ : 1.69 (2H, br s, NH_2), 1.81 (3H, d, $J=1.2$ Hz), 1.85 (3H, d, $J=1.2$ Hz), 2.81 (1H, ddd, $J=15.2$, 7.3, and 1.0 Hz), 3.05 (1H, ddd, $J=15.2$, 4.4, and 1.0 Hz), 3.44 (1H, ddd, $J=7.3$, 4.4, and 3.5 Hz), 3.98 (1H, dd, $J=10.0$ and 3.5 Hz), 5.27 (1H, br d, $J=10.0$ Hz), 6.70-6.91 (1H, m), 6.83 (1H, br s), 6.97-7.19 (2H, m), 8.03 (1H, br s). c) mp 139.0-141.0°C. IR (KBr): 3320, 1683, 1669, 1533 cm^{-1} . ^1H -NMR (CDCl_3) δ : 1.79 (1H, d, $J=2.0$ Hz), 1.81 (1H, d, $J=1.5$ Hz), 2.87 (1H, dd, $J=15.5$ and 7.0 Hz), 3.11 (1H, dd, $J=15.5$ and 4.2 Hz), 3.61 (3H, s), 4.05 (1H, dd, $J=9.8$ and 3.7 Hz), 4.30 (1H, dddd, $J=9.5$, 7.0, 4.2, and 3.7 Hz), 4.75 (1H, br d, $J=9.5$ Hz), 5.21 (1H, br d, $J=9.8$ Hz), 6.68-6.92 (1H, m), 6.85 (1H, br s), 6.96-7.25 (2H, m), 7.93 (1H, br s). d) M. Somei, F. Yamada, Y. Karasawa, and C. Kaneko, *Chemistry Lett.*, **1981**, 615. mp 177.0-178.0°C. IR (KBr): 3140, 3090, 1661, 1616, 1437 cm^{-1} . ^1H -NMR (CDCl_3) δ : 1.57 (1H, s, NH), 1.73 (3H, d, $J=1.2$ Hz), 1.89 (3H, d, $J=1.2$ Hz), 2.51 (3H, s), 2.77 (1H, ddd, $J=15.4$, 10.3, and 1.5 Hz), 2.93-3.22 (2H, m), 4.14 (1H, dd, $J=10.3$ and 3.7 Hz), 5.21 (1H, br d, $J=10.3$ Hz), 6.69-6.87 (1H, m), 6.81 (1H, br s), 6.91-7.15 (2H, m), 8.01 (1H, br s, NH). e) oil. IR (film): 3250, 1617, 1604, 1445 cm^{-1} . ^1H -NMR (pyridine- d_5) δ : 1.92 (3H, d, $J=1.2$ Hz), 2.49 (3H, s), 2.89 (1H, dd, $J=15.4$ and 11.2 Hz), 3.06-3.35 (2H, m), 4.46 (1H, d, $J=12.1$ Hz), 4.54 (1H, dd, $J=10.3$ and 3.7 Hz), 4.78 (1H, d, $J=12.1$ Hz), 5.54 (1H, d, $J=10.3$ Hz), 6.98 (1H, d, $J=6.7$ Hz), 7.05-7.27 (2H, m), 7.34 (1H, d, $J=7.8$ Hz), 11.50 (1H, br s, NH). f) mp 157.0-158.0°C. IR (KBr): 3400, 3100, 2860, 1618, 1607, 1444 cm^{-1} . ^1H -NMR (CDCl_3) δ : 1.63 (3H, br s), 2.56 (3H, s), 2.78 (1H, ddd, $J=15.0$, 10.0, and 1.5 Hz), 2.98 (1H, dd, $J=15.0$ and 4.5 Hz),

3.07 (2H, br s), 3.24-3.49 (1H, m), 3.78-4.08 (1H, m), 5.48 (1H, br s), 6.74-6.98 (1H, m), 6.79 (1H, br s), 6.98-7.18 (2H, m). g) mp 132.0-134.0°C. IR (KBr): 3496, 3246, 1607, 1535, 1370 cm^{-1} . $^1\text{H-NMR}$ (CDCl_3) δ : 1.86 (3H, d, $J=1.5$ Hz), 3.37 (1H, dd, $J=16.0$ and 5.5 Hz), 3.62 (1H, ddd, $J=16.0$, 10.3, and 1.7 Hz), 3.94 (2H, br s), 4.81 (1H, dd, $J=10.3$ and 4.5 Hz), 5.02 (1H, ddd, $J=10.3$, 5.5, and 4.5 Hz), 5.46 (1H, dq, $J=10.3$ and 1.5 Hz), 6.84 (1H, dd, $J=5.8$ and 2.2 Hz), 6.96 (1H, br s), 7.04-7.28 (2H, m), 8.03 (1H, br s). h) oil. IR (film): 1676 (br), 1636, 1544 cm^{-1} . $^1\text{H-NMR}$ (CDCl_3) δ : 1.96 (3H, d, $J=1.5$ Hz), 3.32-3.84 (2H, m), 4.89-5.21 (2H, m), 6.41 (1H, br d, $J=10.2$ Hz), 6.83 (1H, dd, $J=6.2$ and 1.5 Hz), 7.01 (1H, br s), 6.92-7.34 (2H, m), 8.15 (1H, br s, NH), 9.23 (1H, s). i) mp 208.0-210.0°C (dec.). IR (KBr): 3150, 1614, 1597, 1572, 1420 cm^{-1} . $^1\text{H-NMR}$ (pyridine- d_5) δ : 2.06 (3H, d, $J=1.0$ Hz), 2.98 (1H, dd, $J=15.3$ and 7.3 Hz), 3.19 (1H, dd, $J=15.3$ and 3.9 Hz), 3.50-3.77 (1H, m), 4.28 (1H, dd, $J=9.8$ and 3.9 Hz), 4.34 (2H, s), 6.05 (1H, br d, $J=9.8$ Hz), 7.01 (1H, d, $J=6.8$ Hz), 7.09-7.46 (3H, m), 11.55 (1H, br s). j) oil. IR (film): 3350, 1695 cm^{-1} . $^1\text{H-NMR}$ (CDCl_3) δ : 1.86 (3H, d, $J=1.2$ Hz), 2.91 (1H, dd, $J=15.5$ and 6.3 Hz), 3.14 (1H, dd, $J=15.5$ and 4.5 Hz), 3.60 (3H, s), 4.07 (2H, s), 4.14 (1H, dd, $J=9.8$ and 3.7 Hz), 4.20-4.57 (1H, m), 4.89 (1H, br d, $J=9.5$ Hz), 5.56 (1H, br d, $J=9.8$ Hz), 6.64-6.96 (2H, m), 7.00-7.31 (2H, m), 8.04 (1H, br s). k) mp 86.0-88.0°C (recrystallized from CHCl_3). IR (KBr): 3190, 1620, 1605, 1440 cm^{-1} . $^1\text{H-NMR}$ (pyridine- d_5) δ : 2.12 (3H, d, $J=1.0$ Hz), 2.47 (3H, s), 2.95 (1H, dd, $J=15.5$ and 10.0 Hz), 3.10-3.38 (2H, m), 4.28 (2H, s), 4.43 (1H, dd, $J=10.2$ and 3.8 Hz), 6.03 (1H, d, $J=10.2$ Hz), 6.98-7.44 (4H, m), 11.54 (1H, br s).

7. The differences in the direction of oxidation with SeO_2 would be explained by the formation of a complex with the methylamino group in the compound (6), where the SeO_2 lie in the vicinity of Z-methyl group. See also reference 2.
8. Combination of these reagents was first applied for the synthesis of benzo[c]-phenanthridine alkaloids by H. Ishii and co-workers: H. Ishii, I.-S. Chen, S. Ueki, T. Masuda, K. Morita, and T. Ishikawa, Symposium Papers, The 28th Symposium on the Chemistry of Natural Products, Sendai, October 1986, p. 457.

Received, 10th December, 1986