


SYNTHETIC STUDIES ON MITOMYCIN I. SYNTHESIS OF 7-METHOXYMITOSENENE
FROM 6-METHYLINDOLE BY SELECTIVE OXIDATION OF BENZENE PART

Osamu Asano, Shin-ichi Nakatsuka,* and Toshio Goto
Laboratory of Organic Chemistry, Faculty of Agriculture,
Nagoya University, Chikusa, Nagoya 464, Japan

Abstract— Synthesis of 7-methoxymitosene 3, one of the mitosene analog containing common carbon skeleton in mitomycin series was achieved in 19 steps from 6-methylindole 4 by oxidative functionalization methods.


The mitomycins are a class of antibiotics with activity against gram-positive and gram-negative bacteria and also against several kinds of tumors. Mitomycin C is also used clinical anticancer agents. Structures^{1,2,3} of mitomycins were first elucidated in 1962. Since then a lot of synthetic studies⁴ toward mitomycins have been studied and many synthetic methods for mitosene, mitosane analog and so on have been reported. However, the total synthesis of mitomycins was reported only by Kishi and his coworkers.⁵

Although mitomycins have a highly oxidized indole nucleus, there is no useful approach for the synthesis of mitomycins by oxidative functionalization of simple indole. Recently, we have succeeded in the introduction of substituent(s)⁶⁻⁹ on the benzene part of indole derivatives and also in the synthesis of a simple indoloquinone¹⁰ from 6-methylindole. Here, we report the synthesis of 7-methoxymitosene 3 starting from 6-methylindole (4).



mitomycin A : X=OCH₃ (1)

mitomycin C : X=NH₂ (2)




7-methoxymitosene (3)

6-Methylindole **4** was alkylated with NaH and 3-bromopropyl acetate in DMF to afford **5**¹¹ in 89% yield. Carbomethoxy group was introduced at 3-position of **5** in three steps [1) (COCl)₂ in Et₂O, 2) 120°C in tetrachloroethane, 3) MeOH] to give **6**¹² in 52 % overall yield. Acetoxy group of **6** was converted to corresponding tosylate **7**¹³ by two steps [1) 1N NaOH /MeOH, 2) TsCl/Py] in 82 % overall yield. Construction of mitomycin's skeleton was achieved by anion formation at 2-position of **7** with BuLi in THF and intramolecular cyclization to obtain pyrroloindole derivative **8**¹⁴ in 90 % yield. Chloroacetyl group was selectively introduced at 7-position of **8** by treatment with chloroacetyl chloride and AlCl₃ in dichloroethane to afford **9**¹⁵ in 76 % yield. Conversion of chloroacetyl group of **9** to 5-hydroxy compound **10**¹⁶ was achieved by Baeyer-Villiger oxidation of **9** and subsequent hydrolysis of chloroacetoxo group of **10** [1) m-CPBA/Na₂HPO₄/CHCl₃, 2) 1N NaOH in MeOH] in 20 % overall yield. Phenol **11** was oxidized to o-quinone **12**¹⁷ with Fremy's salt [$\cdot\text{O}-\text{N}(\text{SO}_3\text{K})_2$] in acetone-H₂O in 36 % yield. Compound **12** was selectively derived to monomethyl hydroquinone **13**¹⁸ in two steps [1) H₂/10% Pd-C, 2) MeI, K₂CO₃/DMF] in 61% overall yield. Compound **13** was reoxidized with Fremy's salt to produce desired p-quinone **14** in 55 % yield. Thus obtained **14** has the mitomycin's skeleton and the same indoloquinone ring system. Methyl ester group of **14** was converted to carbamate by published procedure⁴ to obtain 7-methoxymitosene **3** [2,3-dihydro-9-hydroxymethyl-7-methoxy-6-methyl-1H-pyrrolo[1,2-a]indole-5,8-dione carbamate] in 52 % overall yield. The spectroscopic data of **3** were completely identical to those of reported 7-methoxymitosene. Thus we could synthesize 7-methoxymitosene **3**, in 19 steps from simple 6-methylindole **4**. Further synthetic studies on mitomycins are under investigation.

REFERENCES AND FOOTNOTES

1. T. Hata, Y. Sano, R. Sugawara, A. Matsumae, K. Kanamori, Y. Shima, and T. Hoshi, *J. Antibiotics*, 1956, **9**, 141, 146.
2. S. Wakaki, K. Marumo, G. Tomioka, E. Shimazu, H. Kato, H. Kamata, S. Kudo, and Y. Fujimoto, *Antibiotics and Chemotherapy*, 1958, **8**, 228.
3. 1) J. S. Webb, D. B. Cosulich, J. H. Mowat, J. B. Patrick, R. W. Broshard, W. E. Meyer, R. P. Williams, C. F. Wolf, W. Fulmor, C. Pidacks, and J. E. Lancaster, *J. Amer. Chem. Soc.*, 1962, **84**, 3185. 2) A. Tulinsky, *ibid.*, 1962, **84**, 3188.



Recently the absolute configuration of mitomycins have been revised.

K. Shirahata and N. Hirayama, *J. Amer. Chem. Soc.*, 1983, **105**, 7199.

U. Hornemann and M. J. Heins, *J. Org. Chem.*, 1985, **50**, 1301.

4. References cited on the previous paper¹⁰.

5. Y. Kishi, *J. Natural. Prod.*, 1979, **42**, 549 and references cited therein.

6. S. Nakatsuka, H. Miyazaki, and T. Goto, *Tetrahedron Lett.*, 1980, **21**, 2817.

7. S. Nakatsuka, H. Miyazaki, and T. Goto, Chemistry Lett., 1981, 407.
8. S. Nakatsuka, O. Asano, and T. Goto, Heterocycles, 1986, 24, 2109.
9. S. Nakatsuka, T. Masuda, O. Asano, T. Teramae, and T. Goto, Tetrahedron Lett., 1986, 27, 4327.
10. S. Nakatsuka, K. Ueda, O. Asano, and T. Goto, Heterocycles, 1987, 26, 65.
11. 5: oil; $^1\text{H-NMR}$ $\delta(\text{CDCl}_3)$ ppm 2.03(3H, s), 2.09 (2H, m), 2.46(3H, s), 4.00(2H, t, $J=7$ Hz), 4.13(2H, t, $J=7$ Hz), 6.39(1H, d, $J=3$ Hz), 6.93(1H, d, $J=3$ Hz), 6.94(1H, d, $J=8$ Hz), 7.06(1H, br.s), 7.45(1H, d, $J=8\text{Hz}$)].
12. 6: mp 75.5-76.0°C; $^1\text{H-NMR}$ $\delta(\text{CDCl}_3)$ ppm 2.03(3H, s), 2.15(2H, m), 2.47(3H, s), 3.87(3H, s), 4.03(2H, t, $J=7$ Hz), 4.18(2H, t, $J=7$ Hz), 7.06(1H, d, $J=9$ Hz), 7.09(1H, s), 7.70(1H, s), 7.99(1H, d, $J=9$ Hz)].
13. 7: mp 112°C; $^1\text{H-NMR}$ $\delta(\text{CDCl}_3)$ ppm 2.20(2H, m), 2.43(3H, s), 2.47(3H, s), 3.88(3H, s), 3.97(2H, t, $J=7$ Hz), 4.19(2H, t, $J=7$ Hz), 6.74(1H, s), 7.07(1H, d, $J=9$ Hz), 7.30(2H, d, $J=8$ Hz), 7.60(1H, s), 7.72(2H, d, $J=8$ Hz), 7.98(1H, d, $J=9$ Hz)].
14. 8: mp 150.5-151.5°C; $^1\text{H-NMR}$ $\delta(\text{CDCl}_3)$ ppm 2.46(3H, s), 2.62(2H, m), 3.26(2H, t, $J=8$ Hz), 3.88(3H, s), 4.08(2H, t, $J=8$ Hz), 7.01(1H, s), 7.04(1H, d, $J=9$ Hz), 7.94(1H, d, $J=9$ Hz)].
15. 9: mp 173°C; $^1\text{H-NMR}$ $\delta(\text{CDCl}_3)$ ppm 2.64(3H, s), 2.68(2H, m), 3.29(2H, t, $J=7$ Hz), 3.90(3H, s), 4.12(2H, t, $J=7$ Hz), 4.82(2H, s), 7.11(1H, s), 8.46(1H, s)].
16. 11: mp 264-265°C; $^1\text{H-NMR}$ $\delta(\text{CDCl}_3\text{-CD}_3\text{OD } 5:1)$ ppm 2.35(3H, s), 2.64(2H, m), 3.24(2H, t, $J=8$ Hz), 3.85(3H, s), 4.07(2H, t, $J=7$ Hz), 7.00(1H, s), 7.40(1H, s)].
17. 12: mp 207-208°C; $^1\text{H-NMR}$ $\delta(\text{CDCl}_3)$ ppm 1.96(3H, d, $J=1$ Hz), 2.60(2H, m), 3.21(2H, t, $J=7$ Hz), 3.83(3H, s), 3.98(2H, t, $J=7$ Hz), 6.83(1H, q, $J=1$ Hz)].
18. 13: mp 190-192°C; $^1\text{H-NMR}$ $\delta(\text{CDCl}_3)$ ppm 2.35(3H, s), 2.61(2H, m), 3.15 (2H, t, $J=7$ Hz), 3.85(3H, s), 3.88(3H, s), 3.96(2H, t, $J=7$ Hz), 6.46(1H, s)].
19. 14: mp 208.5-210°C; $^1\text{H-NMR}$ $\delta(\text{CDCl}_3)$ ppm 1.93(3H, s), 2.58(2H, m), 3.10(2H, t, $J=7.5$ Hz), 3.85(3H, s), 4.04(3H, s), 4.29(2H, t, $J=7.2$ Hz)].

Received, 2nd February, 1987