

METHYL N-ARYLDITHIOCARBAMATES: USEFUL REAGENTS FOR THE ANNE-
LATION OF PYRIMIDINES AND 1,3-OXAZINES TO FIVE-MEMBERED HETEROCYCLIC RINGS

Javier Garín*, María Pilar Loscertales, Enrique Meléndez, Francisco L.
Merchán, Ricardo Rodríguez, and Tomás Tejero

Department of Organic Chemistry, Instituto de Ciencias de Materiales de
Aragón, University of Zaragoza, 50009-Zaragoza, Spain

Abstract- The reaction of methyl N-aryldithiocarbamates with π -excessive
heteroaromatic o-amino acid derivatives leads to the annelation of 4-oxo-
2-thioxopyrimidines, 2,4-dioxopyrimidines or 2-arylamino-1,3-oxazines,
depending on the reaction conditions.

The synthesis of fused bi- and polycyclic compounds by annelation of a pyrimidine
ring to an existing ring constitutes a very large field, which has been recently
reviewed¹. The structure of the required starting compounds is mainly determined
by the nature of the substituents on the pyrimidine ring and, as a general rule,
systems with an amino group adjacent to another functional group are the most wide-
ly used¹. Thus, when a 4-oxopyrimidine is to be synthesized, a ring with an amino
group adjacent to a carboxylic acid derivative is generally employed, while 2-oxo
and 2-thioxo groups are normally introduced by means of isocyanates and ureas or
their sulfur analogues in the case of the latter.

In the present paper, and as a continuation of our studies in the field of fused
heterocyclic compounds², we report a relatively simple general method for the anne-
lation of pyrimidine and 1,3-oxazine rings to five-membered heteroaromatic systems,
by the reaction of the appropriate o-amino acid derivatives with methyl N-aryldi-
thiocarbamates³ 3.

This method has certain advantages compared to previously reported methods:


- Starting compounds 3 are stable solids, obtained in one step from the correspon-
ding anilines.
- The same reagent (i.e. 3) is used to prepare both the 2,4-dioxo and the 4-oxo-
2-thioxo derivatives.
- This procedure can be modified to prepare fused 2-arylamino-1,3-oxazine derivati-
ves which are uncommon. Some 6-amino-4-oxo-1H,4H-pyrazolo [3,4-d] [1,3] oxazines
have in fact been prepared by us⁴, although the corresponding 2-arylamino-4H,

4-oxo-5,6,7,8-tetrahydro[1]benzothieno[2,3-d][1,3]oxazines 15 are, to the best of our knowledge, still unknown.

As can be seen, a large number of structures of potentially active compounds, derived either from benzo[b]thiophene⁵ or from pyrazole -including analogues of the xanthine oxidase inhibitor oxyallopurinol⁶ - can be easily prepared in one step. For the sake of simplicity, the present paper has been arranged according to the most outstanding features of the newly-formed ring and the nomenclature of the final products is based on this criterion.

4-Oxo-2-thioxopyrimidines

Compounds 6 and 7 are easily obtained as shown in Scheme I:



For 1, 4 and 6: X=N, Y=N-C₆H₅, R¹=H

2, 5 and 7: X=C, Y=S, R¹-R²=(CH₂)₄

Ar: see tables

- Scheme I -


Compounds 6, 5-aryl-4-oxo-1-phenyl-6-thioxo-1H-4,5,6,7-tetrahydropyrazolo [3,4-d] pyrimidines, were prepared from 5-amino-4-ethoxycarbonyl-1-phenylpyrazole⁷ 1 and the corresponding methyl N-aryldithiocarbamate 3. A basic medium is necessary in order to achieve greater reactivity of the o-aminoester and unless sodium hydroxide

is used, extensive decomposition occurs and yields are very low.

Compounds 7, 3-aryl-4-oxo-2-thioxo-1,2,3,4,5,6,7,8-octahydro[1]benzothieno[2,3-d]pyrimidines, are similarly obtained by refluxing the mixture for 7 to 10 h.

With longer reaction times several by-products are formed, perhaps due to the rearrangement that 2-aminothiophenes undergo in a basic medium⁸, especially when electron-withdrawing groups are attached to the thiophene ring, as is the case with 2.

The structure of compounds 7 was confirmed by an independent synthesis in which 3 were also used as starting products. These were converted⁹ to the corresponding arylisothiocyanates, which then reacted with 2 to give the thioureas 5. These were isolated and cyclized using ethanolic potassium hydroxide^{10,11}. (Scheme II).



- Scheme II -

The main spectral features of compounds 6 are summarized below.

Ir (Nujol) and ¹H-nmr spectra of compounds 6

Compound	Ar	ν(cm ⁻¹)		δ(ppm); solvent CF ₃ -COOH		
		NH	C=O	H-3	Aromatics	Me
<u>6a</u>	C ₆ H ₅	3120	1725	8.45	7.55-7.10(m,10H)	--
<u>6b</u>	4-Me-C ₆ H ₄	3120	1705	8.50	7.6(s,5H); 7.5-7.0(m,4H)	2.35(s,3H)
<u>6c</u>	4-MeO-C ₆ H ₄	3120	1725	8.50	7.6(s,5H); 7.2(s,4H)	4.0(s,3H)
<u>6d</u>	4-Cl-C ₆ H ₄	3200-3120	1725	8.50	7.7-7.1(m,9H)	--
<u>6e</u>	2,4-di Me-C ₆ H ₃	3350	1725	8.50	7.6(s,5H); 7.3-7.0(m,3H)	2.4(s,3H); 2.1(s,3H)
<u>6f</u>	3,5-di Me-C ₆ H ₃	3120	1725	8.50	7.6(s,5H); 7.3-6.8(m,3H)	2.35(s,6H)


Compounds 7 show medium absorptions in the $3140\text{--}3100\text{ cm}^{-1}$ range ($\nu\text{N-H}$) and strong absorptions around 1230 cm^{-1} , attributable to $\nu(\text{C=S})$. In their nmr spectra, the tetramethylene bridge shows its characteristic multiplets at $\delta\text{ 3.1--2.7 ppm}$ (H-5 and H-8, 4H) and $\delta\text{ 2.1--1.8 ppm}$ (H-6 and H-7, 4H). Additional data with regard to these compounds are shown in the table.

Ir (Nujol) and ^1H -nmr spectra of compounds 7

Compound	Ar	$\nu(\text{C=O})\text{cm}^{-1}$	$\delta\text{ppm; solvent CF}_3\text{COOH}$	
			Ar	Me
<u>7a</u>	C_6H_5	1710	7.8-7.3(m, 5H)	--
<u>7b</u>	4-Me- C_6H_4	1715	7.6-7.2(m, 4H)	2.5(s, 3H)
<u>7c</u>	4-MeO- C_6H_4	1715	7.4-7.0(m, 4H)	4.0(s, 3H)
<u>7d</u>	4-Cl- C_6H_4	1720	7.7-7.2(m, 4H)	--

2,4-Dioxypyrimidines and 2-arylamino-1,3-oxazines

The synthesis of the pyrazole derivatives 11, 6-arylamino-4-oxo-1-phenyl-1H,4H-pyrazolo[3,4-d][1,3]oxazines, and 13, 5-aryl-4,6-dioxo-1-phenyl-4,5,6,7-tetrahydro-1H-pyrazolo[3,4-d]pyrimidines is easily carried out by the reaction of the potassium salt 9 with 3 in the presence of red mercury (II) oxide in dimethylformamide. When the mixture is maintained at room temperature the kinetic isomers 11 are formed after acidification (Scheme III).




The reaction seems to proceed via the thiourea 4, which is desulfurated to the corresponding unstable carbodiimide, which is cyclized to the oxazine anion 10. Acidic treatment affords 11 in moderate to good yields. When the reaction is conducted at reflux temperature, anion 10 in all probability rearranges to the thermodynamically more stable anion 12 which, after acidic work-up, gives 13. Nevertheless, yields are only moderate and an alternative procedure for the synthesis of 13 was devised, namely, direct isomerization of compounds 11 by simply heating in acidic medium. Of all the acids used sulfuric acid turned out to give the best results. The ^1H -nmr spectra of compounds 11 and 13 (in $\text{CF}_3\text{-COOH}$) are quite similar to those of 6 and are not gathered here. The most outstanding difference between 11 and 13 lies in their $\nu(\text{C}=\text{O})$ absorptions, with compounds 11 showing a strong band at a higher wavenumber (lactone type) than 13 (lactam and urea types).

Ir (Nujol) carbonyl absorptions of isomers 11 and 13

Ar	Compound	$\nu(\text{C}=\text{O})\text{cm}^{-1}$	Compound	$\nu(\text{C}_4=\text{O})\text{cm}^{-1}$	$\nu(\text{C}_6=\text{O})$
C_6H_5	<u>11a</u>	1760	<u>13a</u>	1735	1670
4-Me- C_6H_4	<u>11b</u>	1770	<u>13b</u>	1735	1670
4-MeO- C_6H_4	<u>11c</u>	1770	<u>13c</u>	1740	1670
4-Cl- C_6H_4	<u>11d</u>	1780	<u>13d</u>	1740	1660
2,4-diMe- C_6H_3	<u>11e</u>	1780	<u>13e</u>	1740	1660
3,5-diMe- C_6H_3	<u>11f</u>	1760	<u>13f</u>	1740	1675

Depending on reaction conditions the reaction of 3 with sodium salt 14 in the presence of red mercury (II) oxide gives 2-arylamino-4H-4-oxo-5,6,7,8-tetrahydro-[1]benzothieno[2,3-d][1,3]oxazines 15 or 3-aryl-2,4-dioxo-1,2,3,4,5,6,7,8-octahydro-[1]benzothieno[2,3-d]pyrimidines 16. (Scheme IV).



- Scheme IV -

In this case sodium salt was used since it gave better results than the corresponding potassium salt.

The ^1H -nmr spectra of compounds 15 and 16 (in $\text{CF}_3\text{-COOH}$) are very similar to those of 7, the most significant difference being in their ir spectra, as shown below.

Ir (Nujol) carbonyl absorptions of isomers 15 and 16

Ar	Compound	$\nu(\text{C=O})\text{cm}^{-1}$	Compound	$\nu(\text{C}_4=\text{O})\text{cm}^{-1}$	$\nu(\text{C}_2=\text{O})$
C_6H_5	<u>15a</u>	1745	<u>16a</u>	1730	1665
4-Me- C_6H_4	<u>15b</u>	1745	<u>16b</u>	1725	1665
4-MeO- C_6H_4	<u>15c</u>	1745	<u>16c*</u>	--	--
4-Cl- C_6H_4	<u>15d</u>	1745	<u>16d</u>	1730	1665

* Compound 16c could not be isolated.

To sum up, the main advantage of our procedure is that not only pyrimidinediones 13 and 16 can be prepared in one step - a typical reaction with isocyanates^{12,13} - but also that their previously unreported isomers 11 and 15 can be synthesized under smooth conditions. This is of great interest since fused 2-amino-1,3-oxazines are quite elusive compounds¹⁴, mainly due to the fact that reaction conditions tend to favour their thermodynamic isomers.

EXPERIMENTAL

Melting points were determined using a Büchi 510 apparatus and are uncorrected. Ir spectra were recorded on a Perkin-Elmer 283 instrument. ^1H -Nmr spectra were obtained on a Bruker WP-80-CW spectrometer with TMS as internal reference.

5-Aryl-4-oxo-1-phenyl-6-thioxo-1H-4,5,6,7-tetrahydropyrazolo[3,4-d]pyrimidines 6

(General Procedure)

To a solution of 5-amino-4-ethoxycarbonyl-1-phenylpyrazole⁷ 1 (1.155 g, 5 mmol) in DMF (10 ml), aqueous 20 M sodium hydroxide (1 ml) was added. The mixture was stirred for 30 min and then the corresponding methyl N-aryldithiocarbamate³ 3 (5 mmol) in DMF (10 ml) was added dropwise. Once all 3 had been added, the mixture was refluxed for 6-8 h and then cooled. The white precipitate (sodium formiate) was filtered off and the filtrate poured into water (200 ml) and acidified with concentrated hydrochloric acid (until pH=5) in an ice-water bath. The solid thus obtained was filtered off, dried, washed twice with ether and recrystallized.

6a: Yield 54%. Mp 193-195 °C(EtOH/H₂O). Anal. Calcd. for $\text{C}_{17}\text{H}_{12}\text{N}_4\text{O}_2\text{S}$ (320.37): C, 63.73%; H, 3.78; N, 17.49. Found: C, 63.92; H, 3.64; N, 17.69.

6b: Yield 65%. Mp 198-200 °C (EtOH/H₂O). Anal. Calcd. for C₁₈H₁₄N₄OS (334.40): C, 64.65%; H, 4.22; N, 16.76. Found: C, 64.45; H, 4.08; N, 16.95.

6c: Yield 50%. Mp 235-237 °C (EtOH/H₂O). Anal. Calcd. for C₁₈H₁₄N₄O₂S (350.40): C, 61.70%; H, 4.03; N, 15.99. Found: C, 61.88; H, 3.89; N, 16.16.

6d: Yield 58%. Mp 230-232 °C (EtOH). Anal. Calcd. for C₁₇H₁₁ClN₄OS (354.82): C, 57.54%; H, 3.13; N, 15.79. Found: C, 57.38; H, 3.30; N, 15.68.

6e: Yield 60%. Mp 123-125 °C (EtOH/H₂O). Anal. Calcd. for C₁₉H₁₆N₄OS (348.42): C, 65.49%; H, 4.63; N, 16.08. Found: C, 65.55; H, 4.70; N, 16.00.

6f: Yield 55%. Mp 131-133 °C (EtOH/H₂O). Anal. Calcd. for C₁₉H₁₆N₄OS (348.42): C, 65.49%; H, 4.63; N, 16.08. Found: C, 65.46; H, 4.69; N, 16.10.

N-(3-Ethoxycarbonyl-4,5,6,7-tetrahydrobenzo[b]thien-2-yl)-N'-arylthioureas 5

(General Procedure)

A mixture of 2-amino-3-ethoxycarbonyl-4,5,6,7-tetrahydrobenzo[b]thiophene 2 (1.125 g, 5 mmol) and the corresponding arylisothiocyanate⁹ 8 (5 mmol) in ethanol (5 ml) was refluxed for 8-10 h and then cooled. The resulting precipitate was filtered off, washed with ethanol and dried.

5a: Yield 85%. Mp 182-183 °C. Anal. Calcd. for C₁₈H₂₀N₂O₂S₂ (360.49): C, 59.97%; H, 5.59; N, 7.77. Found: C, 60.06; H, 5.66; N, 7.69.

5b: Yield 75%. Mp 160-161 °C. Anal. Calcd. for C₁₉H₂₂N₂O₂S₂ (374.51): C, 60.93%; H, 5.92; N, 7.48. Found: C, 60.99; H, 5.98; N, 7.39.

5c: Yield 80%. Mp 174-176 °C. Anal. Calcd. for C₁₉H₂₂N₂O₃S₂ (390.51): C, 58.43%; H, 5.68; N, 7.18. Found: C, 58.38; H, 5.72; N, 7.15.

5d: Yield 78%. Mp 205-207 °C. Anal. Calcd. for C₁₈H₁₉ClN₂O₂S₂ (394.93): C, 54.74%; H, 4.85; N, 7.09. Found: C, 54.70; H, 4.81; N, 7.13.

3-Aryl-4-oxo-2-thioxo-1,2,3,4,5,6,7,8-octahydro[1]benzothieno[2,3-d]pyrimidines 7

(General Procedure)

METHOD A: Potassium hydroxide (0.28 g, 5 mmol) in ethanol (25 ml) was added to a solution of the corresponding thiourea 5 (5 mmol) in ethanol (25 ml). The mixture was refluxed for 2 h, cooled, poured into water (500 ml) -in an ice-water bath- and neutralized to litmus with concentrated hydrochloric acid. The white solid was filtered off, washed first with water then with ether and dried.

METHOD B: Aqueous 10M sodium hydroxide (0.5 ml) was added to a solution of 2-amino-3-ethoxycarbonyl-4,5,6,7-tetrahydrobenzo[b]thiophene 2 (1.125 g, 5 mmol) in DMF (10 ml). The mixture was stirred for 30 min and then the corresponding methyl N-aryldithiocarbamate 3 (5 mmol) in DMF (10 ml) was added dropwise. The mixture

was refluxed for 7-10 h, cooled, filtered and the filtrate was poured into cold water (200 ml) and acidified with concentrated hydrochloric acid (until pH=4-5). The precipitate was filtered off, washed with water, dried and recrystallized.

7a: Yield 97%(A), 65%(B). Mp 284-286 °C(CH₃CN). Anal. Calcd. for C₁₆H₁₄N₂O₂S₂ (314.42): C, 61.12%; H, 4.49; N, 8.91. Found: C, 61.07; H, 4.54; N, 8.99.

7b: Yield 93%(A), 75%(B). Mp 276-278 °C(CH₃CN). Anal. Calcd. for C₁₇H₁₆N₂O₂S₂ (328.45): C, 62.16%; H, 4.91; N, 8.53. Found: C, 62.19; H, 4.87; N, 8.59.

7c: Yield 98%(A), 74%(B). Mp 278-280 °C(CH₃CN). Anal. Calcd. for C₁₇H₁₆N₂O₂S₂ (344.45): C, 59.27%; H, 4.68; N, 8.13. Found: C, 59.33; H, 4.61; N, 8.09.

7d: Yield 93%(A), 80%(B). Mp 286-288 °C(CH₃CN). Anal. Calcd. for C₁₆H₁₃ClN₂O₂S₂ (348.87): C, 55.08%; H, 3.76; N, 8.03. Found: C, 55.12; H, 3.74; N, 8.07.

6-Arylamino-4-oxo-1-phenyl-1H,4H-pyrazolo[3,4-d][1,3]oxazines 11 (General Procedure)

To a suspension of potassium 5-amino-1-phenylpyrazole-4-carboxylate 9 (0.723 g, 3 mmol) and red mercury (II) oxide (0.756 g, 3.5 mmol) in DMF (10 ml), a solution of the corresponding methyl N-aryldithiocarbamate 3 (3 mmol) in DMF (10 ml) was added dropwise. The mixture was stirred for 15 h at room temperature and then filtered. The filtrate was poured into water (200 ml) -ice-water bath- and acidified with concentrated hydrochloric acid (until pH=5). The crude solid was filtered off, dried, washed with ether and recrystallized from acetonitrile.

11a: Yield 65%. Mp 220-222 °C. Anal. Calcd. for C₁₇H₁₂N₄O₂ (304.30): C, 67.09%; H, 3.98; N, 18.41. Found: C, 67.13; H, 4.01; N, 18.36.

11b: Yield 65%. Mp 222-224 °C. Anal. Calcd. for C₁₈H₁₄N₄O₂ (318.32): C, 67.91%; H, 4.43; N, 17.60. Found: C, 67.95; H, 4.40; N, 17.56.

11c: Yield 64%. Mp 209-211 °C. Anal. Calcd. for C₁₈H₁₄N₄O₃ (334.32): C, 64.66%; H, 4.22; N, 16.76. Found: C, 64.70; H, 4.28; N, 16.71.

11d: Yield 55%. Mp 227-229 °C. Anal. Calcd. for C₁₇H₁₁ClN₄O₂ (338.76): C, 60.27%; H, 3.27; N, 16.54. Found: C, 60.31; H, 3.33; N, 16.48.

11e: Yield 34%. Mp 165-167 °C. Anal. Calcd. for C₁₉H₁₆N₄O₂ (332.35): C, 68.66%; H, 4.85; N, 16.86. Found: C, 68.60; H, 4.88; N, 16.93.

11f: Yield 65%. Mp 209-211 °C. Anal. Calcd. for C₁₉H₁₆N₄O₂ (332.35): C, 68.66%; H, 4.85; N, 16.86. Found: C, 68.71; H, 4.83; N, 16.90.

5-Aryl-4,6-dioxo-1-phenyl-4,5,6,7-tetrahydro-1H-pyrazolo[3,4-d]pyrimidines 13

(General Procedure)

METHOD A: This method is identical to the one described for the synthesis of compounds 11, except that after the addition of 3 the reaction mixture was heated

under reflux for 4 h.

METHOD B: A solution of the corresponding compound 11 (2 mmol) in DMF (20 ml) containing four drops of concentrated sulfuric acid was heated at 100 °C for 2-3 h, then cooled and poured into water (100 ml). The mixture was stirred for 30 min and the precipitate was filtered off, dried, washed with ether and recrystallized.

13a: Yield 25%(A), 60%(B). Mp 225-227 °C(n-butanol). Anal. Calcd. for $C_{17}H_{12}N_4O_2$ (304.30): C, 67.09%; H, 3.98; N, 18.41. Found: C, 67.02; H, 3.92; N, 18.39.

13b: Yield 24%(A), 52%(B). Mp 275-277 °C (n-butanol). Anal. Calcd. for $C_{18}H_{14}N_4O_2$ (318.32): C, 67.91%; H, 4.43; N, 17.60. Found: C, 67.87; H, 4.48; N, 17.55.

13c: Yield 54%(A), 67%(B). Mp 208-210 °C. Anal. Calcd. for $C_{18}H_{14}N_4O_3$ (334.32): C, 64.66%; H, 4.22; N, 16.76. Found: C, 64.60; H, 4.29; N, 16.80.

13d: Yield 49%(A), 75%(B). Mp 256-258 °C(n-butanol). Anal. Calcd. for $C_{17}H_{11}ClN_4O_2$ (338.74): C, 60.27%; H, 3.27; N, 16.54. Found: C, 60.23; H, 3.24; N, 16.59.

13e: Yield 30%(A), 51%(B). Mp 230-232 °C. Anal. Calcd. for $C_{19}H_{16}N_4O_2$ (332.35): C, 68.66%; H, 4.85; N, 16.86. Found: C, 68.59; H, 4.90; N, 16.83.

13f: Yield 31%(A), 60%(B). Mp 263-265 °C(n-butanol). Anal. Calcd. for $C_{19}H_{16}N_4O_2$ (332.35): C, 68.66%; H, 4.85; N, 16.86. Found: C, 68.62; H, 4.79; N, 16.88.

2-Arylamino-4H,4-oxo-5,6,7,8-tetrahydro[1]benzothieno[2,3-d][1,3]oxazines 15 (General Procedure)

To a suspension of sodium 2-amino-4,5,6,7-tetrahydrobenzo[b]thiophene-3-carboxylate 14 (1.095 g, 5 mmol) and red mercury (II) oxide (1.296 g, 6 mmol) in DMF (20 ml), a solution of the corresponding methyl N-aryldithiocarbamate 3 (5 mmol) in DMF (20 ml) was added dropwise. The mixture was then stirred for 20-24 h at room temperature and filtered. The filtrate was poured into cold water (400 ml) and acidified until pH=5 with concentrated hydrochloric acid. The resulting precipitate was filtered off, dried and recrystallized from n-butanol.

15a: Yield 72%. Mp 218-220 °C. Anal. Calcd. for $C_{16}H_{14}N_2O_2S$ (298.37): C, 64.41%; H, 4.73; N, 9.39. Found: C, 64.45; H, 4.79; N, 9.31.

15b: Yield 70%. Mp 235-237 °C. Anal. Calcd. for $C_{17}H_{16}N_2O_2S$ (312.39): C, 65.36%; H, 5.16; N, 8.97. Found: C, 65.40; H, 5.11; N, 8.93.

15c: Yield 75%. Mp 226-228 °C. Anal. Calcd. for $C_{17}H_{16}N_2O_3S$ (328.39): C, 62.18%; H, 4.91; N, 8.53. Found: C, 62.10; H, 4.98; N, 8.48.

15d: Yield 81%. Mp 243-245 °C. Anal. Calcd. for $C_{16}H_{13}ClN_2O_2S$ (332.81): C, 57.74%; H, 3.94; N, 8.42. Found: C, 57.69; H, 3.86; N, 8.50.

3-Aryl-2,4-dioxo-1,2,3,4,5,6,7,8-octahydro[1]benzothieno[2,3-d]pyrimidines 16

(General Procedure)

The procedure is identical with the one described for the synthesis of compounds 15, except that after the addition of 3 the reaction mixture was heated at 100°C for 8 h.

16a: Yield 70%. Mp 278-280 °C. Anal. Calcd. for $C_{16}H_{14}N_2O_2S$ (298.39): C, 64.41%; H, 4.73; N, 9.39. Found: C, 64.48; H, 4.79; N, 9.35.

16b: Yield 74%. Mp 284-286 °C. Anal. Calcd. for $C_{17}H_{16}N_2O_2S$ (312.38): C, 65.36%; H, 5.16; N, 8.97. Found: C, 65.29; H, 5.19; N, 8.94.

16d: Yield 70%. Mp 272-274 °C. Anal. Calcd. for $C_{16}H_{13}ClN_2O_2S$ (332.81): C, 57.74%; H, 3.94; N, 8.42. Found: C, 57.79; H, 3.99; N, 8.46.

ACKNOWLEDGEMENT

We would like to express our gratitude to the Comisión Asesora de Investigación Científica y Técnica (Spain) for its financial support.

REFERENCES

1. A. Albert, Adv. Heterocyclic Chem., 1982, 32, 1
2. J. Garín, E. Meléndez, F.L. Merchán, and T. Tejero, Heterocycles, 1986, 24, 87
3. J. Garín, V. Martínez, J. Mayoral, E. Meléndez, and F. Merchán, Synthesis, 1981, 961
4. F.L. Merchán, Synthesis, 1981, 965
5. R.M. Scrowston, Adv. Heterocyclic Chem., 1981, 29, 171
6. B.G. Hildick and G. Shaw, J. Chem. Soc. (C), 1971, 1610
7. P. Schmidt and J. Druey, Helv. Chim. Acta, 1956, 39, 986
8. O. Meth-Cohn and B. Narine, J. Chem. Res. (S), 1977, 294; (M), 1977, 3262
9. C.S. Pak, I.K. Youn, and Y.S. Lee, Synthesis, 1982, 969
10. M.B. Devani, C.J. Shishoo, U.S. Pathak, S.H. Parikh, G.F. Shah, and A.C. Padhya, J. Pharm. Sci., 1976, 65, 660
11. S.M. Khripak, A.A. Dobosh, and I.V. Smolanka, Khim. Geterotsikl. Soedin., 1973, 567; Chem. Abst., 1973, 79, 32005
12. L. Capuano, M. Welter, and R. Zander, Chem. Ber., 1969, 102, 3698
13. S. Sarangan and S. Somasekhara, J. Indian Chem. Soc., 1976, 53, 426
14. J. Garín, E. Meléndez, F.L. Merchán, and T. Tejero, Synthesis, 1984, 520

Received, 21st January, 1987