

CONVENIENT RUTHENIUM-COMPLEX CATALYSED SYNTHESIS OF ENTEROLACTONE
FROM THE CORRESPONDING DIBENZYLIDENE SUCCINIC ACID MOIETY

Massimo Bambagiotti-Alberti*, Silvia A. Coran, Franco P. Vincieri,
and Pierandrea Lo Nostro¹

Dipartimento di Scienze Farmaceutiche dell'Università
Via Gino Capponi 9, 50121 Firenze, Italy

Abstract - A convenient synthesis of enterolactone is outlined consisting mainly of a ruthenium carbonyl-hydride complex catalysed hydrogenation of bis(3-methoxybenzylidene)succinic acid.


The recent discovery of enterolactone (2,3-bis(3-hydroxybenzyl)-4-butanolide) in human and various animal excretions²⁻⁶ has prompted a renewed interest in the lignans, a well-known class of plant natural products⁷⁻¹⁰ to which the above mentioned compound of animal origin can also be ascribed.

Many plant lignans exhibit a variety of biological activities¹¹⁻¹³ the most important being proven anticancer capabilities¹⁴⁻²³. Consequently, a great wealth of research has been undertaken in the last five years on the role of enterolactone in human metabolism, which has also finally been regarded as a sort of endogenous protective agent connected to certain types of cancer²⁴. Shortage of synthetic enterolactone for extensive biological tests has been, however, a common complaint in almost every paper concerned with this topic. To meet this requirement several syntheses based on multi-step procedures have been promptly proposed²⁵⁻²⁷.

In order to obtain enterolactone (1) we exploited the ability of the soluble ruthenium carbonyl-hydride complex $H_4Ru_4(CO)_8(PBu_3)_4$ to catalyse the gamma-lactone ring formation^{28,29} starting from the succinic acid moiety, obtained via the Stobbe condensation, in one step under usual high pressure catalytic hydrogenation conditions.

Bis(3-methoxybenzylidene)succinic acid (2) (mp 183-185°C from MeOH; ir (nujol): 3200-2200, 1683 cm^{-1} ; ¹H nmr (60 MHz, DMSO-d₆): 7.85(2H, =CH), 7.5-6.8 (8H, Ar-H), 3.70 (6H, OCH₃); ms: M⁺ 354(4), 336(56), 291(19), 264(12), 229(100), 108(65) m/z) was obtained by an improved Stobbe condensation in 50.5% yield (13.1g) by refluxing (2 h) in anhydrous toluene 3-methoxybenzaldehyde (21g, 154mmol), dimethyl succinate (10.7g, 73mmol) and NaH (3.6g, 150mmol, dry powder). A catalytic amount of prepared Ru-catalyst (50mg)³⁰ and 2 (12.5g) were dissolved in toluene and the

solution placed in a high pressure autoclave. After 24 h at 170°C under hydrogen at 270-300 bars, a complete conversion to 2,3-bis(3-methoxybenzyl)-4-butanolide (3) (Found: M^+ 326.1518. $C_{20}H_{22}O_4$ requires M^+ 326.1518) was achieved by the simultaneous occurrence of double-bond saturation and butanolide formation. The reaction course was advantageously followed by mass spectrometry in the "direct electron impact" mode³¹ allowing direct periodic monitoring of the crude reaction mixture (M^+ of 2 = m/z 354, M^+ of 3 = m/z 326). Gas chromatography (fused silica capillary column and PTV injector) and mass spectrometry (EI) revealed 3 as a 4:1 mixture of the trans- and cis-isomers the more abundant of which being the trans-form.



The toluene solution was dried and evaporated, the residue (11.5g) was treated with cold 1% methanolic potassium hydroxide solution (24 h) to obtain a complete conversion to the trans-form³². The exhausted ruthenium-complex catalyst was finally removed by solvent extraction (hexane, MeOH-aq 90%) giving trans-1 dimethylether in 49% overall yield (colorless gum from dry column chromatography (silica gel and CH_2Cl_2); ir (neat): 2838, 1771 cm^{-1} ; 1H nmr (200 MHz, $CDCl_3$): 2.40-2.70 (m, H-2, H-3, H-5a, H-5b), 2.50 (dd, H-6b), 3.06 (dd, H-6a), 3.75 (6H, -OCH₃), 3.86 (dd, H-4b), 4.10 (dd, H-4a), 6.5-7.3 (8H, arom.); ms: M^+ 326(38), 205(15), 191(3), 159(13), 147(27), 135(9), 122(100), 121(60), 117(7), 107(8), 91(31), 77(12)). Demethylation of this dimethylether with boron tribromide²⁵ gave

trans-1 (8.2g) with spectroscopic data in agreement with those reported in the literature^{25,33}.

Further work is currently in progress for the full exploitation of this process to synthesize other lignans having a butanolide ring in their molecule.

ACKNOWLEDGMENTS

This research was supported by the Italian Consiglio Nazionale delle Ricerche within the program "Progetti Finalizzati Chimica Fine e Secondaria, Ca2".

Thanks are due to Profs. M.Bianchi and U.Matteoli, of this University, for helpful discussion and for making hydrogenation apparatus available.

REFERENCES

1. Taken partly from his Tesi di Laurea in Chimica
2. S.R.Stitch, J.K.Toumba, M.B.Groen, C.W.Funke, J.Leemhuis, J.Vink and G.F.Woods, *Nature*, 1980, **287**, 738.
3. K.D.R.Setchell, A.M.Lawson, F.L.Mitchell, H.Adlercreutz, D.N.Kirk and M.Axelsson, *Nature*, 1980, **287**, 740.
4. H.Adlercreutz, T.Fotsis, R.Heikkinen, J.T. Dwyer, B.R.Goldin, S.L. Gorbach, A.M.Lawson and K.D.R.Setchell, *Med.Biol.*, 1981, **59**, 259.
5. S.P.Borriello, K.D.R. Setchell, M.Axelsson and A.M. Lawson, *J. Applied Bacteriology*, 1985, **58**, 37.
6. K.D.R.Setchell, A.M.Lawson, E.Conwaj, N.F.Taylor, D.M.Kirk, G.F.Cooley, R.D.Farrant, S.Wynn and M.Axelsson, *Biochem.J.*, 1981, **447**.
7. C.B.S.Rao, "The Chemistry of Lignans", Andhra University Press, Waltair, 1978.
8. W.M.Hearon and W.S. MacGregor, *Chem. Rev.*, 1955, **55**, 957.
9. R.S.Ward, *Chem. Soc. Rev.*, 1982, **11**, 75.
10. A.Chatterjee, A.Banerji and J.Banerji, *Proc.Indian Acad.Sci.(Chem.Sci.)*, 1984, **93**, 1031.
11. M.Axelsson, J.Sjovall, B.E.Gustafsson and K.D.R.Setchell, *Nature*, 1982, **298**, 659.
12. T.Markanen, M.Markinen, E.Maunuksela and P.Himanen, *Drug Under Exp. Clin. Res.*, 1981, **7**, 711.
13. W.D.MacRae and G.H.Towers, *Phytochemistry*, 1984, **23**, 1207.
14. K.Tomioka and K.Koga, *Tetrahedron Lett.*, 1979, 3315.
15. J.Leiter, V.Dowing, J.L.Hartwell and M.J.Shear, *J.Nat.Cancer Inst.*, 1950, **10**, 1273.
16. P.B.McDoniel and J.R.Cole, *J.Pharm.Sci.*, 1972, **61**, 1992.
17. H.Stahelin, *Eur. J. Cancer*, 1973, **9**, 215.
18. S.J.Torrance, J.J.Hoffman and J.R.Cole, *J.Pharm.Sci.*, 1979, **68**, 664.
19. K.N.Campbell, J.A.Cella and B.K.Campbell, *J.Am.Chem.Soc.*, 1953, **75**, 4681.
20. J.R.Cole, E.Bianchi and E.R.Trumbull, *J.Pharm.Sci.*, 1969, **58**, 175.
21. E.M.Greenspan, J.Leiter and M.J.Shear, *J.Nat. Cancer Inst.*, 1950, **10**, 1295.

- 22.D.A.Cairnes, O.Ekundayo and D.G.I.Kingston, *J.Nat. Prod.*, 1980, **43**, 495.
- 23.L.Bézanger-Beauquesne, *Plantes méd. Phytothér.*, 1982, **16**, 206.
- 24.H.Adlercreutz, R.Heikkinen, M.Woods, T.Fotsis, J.T.Dwyer, B.Goldin and S.L.Gorbach, *Lancet*, 1982, 1295.
- 25.A.Pelter, R.S.Ward, P.Satyanarayana and P.Collins, *J.Chem.Soc.Perkin Trans.I*, 1983, 643.
- 26.G.Cooley, F.D.Farrant, D.N.Kirk and S.Wynn, *Tetrahedron Lett.*, 1981, **22**, 349.
- 27.M.B.Groen and J.Leemhuis, *Tetrahedron Lett.*, 1980, **21**, 5043.
- 28.M.Bianchi, G.Menchi, F.Francalanci, F.Piacenti, U.Matteoli, P.Frediani and C.Botteghi, *J. Organomet. Chem.*, 1980, **188**, 109.
- 29.M.Bambagiotti-Alberti, S.A.Coran, V.Giannellini and F.F.Vincieri, *Atti XV Congresso Nazionale S.C.I.*, Grado, 1984, p.146.
- 30.F.Piacenti, M.Bianchi, P.Frediani and E.Benedetti, *Inorg.Chem.*, 1971, **10**, 2759.
- 31.P.Traldi, U.Vettori and F.Dragoni, *Org.Mass Spectrom.*, 1982, **17**, 587.
- 32.R.D.Haworth and J.R.Atkinson, *J.Chem.Soc.*, 1938, 797.
- 33.J.Harmatha, M.Budesinsky and A.Trka, *Collect.Czech.Chem.Comm.*, 1982, **47**, 644.

Received, 3rd March, 1987