

IWAMIDE: A NEW PHENOLIC AMIDE DERIVED FROM A QUATERNARY
BENZO[c]PHENANTHRIDINE ALKALOID¹⁾


Tsutomu Ishikawa and Hisashi Ishii*

Faculty of Pharmaceutical Sciences, Chiba University,
1-33, Yayoi-cho, Chiba, Japan

The structure of iwamide (I) was established by the fact that decarine (II) was chemically transformed into iwamide (I) using Baeyer-Villiger like oxidation of an immonium group.


In the course of our studies on the chemical constituents of Rutaceous plants, we²⁾ have isolated a new phenolic amide (I) as a minor component of the bark of *Xanthoxylum arnottianum* Maxim. (Japanese name: Iwa-zansho), in 0.00086 % yield. We wish to designate this compound as iwamide (I) and report here its structural establishment due to chemical interconversion of decarine³⁾ (II), a tertiary benzo[c]phenanthridine alkaloid, into iwamide (I) using Baeyer-Villiger like oxidation of an immonium group.

Iwamide (I) is a phenolic amide isolated as colourless prisms, mp 271-273° (from CHCl₃-MeOH). [C₂₀H₁₇O₆N⁴⁾ (M⁺: at m/e




(III) : $\text{R}_1=\text{OCH}_3$, $\text{R}_2=\text{H}$

(IV) : $\text{R}_1=\text{H}$, $\text{R}_2=\text{OCH}_3$



(II)



(V) : $\text{R}_1=\text{OCH}_3$, $\text{R}_2=\text{H}$

(VI) : $\text{R}_1=\text{H}$, $\text{R}_2=\text{OCH}_3$

367)]. It shows the following spectral properties: $\text{ir } \nu_{\text{max}}^{\text{KBr}} \text{ cm}^{-1}$: 3480(OH), 1646(C=O); $\text{nmr}(\text{CDCl}_3 + \text{CD}_3\text{OD}) \delta$: 2.97(3H, s, NCH_3), 3.82(3H, s, OCH_3), 6.05(2H, s, OCH_2O), 6.43(1H, s, $J=9.0$ Hz, aromatic H), 6.66(1H, d, $J=9.0$ Hz, aromatic H), 7.00(1H, s, aromatic H), 7.18(1H, s, aromatic H), 7.27(1H, d, $J=9.0$ Hz, aromatic H), 7.71(1H, d, $J=9.0$ Hz, aromatic H), 8.07(1H, s, CHO).

In the previous paper,⁵⁾ we reported the natural occurrence of a new type of amides, arnottianamide (III) and isoarnottianamide (IV), which were supposed to be formed by Baeyer-Villiger like oxidation of an immonium group of quaternary benzo[c]-phenanthridine alkaloids, chelerythrine (V) and nitidine (VI) respectively. We, furthermore, showed that this type of reaction took place in a test tube.


The presence of a formyl proton at δ 8.07 as a 1H singlet and an N-methyl signal at δ 2.97 as a 3H singlet in the NMR spectrum of iwamide (I) suggested that this amide (I) could be

belonged to the arnottianamide-isoarnottianamide group. Taking into consideration that decarine³⁾ (II) was obtained as a phenolic tertiary base from the same plant,⁵⁾ we could imagine that iwamide (I) was formed by quaternarization of decarine (II) followed by Baeyer-Villiger like oxidation of an immonium group in the plant body.

Treatment of decarine (II) with benzylchloride gave benzyl decarine (VII) as pale yellow needles, mp 218-220° (from benzene-MeOH), in 81.3 % yield. $[C_{26}H_{19}O_4N^4]$; nmr(CDCl₃) δ : 4.12(3H, s, OCH₃), 5.24(2H, s, OCH₂Ph), 6.04(2H, s, OCH₂O), 7.20(1H, s, aromatic H), 7.30-7.60(6H, m, aromatic H), 7.75(1H, d, J=9.0 Hz, aromatic H), 8.21(1H, d, J=9.0 Hz, aromatic H), 8.24(1H, d, J=9.0 Hz, aromatic H), 8.66(1H, s, aromatic H), 9.68(1H, s, aromatic H)].

Since quaternarization of a nitrogen of a pyridine nucleus with an alkyl reagent gave sometimes a mixture of a quaternary base and salt of the starting tertiary base, we attempt to improve this step of the reaction sequence. A solution of benzyl decarine (VII) in hexamethylphosphoric triamide (HMPA) was treated with NaBH₄ and dimethyl sulfate to give N-methyl dihydro-derivative (VIII) as colourless fine needles, mp 183.5-185.5° (from benzene-MeOH), in 98.8 % yield. $[C_{27}H_{23}O_4N^4]$; nmr(CDCl₃) δ : 2.58(3H, s, NCH₃), 3.91(3H, s, OCH₃), 4.29(2H, s, ArCH₂N<), 5.15(2H, s, OCH₂Ph), 6.00(2H, s, OCH₂O), 6.95(1H, d, J=9.0 Hz, aromatic H), 7.08(1H, s, aromatic H), 7.30-7.55(7H, m, aromatic H), 7.65(1H, s, aromatic H), 7.67(1H, d, J=9.0 Hz, aromatic H)].

Oxidation of N-methyl dihydro-derivative (VIII) with DDQ



gave a desired quaternary salt which was obtained as sulfate (IX), orange needles, mp $260-265^\circ$ (from MeOH), in 50.4 % yield. [nmr($\text{CF}_3\text{CO}_2\text{H}$) δ : 4.28(3H, s, OCH_3), 4.93(3H, s, NCH_3), 5.29(2H, s, OCH_2Ph), 6.06(2H, s, OCH_2O), 7.18-7.36(6H, m, aromatic H), 7.84(1H, s, aromatic H), 7.96(2H, d, $J=8.0$ Hz, aromatic H), 8.33(2H, d, $J=8.0$ Hz, aromatic H), 9.51(1H, s, $\text{ArCH}=\text{N}^+$)].

Treatment of the sulfate (IX) in HMPA with m -chloroperben-

zoic acid afforded benzyl iwamide (X) as colourless prisms, mp 249-252° (from CHCl_3 -MeOH), in 77.9 % yield. $[\text{C}_{27}\text{H}_{23}\text{O}_6\text{N}^4] (\text{M}^+)$: at m/e 457); ir $\nu_{\text{max}}^{\text{KBr}}$ 1675 (C=O); nmr (CDCl_3) δ : 2.99 (3H, s, NCH_3), 3.97 (3H, s, OCH_3), 5.12 (2H, s, OCH_2Ph), 5.99 (1H, s, OH), 6.05 (2H, s, OCH_2O), 6.58 (1H, d, $J=8.5$ Hz, aromatic H), 6.78 (1H, d, $J=8.5$ Hz, aromatic H), 7.08 (1H, s, aromatic H), 7.19 (1H, s, aromatic H), 7.36-7.53 (6H, m, aromatic H), 7.73 (1H, d, $J=8.5$ Hz, aromatic H), 8.17 (1H, s, CHO)].

A benzyl-protecting group was removed via hydrogenolysis using 5 % palladium on charcol to give colourless prisms, mp 267-268.5°, in 97.6 % yield. This was identified with a sample of iwamide (I) which was obtained from the natural source. This result was enough to establish the structure of iwamide to be shown by the formula (I).

REFERENCES

- 1 This paper constitutes Part XXXII of Chemical Constituents of Rutaceous Plants. [Part XXXI: H. Ishii, T. Ishikawa, S.-T. Lu, and I.-S. Chen, Yakugaku Zasshi, submitted for publication].
- 2 H. Ishii, T. Ishikawa, and J. Haginiwa, Yakugaku Zasshi, in preparation.
- 3 J. Vaquett, J. L. Pousset, R. R. Paris, and A. Cavé, Phytochemistry, 1974, 13, 1257.
- 4 The compound gave satisfactory elemental analysis for the formula given.
- 5 H. Ishii, T. Ishikawa, S.-T. Lu, and I.-S. Chen, Tetrahedron letters, 1976, 1203.

Received, 2nd August, 1976