

Reactions of Stable Ylidic 1-Thianaphthalene with Electrophiles.¹

A New Ring Expansion Reaction

Mikio Hori,* Tadashi Kataoka, Hiroshi Shimizu, and Harumi Aoki
Gifu College of Pharmacy, Mitahora, Gifu 502, Japan

Reactions of 2-cyano-1-methyl-4-phenyl-1-thianaphthalene (I) with electrophiles, such as tetracyanoethylene, maleic anhydride, and 4-phenyl-1-thianaphthylum perchlorate gave the dimeric compound (II) of I, but not any cycloaddition products. However, the treatment of I and dimethyl acetylenedicarboxylate afforded a new ring expansion product (III) having sulphur-containing nine-membered ring.

Recently, much attention has been focussed on thiabenzenes in organic chemistry.^{2a,3a} Some stable ylidic 1,2-disubstituted 2-thianaphthalenes,^{2b,3b} and 9,10-disubstituted 10-thiananthracenes^{2c,2d} have been reported. However, report on the title ring system has not been done.

In this communication, we wish to describe the reactions of a typical 1-thianaphthalene derivative (I)¹ with electrophiles, such as tetracyanoethylene (TCNE), maleic anhydride, 4-phenyl-1-thianaphthylum perchlorate, and dimethyl acetylenedicarboxylate (DMAD). And also, especially, discussion will be made on a new ring expansion reaction using I with DMAD.

Stirring I with TCNE in benzene at room temperature under nitrogen stream for 12 hr gave dimeric compound (II) of I in 9.3 % yield, the recovered TCNE and black resins of undetermined structure. The physical data of II are as follows : m.p. 279-281° ($C_{34}H_{26}N_2S_2$) as orange needles (benzene-pet.ether) : mass (m/e) : 526 (M^+) : ir (KBr) ν max cm^{-1} ; 2225 (CN) ; nmr (C_6D_6) ; δ 1.90 (3H, s, S-CH₃), 6.85-7.50 (10H, m, olefinic and aromatic H). Similarly, II was also obtained by the treatment of I with maleic anhydride in benzene for 30 hr in 10 % yield and with 4-phenyl-1-thianaphthylum perchlorate in CH_2Cl_2 for 6 hr in 30 % yield besides the recovered 4-phenyl-1-thianaphthylum perchlorate in 77.3 %, but any isolable cycloaddition products were not obtained.




Chart 1

From the above data, the authors propose the mechanism for the reaction of I with these electrophiles as shown in Chart 1.

On the other hand, the treatment of I with 2 eq DMAD in a manner similar to the case of TCNE, yielded an interesting ring expansion product (III) : m.p. 142-149° ($C_{23}H_{19}O_4NS$) as colorless prisms (ether) : mass (m/e) 405 (M^+) : ir (KBr) ν max cm^{-1} ; 2220 (CN), 1740 and 1710 (CO) : nmr ($CDCl_3$) δ 2.98 (1H, d,d, $J_{ab} = 15$ Hz, $J_{ac} = 9$ Hz, H_a), 3.68 (3H, s, CO_2CH_3), 3.71 (1H, broad d, $J_{ab} = 15$ Hz, $J_{bc} = 0$ Hz, H_b), 3.72 (3H, s, CO_2CH_3), 4.20 (1H, broad d, $J_{ac} = 9$ Hz, $J_{bc} = 0$ Hz, H_c), 6.90 (1H, s, olefinic H), and 7.20-8.00 (9H, m, aromatic H). Subsequently, III was successively oxidized with m-chloroperbenzoic acid in CH_2Cl_2 over night at 0° to give sulfone derivative (IV) in 77.8 % yield : m.p. 192-193° ($C_{23}H_{19}O_6NS$) as colorless prisms (ether) : ir (KBr) ν max cm^{-1} ; 2225 (CN), 1740 (CO), 1318 and 1058 (SO_2) : nmr ($CDCl_3$) δ 3.45 (1H, d,d, $J_{ab} = 17$ Hz, $J_{ac} = 10$ Hz, H_a), 3.78 (6H, s, CO_2CH_3), 4.28 (1H, broad d, $J_{ab} = 17$ Hz, $J_{bc} = 0$ Hz, H_b), 4.50 (1H, broad d, $J_{ac} = 10$ Hz, $J_{bc} = 0$ Hz, H_c), 6.83 (1H, s, olefinic H), 7.30-8.30 (9H, m, aromatic H). However, it is difficult to determine the stereochemical assignments of cis and trans isomers of III. Quite recently, Vedejs et al.⁴ reported on synthesis of eight-membered ring compound having a cis double bond as major product starting from sulphur ylide containing five-membered ring. Thus, III can be assigned to the cis form by the stereochemical requirement on the transition state.

The formation of III is presumed to proceed through the five-centered transition state as shown in Chart 2.




Chart 2

Finally, concerning with the intermediate A in Chart 1 :

When E^- () is rather stable, the reaction might proceed to give the dimeric product. When E^- () is more basic than the above described species, the intramolecular deprotonation might occur to give the authors' intermediate ylide in Chart 2.

Further related studies directed toward the applications to synthetic and theoretical organic chemistry are also in progress.

ACKNOWLEDGMENT. The authors deeply wishes to thank the Ministry of Education for their kind financial help for this research work.

REFERENCES.

- 1 A part of this work was presented at the 96th Annual Meeting of the Pharmaceutical Society of Japan, Nagoya, April, 1976, Abstracts of Paper II p. 42.
- 2 a) M. Hori, and H. Shimizu, FARUMASHIA, 12, 468 (1976) ; b) M. Hori, T. Kataoka, H. Shimizu, K. Narita, S. Ohno, and H. Aoki, Chem. Lett., 1974, 1101 ; c) M. Hori, T. Kataoka, H. Shimizu, S. Ohno, K. Narita, and H. Aoki, Fifth International Congress of Heterocyclic Chemistry, Ljubliana (Yugoslavia), July, 1975, Abstracts of Papers p. 299 ; d) M. Hori, T. Kataoka, H. Shimizu, S. Ohno, M. Okitsu, H. Matsushita, and T. Miyasako, VIIth International Symposium on Organic Sulphur Chemistry, Hamburg (Federal Republic of Germany), July, 1976, Abstracts of Papers p. 172.
- 3 a) G. H. Senkler Jr., B. E. Maryanoff, J. Stackhouse, J. D. Andose, and K. Mislow, " Organic Sulphur Chemistry ", ed. by C. J. M. Stirling, Butterworths & Co (Publishers) London and Boston, pp. 157-179 ; b) G.H. Senkler Jr., J. Stackhouse, B. E. Maryanoff, and K. Mislow, J. Amer. Chem. Soc., 96, 5650 (1974) ; B. E. Maryanoff, J. Stackhouse, G. H. Senkler Jr.,

and K. Mislow, ibid. 97, 2718 (1975).

4 E. Vedejs, and J. P. Hagen, ibid. 97, 6875 (1975).

Received, 19th August, 1976