

OXIDATIVE RING CLEAVAGE OF CORALYNE ALKALOID

Yoshikazu Kondo*, Hideo Inoue, and Jiro Imai


Pharmaceutical Institute, Tohoku University, Aobayama, Sendai, Japan

Oxygenation of coralyne alkaloid, a nonenzymic model of the oxygenase, has been investigated. A solution of dihydrocoralyne 1 or 13-hydroxycoralyinium betaine 2 in the presence of sodium alkoxide was allowed to contact with air under illumination of visible light to give 6,7-dimethoxyisoquinolone-1 3 and the corresponding 3-methyl-3-alkoxy-5,6-dimethoxyphthalide 4 or 5. When the same autoxidation was carried out in the presence of borohydride anion, 3 and 3-methyl-5,6-dimethoxyphthalide 6 were obtained along with 13-hydroxycoralydine. Oxygenation of 2 in the presence of cuprous chloride led to 2'-acetylpapaveraldine 11. On the other hand, irradiation of 2 in chloroform in the presence of Rose Bengal gave a 1:1 mixture of 11 and 2'-carbomethoxy-papaveraldine 12.

The mechanisms were also discussed.

The existence of the isoquinolone alkaloids together with the major berberine alkaloids in plants has been reported, and the possibility was recognized that the isoquinolone might be a result of enzymic oxidation of these major alkaloids¹. However, the mode of oxidation of the berberine alkaloids in plants is little known at the present time. In this communication we report the oxygenation of coralyne alkaloid, a nonenzymic model of the dioxygenase², in which the products

are dependent on a different excitation means of oxygen.




Scheme 1

A solution of dihydrocoralyne³ 1 in hot methanol was kept for 2 hr at pH 8 and to the solution subsequently was added sodium methoxide. The solution was allowed to stand under illumination of a photoflood lamp to give 6,7-dimethoxyisoquinolone-1 3, C₁₁H₁₁O₃N, mp 244-245° (lit.⁴, mp 237°), and 3-methyl-3,5,6-trimethoxyphthalide⁵ 4, C₁₂H₁₄O₄, mp 150-152°, in moderate yields. 3: IR (KBr) cm⁻¹: 3450 (>NH), 1658 and 1632 (amido carbonyl). UV (EtOH) nm (log ε): 246 (4.26), 253 (sh, 4.00), 268 (3.45), 278 (3.48), 290 (3.51), 310 (3.25), 322 (3.35), 335 (3.24). NMR (CDCl₃) ppm: 1.75 (1H, br.s, >NH), 4.01 (3H, s, -OCH₃), 4.04 (3H, s, -OCH₃), 6.46 (1H, d, J=8 Hz, Ar-CH=CH-N<), 6.88 (1H, s, Ar-H), 7.10 (1H, br. d, J=8Hz, Ar-CH=CH-N<), 7.75 (1H, s, Ar-H). MS m/e: 205 (M⁺). 4: IR (KBr) cm⁻¹: 1745 (lactone carbonyl). UV (EtOH) nm (log ε): 221 (4.14), 257 (3.71), 293 (3.60), 300 (sh, 3.57). NMR (CDCl₃) ppm: 1.80 (3H, s, -CH₃), 3.07 (3H, s, -OCH₃), 3.93 (3H, s, Ar-OCH₃), 6.80 (1H, s, Ar-H), 7.20 (1H, s, Ar-H). MS m/e: 238 (M⁺). Similar oxidation accompanied by cleavage of the hetero-ring was observed by the addition of caustic alkali in aqueous solution, but the corresponding phthalide could not be isolated.

It is considered that the oxidative cleavage of 1 proceeds *via* the initial formation of the phenolbetaine 2. Our previous work has demonstrated that 1 is oxidized to 2 in quantitative yield under the physiological conditions³. A solution of 2 in methanol containing sodium methoxide was allowed to contact with air under illumination of visible light to give 3 and 4 in approximately 50% and 30% yields, respectively (Scheme 1). When sodium ethoxide-ethanol solution was used in the place of sodium methoxide, the autooxidation of 2 gave 3 and 3-ethoxy-3-methyl-5,6-dimethoxyphthalide 5, C₁₃H₁₆O₄, oil, in approximately 45% and 30% yields, respectively. 5: IR (CHCl₃) cm⁻¹: 1753 (lactone carbonyl). UV (MeOH) nm (log ε): 223 (4.14), 261 (3.62), 293 (3.50), 300 (sh, 3.48). NMR (CDCl₃) ppm: 1.15 (3H, t, J=8 Hz, -OCH₂CH₃), 1.82 (3H, s, -CH₃), 3.10 (1H, dq, J_{gem}=9 Hz, J_{vic}=8 Hz, -OCHH-CH₃), 3.44 (1H, dq, J_{gem}=9 Hz, J_{vic}=8 Hz, -OCHH-CH₃), 6.90 (1H, s, Ar-H), 7.30 (1H, s, Ar-H). MS m/e: 250 (M⁺).


The autoxidative degradation of 2 is rationalized by addition of singlet oxygen to form a peroxide intermediate 7, which could cleave *via* 1,2-dioxetane⁶ 8 and hydrolyze by concomitant attack of the nucleophile to 3 and the corresponding phthalide (Scheme 2). The mechanism is further supported by the fact that the oxygenation of 2 was carried out in methanol in the presence of sodium borohydride to give 3 and 3-methyl-5,6-dimethoxyphthalide 6, C₁₁H₁₂O₄, mp 101-102° (lit.⁷, 100-101°), together with 13-hydroxycoralydine, C₂₂H₂₇O₄N, mp 185-187°, in 26%, 16%, and 4% yields, respectively. 6: IR (KBr) cm⁻¹: 1730 (lactone carbonyl). UV (EtOH) nm (log ε): 221 (4.11), 257 (3.67), 293 (3.58), 300 (sh, 3.52). NMR (CDCl₃) ppm: 1.60 (3H, d, J=7 Hz, >CH-CH₃), 3.92 (3H, s, -OCH₃), 3.96 (3H, s, -OCH₃), 5.43 (1H, q, J=7 Hz, Ar-CH-O), 6.98 (1H, s, Ar-H), 7.22 (1H, s, Ar-H). In the final step of the mechanism, the borohydride anion attacks to the amido-lactone intermediate 10 to give the phthalide 6. When the same reaction was carried out in the solution previously deactivated by acetone, the



Scheme 2

products, 3 and 4, were exactly analogous to the autoxidation in sodium methoxide-methanol solution.

Of interest the oxidation of 2 in DMF in the presence of cuprous chloride which catalyzed the activation of molecular oxygen^{8,9}, yielded 2'-acetylpapaveraldine³ 11 in quantitative yield. Since single oxygen is inserted in 2, the most likely intermediate, the peroxide 13, formed *via* rearrangement of 7, should exist before final ring opening (Scheme 3). When irradiation of 2 in chloroform in the presence of Rose Bengal was performed, a 1:1 mixture of 11 and 2'-carbomethoxypapaveraldine 12, C₂₂H₂₁O₇N, mp 210-212°, was obtained. 12: IR (KBr) cm⁻¹: 1705 (ester carbonyl), 1675 (Ar-CO-Ar), 1438 and 1355 (carbomethoxy). UV (EtOH) nm (log ε): 235 (4.43), 260 (sh, 4.20), 298 (3.70), 343 (3.68). NMR (CDCl₃) ppm: 3.20 (3H, s, -COOCH₃), 4.00 (6H, s, -OCH₃), 4.07 (3H, s, -OCH₃), 4.13 (3H, s, -OCH₃), 7.08 (1H, s, Ar-H), 7.18 (1H, s, Ar-H), 7.20 (1H, s, Ar-H).



8.64 (1H, s, Ar-H), 7.55 (1H, d, $J=7$ Hz, $-\text{CH}=\text{CH}-\text{N}<$), 8.54 (1H, d, $J=7$ Hz, $-\text{CH}=\text{CH}-\text{N}<$). MS m/e : 411 (M^+), 396, 380, 368, 352. The mechanism of formation of 12 involves simultaneous cyclization of 13 to a three-membered cyclic peroxide 14 (Scheme 3).

The analogous oxidative ring cleavage has been extended to the berberine alkaloids¹⁰.

ACKNOWLEDGEMENT The authors wish to thank Professor A. Nishinaga, Faculty of Engineering, Kyoto University, for valuable discussions.

REFERENCES AND NOTES

- 1 R. W. Duskoch, P. L. Schiff, Jr., and J. L. Beal, *Tetrahedron*, 1969, 25, 469.
- 2 Reviews: V. Urlich, *Angew. Chem.*, 1972, 84, 689; A. Nishinaga, *J. Syn. Org. Chem. Japan*, 1976, 34, 856 and references cited therein.
- 3 J. Imai and Y. Kondo, *Heterocycles*, 1976, 5, 153.
- 4 A. M. Deryckere and F. G. F. Eloy, *Ger. Offen.*, 1,933,870 (*CA*, 1970, 72, P90317x).
- 5 All new compounds gave satisfactory elemental analysis.
- 6 K. Orito, R. H. Manske, and R. Rodrigo, *J. Am. Chem. Soc.*, 1974, 96, 1944.
- 7 E. Maekawa and Y. Sumimoto, *Bull. Chem. Soc. Japan*, 1960, 33, 941.
- 8 J. Tsuji and H. Takayanagi, *J. Am. Chem. Soc.*, 1974, 96, 7349; J. Tsuji, H. Takayanagi, and I. Sakai, *Tetrahedron Letters*, 1975, 1245; J. Tsuji and H. Takayanagi, *Tetrahedron Letters*, 1976, 1365.
- 9 Recently, Rogić et al. demonstrated that the oxidative cleavage of catechol (phenol) to *cis*, *cis*-muconate derivative was catalysed by cupric ion in pyridine containing methanol and water under the anaerobic conditions. They claimed that one of the two new introduced oxygen atoms originated from solvent water. M. M. Rogić, T. R. Demmin, and W. B. Hammond, *J. Am. Chem. Soc.*, 1976, 98, 7441.
- 10 To be published.

Received, 1st February, 1977