

[Dedicated to Professor Robert B. Woodward on the occasion of his 60th birthday.]

THE BIRCH REDUCTION OF QUERCETIN METHYL ETHERS

James G. Sweeny, Terence Radford, and Guillermo A. Iacobucci*


Corporate Research and Development Department, The Coca-Cola Company, P. O. Drawer 1734, Atlanta, GA 30301, U.S.A.

The sodium-ammonia reduction of 3,3',4',5,7-pentamethylquercetin (I) afforded the dihydrochalcone (II) in 54% yield. In contrast, the reduction of 3',4',5,7-tetramethylquercetin (III) yielded the α -hydroxy-dihydrochalcone (IV) (64% yield) as major product. The reactions are believed to proceed through the intermediary formation of 2,3-trans-dihydroquercetins.

Regiospecific 1,2-reductions of flavonols, either at the carbonyl or the $\Delta^{2,3}$ -double bond, are known to proceed poorly as a result of electronic and steric interactions¹. We would like to communicate that the heterocyclic ring of 3,3',4',5,7-pentamethylquercetin (I) is smoothly hydrogenolyzed by sodium in liquid ammonia to yield 2'-hydroxy-3,4,4',6'-tetramethoxydihydrochalcone (II), with the loss of the methoxy group originally attached to the flavone C₃-position.


To a suspension of 1 g (2.7 mmol) (I) in liquid ammonia (40 ml), sodium (400 mg, 17.4 mmol) was added in three portions over 10 minutes. After stirring for one hour the mixture was quenched with ammonium chloride; the ammonia was evaporated, and the residue was partitioned between 1% acetic acid and chloroform. The organic layer afforded a brown oil, which was crystallized from methanol to yield 518 mg (54%) of II. m.p. 124-5° (lit.² 125-6°). NMR (CDCl₃): 4H m at δ 2.8-3.4 ppm (2 CH₂), 12H s at δ 3.8 ppm (4OCH₃), 2H q at

δ 6.0 ppm (C_3H and C_5H) and 3H s at δ 6.8 ppm (C_2H , C_5H and C_6H). MS (m/e, relative intensity): M^+ 346 (57), 181 (100), 164 (85) and 151 (42).




(I) R = OCH_3

(III) R = OH



(II) R = H


(IV) R = OH




(V) R = OCH_3

(VI) R = OH


When 3',4',5,7-tetramethylquercetin (III)³ was reduced with excess sodium as above, however, the corresponding 2', α -dihydroxy-3,4,4',6'-tetramethoxydihydrochalcone (IV) was isolated from preparative TLC on silica gel (eluant: 50% EtOAc-hexane) in 64% yield, m.p. 118-20° (MeOH). (Calc. for $C_{19}H_{22}O_7$: C, 62.97; H, 6.12. Found: C, 62.75; H, 6.16.) The NMR ($CDCl_3$) of IV showed a 2H multiplet at δ 2.75-3.05 ppm for the benzyl CH_2 and a broad 1H quartet at δ 5.3-5.5 ppm for the proton α - to the keto group. The mass spectrum included fragments at m/e 344 (M^+-H_2O), 211, 181, and 151, which were assigned as follows:




m/e 344



m/e 211



m/e 181



m/e 151

Furthermore, the Birch reduction of the corresponding 2,3-trans-dihydroquercetins V and VI afforded, as expected, the dihydrochalcones II and IV in 31 and 23% yield, respectively. From these results, one plausible mechanism would involve first the two-electron reduction of the flavonol to the 2,3-dihydro derivative, followed by base catalyzed opening to the α -substituted chalcone and further reduction to the final products.

α -Hydroxydihydrochalcones have been found in nature quite recently.^{4,5} The Birch reduction of readily available flavonols seems a convenient route for the synthesis of specific members of this group of natural products.

REFERENCES

- 1 T. A. Geissman (Ed.), "The Chemistry of Flavonoid Compounds", MacMillan, New York, 1962.
- 2 K. Freudenberg and E. Cohn, Chem. Ber., 1923, 56B, 2127.
- 3 G. Janzsó, F. Kállay, and I. Koczor, Tetrahedron, 1966, 22, 2909.
- 4 Y. N. Shukla, J. S. Tandon, and M. M. Dhar, Ind. J. Chem., 1973, 11, 720.
- 5 D. Bhakuni, M. Bittner, M. Silva, and P. G. Sammes, Phytochemistry, 1973, 12, 2777.

Received, 15th March, 1977