

STUDIES ON POTENTIAL ANTITUMOR AGENTS (II).

Thiosemicarbazones of p-Bromophenyl- and
o-Chlorophenylpyridine-2-carboxaldehydes

Lian-Fang Lin , Shwu-Jiuan Lee , and Chao-Tung Chen*

Institute of Chemistry, Academia Sinica, Nankang, Taipei, Taiwan 115
Republic of China

Thiosemicarbazones of eight p-bromophenyl- and o-chlorophenylpyridine-2-carboxaldehydes have been synthesized. Tumor-inhibitory potency and host toxicity of these compounds were assessed in mice bearing Sarcoma 180 ascites cells.

4-(m-Aminophenyl)pyridine-2-carboxaldehyde thiosemicarbazone was reported to have potential antitumor activity on mice bearing Sarcoma 180 ascites cells.¹ A previous communication from This Laboratory² described synthesis of thiosemicarbazones of p-chlorophenyl- and m-chlorophenylpyridine-2-carboxaldehydes. The present study was undertaken to prepare further thiosemicarbazones of p-bromophenyl- and o-chlorophenylpyridine-2-carboxaldehydes for biological evaluation.

p-Bromophenyl-2-picoline were synthesized by a procedure similar to the arylation of 2-picoline with diazotized p-chloroaniline² (scheme 1). p-Bromoaniline (103.2 g) was diazotized at 0° and the resulting solution was added dropwise to 2-picoline (300 ml) at 40°. The reaction mixture was kept at 80° for 30 minutes and worked up as usual² to obtain 38 g of a semisolid, which was chromatographed on silica gel and on neutral alumina. Four isomeric p-bromophenyl-2-picoline (1-4) were obtained pure. The isomers were differentiated by nmr studies

(Table I). These arylpicolines (1-4) were oxidized with selenium dioxide in refluxing dioxane to p-bromophenylpyridine-2-carboxaldehydes (5-8), which were converted to thiosemicarbazones (9-12). These compounds are listed in Table II.

o-Chlorophenyl-2-picolines resulted from the reaction of 2-picoline and diazotized o-chloroaniline in a similar fashion except that the reaction mixture was maintained at pH 4-5. This reaction gave, after chromatography, 3-, 4-, 6- and a mixture of 3- and 5-(o-chlorophenyl)-2-picolines (13-16). Selenium dioxide oxidation of these methyl compounds gave corresponding aldehydes (17-20), the mixture of 3- and 5- isomers being separated by chromatography. The aldehydes were converted to thiosemicarbazones (21-24). These results are listed in Table III.

Biological evaluation of these thiosemicarbazones on mice bearing Sarcoma 180 ascites cells showed 4-(o-chlorophenyl)-pyridine-2-carboxaldehyde thiosemicarbazone to be the best.

Scheme I.





Table I. NMR Data*

Compd	Position of aryl group substitution	2-Methyl protons (3H, s)	Protons on phenyl and pyridine rings	H-6 on pyridine ring (1H, dd)	Proton of aldehyde CHO (1H, s)
1	3	2.45	7.03-7.73 (6H, m)	8.53 J=5,2 Hz	
2	4	2.53	7.13-7.70 (6H, m)	8.53 J=5, 1 Hz	
3	5	2.53	7.06-7.79 (6H, m)	8.76 J=2,1 Hz	
4	6	2.56	7.03-8.09 (7H, m)		
5	3		7.00-7.83 (6H, m)	8.82 J=5,2 Hz	10.05
6	4		7.20-8.13 (6H, m)	8.75 J=5,1 Hz	10.05
7	5		7.30-8.23 (6H, m)	9.00 J=2,1 Hz	10.10
8	6		7.13-8.03 (7H, m)		10.03
13	3	2.34	7.10-7.90 (6H, m)	8.65 J=5,2 Hz	
14	4	2.63	7.13-7.73 (6H, m)	8.71 J=5,1 Hz	
16	6	2.61	7.10-7.90 (7H, m)		
17	3		7.03-7.83 (6H, m)	8.78 J=5,2 Hz	9.91
18	4		7.06-8.06 (6H, m)	8.77 J=5,1 Hz	10.05
19	5		7.13-8.00 (6H, m)	8.73 J=2,1 Hz	9.98
20	6		7.10-8.00 (7H, m)		10.04

*NMR Spectra were recorded on a JEOL-C-60-HL High Resolution NMR Instrument. The chemical shifts are reported in ppm downfield from internal TMS. Compounds 1, 2, 3, 4, 19 were dissolved in CCl_4 , all others in CDCl_3 .

Table II. Compounds of p-Bromophenyl Series




Compd ^a	R	Position of p-Bromophenyl Substitution	Mp (Bp) °C	% Yield	Formula	Mass Spectra (M ⁺)
1	CH ₃	3	50.5-51.5	13.7 ^b	C ₁₂ H ₁₀ BrN	247
2	CH ₃	4	75-76.2	20.7 ^b	C ₁₂ H ₁₀ BrN	247
3	CH ₃	5	100-101	6.2 ^b	C ₁₂ H ₁₀ BrN	247
4	CH ₃	6	75	59.4 ^b	C ₁₂ H ₁₀ BrN	247
5	CHO	3	87-88	78(196) ^c	C ₁₂ H ₈ BrNO	261
6	CHO	4	101-102	47(286) ^c	C ₁₂ H ₈ BrNO	261
7	CHO	5	137-138.5	52.6(135) ^c	C ₁₂ H ₈ BrNO	261
8	CHO	6	84-85.5	72(297) ^c	C ₁₂ H ₈ BrNO	261
9	CH=NNHCSNH ₂	3	218.5-220 (dec)	87.3	C ₁₃ H ₁₁ BrN ₄ S	
10	CH=NNHCSNH ₂	4	217.5-220 (dec)	81	C ₁₃ H ₁₁ BrN ₄ S	
11	CH=NNHCSNH ₂	5	235-237 (dec)	77	C ₁₃ H ₁₁ BrN ₄ S	
12	CH=NNHCSNH ₂	6	217-219 (dec)	79	C ₁₃ H ₁₁ BrN ₄ S	

a. All compounds are new.

b. Yields are based on the recovery of various isomers from a mixture obtained in 7.0% yield from the arylation of 2-picoline with p-bromoaniline.

c. Reflux time (hours) in selenium dioxide oxidation of the 2-picolines.

Table III. Compounds of o-Chlorophenyl Series



Compd ^a	R	Position of o-Chlorophenyl Substitution	Mp (Bp) °C	% Yield	Formula	Mass Spectra (M ⁺)
13	CH ₃	3	>300(Bp)	35.5 ^b	C ₁₂ H ₁₀ ClN	203
14	CH ₃	4	37-38.5	9.6 ^b	C ₁₂ H ₁₀ ClN	203
15	CH ₃	3,5 ^d	>300(Bp)	6.1 ^b	C ₁₂ H ₁₀ ClN	203
16	CH ₃	6	>300(Bp)	48.8 ^b	C ₁₂ H ₁₀ ClN	203
17	CHO	3	81-82	70.1(103) ^c	C ₁₂ H ₈ ClNO	217
18	CHO	4	68-70	43.4(110) ^c	C ₁₂ H ₈ ClNO	217
19	CHO	5	68-69.5	57.5(102) ^c	C ₁₂ H ₈ ClNO	217
20	CHO	6	86-86.5	78(147) ^c	C ₁₂ H ₈ ClNO	217
21	CH=NNHCSNH ₂	3	204-205 (dec)	88	C ₁₃ H ₁₁ ClN ₄ S	
22	CH=NNHCSNH ₂	4	207-208.5 (dec)	86	C ₁₃ H ₁₁ ClN ₄ S	
23	CH=NNHCSNH ₂	5	204-205 (dec)	91	C ₁₃ H ₁₁ ClN ₄ S	
24	CH=NNHCSNH ₂	6	187-188.5 (dec)	85	C ₁₃ H ₁₁ ClN ₄ S	

a. All compounds are new.

b. Yields are based on the recovery of various isomers from a mixture obtained in 13.5% yield from the arylation of 2-picoline with o-chloroaniline.

c. Reflux time (hours) in selenium dioxide oxidation of the 2-picolines.

d. Mixture of 3- and 5- isomers.

REFERENCES

1. K. C. Agrawal, A. J. Lin, B. A. Booth, J. R. Wheaton, and A. C. Sartorell, J. Med. Chem., 17, 631 (1974).
2. C. T. Chen, S. F. Chen, and S. J. Lee, Heterocycles, 5, 239 (1976); Bull. Inst. Chem. Academia Sinica, No. 24, p. 53, 1977.

ACKNOWLEDGEMENT

We thank National Science Council for a fund through Chemistry Research Center, National Taiwan University.

Received, 23rd July, 1977