

ISOLATION OF AN INSECT ANTIFEEDANT N-METHYLFLINDERSINE
AND SEVERAL BENZ[C]PHENANTHRIDINE ALKALOIDS FROM
EAST AFRICAN PLANTS; A COMMENT ON CHELERYTHRINE

Frances Y. Chou, Kurt Hostettmann, Isao Kubo, Koji Nakanishi*
Department of Chemistry, Columbia University, New York, N. Y. 10027

and

Makoto Taniguchi
Department of Biology, Faculty of Science, Osaka City University,
Sumiyoshi-ku, Osaka 558, Japan

Several benzphenanthridine type alkaloids have been isolated by following insect antifeedant assays for fractionation. There is a possibility that the well-known chelerythrine could be an artefact, at least in the present case.

In continuation with our studies on insect antifeedants^{1,2} and medicinally active compounds contained in plants, we have investigated the root barks of Fagara chalybea³, F. holstii (Rutaceae) and Xylocarpus granatum (Meliaceae)^{16,4}, all of which taste bitter and are used for medicinal treatment in East Africa⁵.

The antifeedant bioassays of the extracts were carried out by the leaf-disk method^{1f,1g,6} employing: (i) African armyworms Spodoptera exempta (monophagous) and S. littoralis (polyphagous) on leaves of Zea mays and Ricinus communis respectively, and (ii) the Mexican bean beetle, Epilachna varivestis on leaves of Bountiful stringbeans. This led to the isolation and characterization of N-methylflindersine 1⁷ as the principle responsible for insect antifeedant activity (Table 1). Thus it was found that Zea mays leaves which were dipped in 100 ppm acetone solution of 1 for 2 seconds⁸ inhibited the feeding of S. exempta as compared to untreated control leaves; this corresponds to a concentration of ca. 1 $\mu\text{g}/\text{cm}^2$.

N-Methylflindersine 1 was also found to be responsible for the antimicrobial activity exhibited by the original 60% aqueous methanol extracts of the root barks. Namely, it was active against the yeast Candida albicans at 50 ppm (minimum inhibitory concentration, MIC) and the fungi Rhizopus delemay at 100 ppm; furthermore, when tested against Trichophyton mentagrophytes it was twice as active as the antibiotic nystatin⁹. However, 1 was only weakly active against gram positive bacteria, e.g., Staphylococcus aureus, Diplococcus pneumoniae and Protens vulgaris, and inactive against gram negative bacteria. In the cytotoxicity test with KB tissue culture, the ED₅₀ was 24 µg/ml¹⁰, and according to a modified method¹¹ of the Ames test¹² employing strain TA-100 of Salmonella typhimurium, 400 revertants per plate could be counted at a concentration of 250 µg/plate in the presence of S-9 mix¹³.

During the process of isolation of the active factor, several other known alkaloids of the benz[c]phenanthridine series were also characterized. As shown in Table 1, and Fig.1 these were dihydrochelerythrine¹⁴, chelerythrine^{15,16}, arnottianamide¹⁷, acetonvldihydrochelerythrine¹⁸, and nitidine^{16,19}.

It has been reported that chelerythrine and nitidine both exhibit strong antitumor activity but are too toxic for human use²⁰.


Liquid chromatography was extensively used in all isolations. In the case of F. chalybea constituents, preparative liquid chromatography was applied directly for the first time to the crude hexane extract³. Upon standing, a precipitate of crystalline dihydrochelerythrine 2 was obtained in a yield of about 0.1%. Prep LC of the residual 3.5 g (corresponding to ca. 150 g of the dried bark) afforded the following compounds in a total working period of 2 hours: 1 g of a mixture of hydrocarbons, 850 mg of germacrone, 120 mg of a further amount of dihydrochelerythrine 2, 180 mg of a mixture of several compounds, 40 mg of a terpene and 125 mg of N-methylflindersine³. In contrast, conventional open column chromatography required 2 weeks using the same amount of the hexane extract, moreover, the unstable dihydrochelerythrine was not isolated by this method probably owing to decomposition while it was left on silica gel column for the

extended period. Arnottianamide 4 and the quaternary ammonium salts, chelerythrine 3 and nitidine 6 were obtained by passage of the chloroform extract through an open silica gel column.

The high-pressure liquid chromatography (hplc) results shown in Fig. 2 are worthy of comments. Namely, when X. granatum was extracted directly with hexane, the extract showed the hplc peak of dihydrochelerythrine 2, (Fig. 2). However, when treated with chloroform, this peak was replaced by that due to chelerythrine 3. In fact dihydrochelerythrine 2 is extremely unstable in solvents like chloroform, methanol and ethanol and is readily converted under normal laboratory conditions into chelerythrine 3 due to the action of air and light: the conversion is much slower, however, in hexane when applied on the plates, the colorless 2 is converted into the colored 3 within several minutes, similarly, the newly chopped bark of F. chalybea is colorless but it turns deep yellow after a few minutes, presumably due to the oxidation of 2 to 3. On the other hand, chelerythrine is readily reduced to the dihydro compound by sodium borohydride. Acetonyldihydrochelerythrine was only seen on the hplc trace when the root bark was extracted with acetone (Fig.2). These results imply that the well documented chelerythrine¹⁵ could be an artefact, acetonyldihydrochelerythrine has already been suggested to be an artefact¹⁸. As described above, arnottianamide 4 was obtained by passage through silica gel of the chloroform extract of F. chalybea. It was first isolated and characterized by Ishii and coworkers from the bark extract of Xanthoxylum arnottianum and they clearly showed that it could be produced from chelerythrine by oxidation with metaperchloroperbenzoic acid; an elegant oxidation mechanism has also been forwarded¹⁷. Due to the very facile transformations between 2 and 3, it would not be easy to disprove the presence of chelerythrine in nature, but at least it is safe to conclude that dihydrochelerythrine is by far the major constituent.

The structure of all compounds described in this communication were determined by a combination of spectroscopic methods including NOE measurements in the PMR to determine the substitution patterns of substituents on the aromatic ring.

Figure 1






Fig.2. Hplc of *X. granatum* extracts.
 μ -Porasil, 25:75 (ether:n-hexane).

TABLE 1

	<u>Fagara chalybea</u>	<u>F. holstii</u>	<u>Xylocarpus granatum</u>
1 N-methylflindersine ⁷	0.058% ^a	0.03	0.03 ^b
2 dihydrochelerythrine ¹⁴	0.18%	0.09	-
3 chelerythrine ^{15, 16}	0.0012%	-	-
4 arnottianamide ¹⁷	0.006%	-	-
5 acetonyldihydrochelerythrine ¹⁸	-	-	0.64
6 nitidine ^{16, 19}	0.0024%	-	-

a. Dry weight yield

b. Acetone was used for extraction.

ACKNOWLEDGEMENT

The authors are grateful to Mr. J. B. Gillet, East African Herbarium, Nairobi, and Dr. J. O. Kokwaro, Department of Botany, University of Nairobi, for identification of plant specimen, to Mr. Andrew Chapya, International Centre of Insect Physiology and Ecology (ICIPE), Nairobi, for assistance in plant collection and extraction. We are also grateful to Mr. I. Miura, Columbia University, for measurement of NMR Spectra, to Mr. J. Miller, Columbia University for bioassays, and in addition to the scientists quoted, in respective references, to the Eli Lilly Company for carrying out some of the bioassays. The studies were initiated during the stay of I. Kubo at ICIPE (May, 1974 to December 1975). We acknowledge support by NIH Grant AI 10187.

One of the authors (K. Hostettmann) expresses his gratitude to the Fonds National Suisse de la Recherche Scientifique for financial support, (Grant 820.425.76).

REFERENCES

- 1) a) P.R. Zanno, I.Miura, K.Nakanishi, and D.L.Elder, J.Am.Chem. Soc. 97 1975 (1975); b) I.Kubo, I.Miura, and K. Nakanishi, J.Am.Chem.Soc., 98, 6704 (1976); c) I.Kubo, Y-W. Lee, V.Balogh-Nair, K.Nakanishi, and A. Chapaya, J.C.S. Chem.Comm., 949 (1976); d) I. Kubo, S.P.Tanis, Y-W. Lee, I.Miura, K.Nakanishi, and A. Chapaya, Heterocycles, 5, 485 (1976); e) I.Kubo, Y-W.Lee, M.Pettei, F.Pilkiewicz, and K.Nakanishi, J.C.S.Chem.Comm., 1013 (1976); f) K. Nakanishi, Pontifical Acad.Sci., in press; g) I.Kubo and K.Nakanishi, ACS Symposium Series, The Chemical Basis for Host Plant Resistance to Pests(P.A.Hedin,ed), in press.
- 2) a) I.Kubo, K.Nakanishi, T. Kamikawa, T.Isobe, and T.Kubota, Chem.Lett., 99 (1977); b) I.Kubo, I.Miura, K.Nakanishi, T. Kamikawa, T.Isobe, and T.Kubota, J.C.S. Chem.Comm., 555,(1977).
- 3) K. Hostettmann, M.J.Pettei, I.Kubo, and K.Nakanishi, Helv. Chem. Acta, 60, 670 (1977).
- 4) In Ref. 16 we had reported the plant to be Xylocarpus moluscensis but according to information from Prof. D.A.H. Taylor, it should be X.granatum. We are following Prof. Taylor's suggestion, but since the literature is somewhat confused, the correct identity is being checked.
- 5) a) J.M.Watt and M.G. Breyer-Brandwijk, Medicinal and Poisonous Plants of Southern and East Africa, (1962), E.S. Livingstone Ltd., Edinburgh and London,; b) J.O. Kokwaro, Medicinal Plants of East Africa (1976), East African Literature Bureau, Nairobi.
- 6) see also: K.Wada and K.Munakata, Agr. Food Chem., 17, 471 (1968).
- 7) C.D.Adams, D.R.Taylor, and J.M.Warner, Phytochemistry 12, 1359 (1973).
- 8) The 2 second restriction in the period of dipping is to avoid extraction and loss of the natural phagostimulants from the leaves.

- 9) We are grateful to Suntory Co., Ltd., Osaka for this bioassay.
- 10) We are grateful to Prof.F.J.Schmitz, University of Oklahoma, for having the KB tests run at National Cancer Institute,Md.
- 11) M.Nagao, T.Yahagi, Y.Seino, and T.Sugimura, Cancer Lett. in press.
- 12) B.N.Ames, J.McCann, and E.Yamasaki, Mutation Res., 31, 347 (1975).
- 13) M.Nagao, National Cancer Center, Tokyo, private communication.
- 14) P.J.Scheuer, M.Y.Chang, and C.E.Swanholm, J.Org.Chem. 27, 1472 (1962).
- 15) R.H.F.Manske, in The Alkaloids (R.H.F.Manske and H.L.Holmes, eds.) Vol. IV, p253 (1954). Academic Press, New York.
- 16) F.Fish and P.G.Waterman, Phytochemistry 11, 3007 (1972).
- 17) H.Ishii and T.Ishikawa, Tetrahedron Lett., 1203 (1976).
- 18) P.D. Desai, T. R. Govindachari, K.Nagarajan and N.Viswanathan, Indian J.Chemy. 5, 42 (1967).
- 19) H.R.Arthur, W.H.Hui, and Y.L.Ng, J.Chem.Soc., 1840 (1959).
- 20) G.A. Cordell and N.R. Farnsworth, Heterocycles, 4, 393 (1976).

Received, 7th October, 1977