



PARTIAL SYNTHESIS OF 3-DEOXYDIHYDROMORPHINE FROM (-)-4-HYDROXY-6-KETO-N-METHYLMORPHINAN¹

Fu-Lian Hsu, Arthur E. Jacobson, Kenner C. Rice and Arnold Brossi*



Section on Medicinal Chemistry, Laboratory of Chemistry, National Institute of Arthritis, Metabolism and Digestive Diseases, National Institutes of Health, Bethesda, Maryland 20205, USA.

Abstract - Conversion of (-)-3-deoxydihydromorphine (2) into (-)-4-hydroxy-6-keto-N-methylmorphinan (5) and reconversion of 5 into 2 via the ketone 4 is described. The morphinan structure of 5 was proven by reduction to the known (-)-4-hydroxy-N-methylmorphinan (7).

3-Deoxydihydromorphine (2) and 3-deoxydihydromorphinone (4), easily obtainable from dihydromorphine (1)², represent valuable intermediates for the preparation of a variety of antinociceptive 3-deoxyopioids.² In addition the ketone 4, with the ether bridge linked at the α -position of the carbonyl group, constitutes an important entry into the relatively unexplored class of 4-hydroxymorphinans.³ Such a conversion could be accomplished as follows:



1. R₁ = R₃ = OH, R₂ = H
2. R₁ = R₂ = H, R₃ = OH
3. R₁ = R₃ = H, R₂ = OH



Reduction of ketone 4 with freshly prepared aluminium amalgam⁴ afforded the ketomorphinan 5 in 85-88% yield: mp 119-121°; $[\alpha]_D^{20}$ -142° (1.02, CHCl₃); ir ($\gamma_{\max}$, CHCl₃, cm⁻¹) 1710 (C=O), 3300 (OH); nmr (δ , CDCl₃) 6.94 (1H, dd, ArH, J = 8, 8 Hz), 6.64 (2H, m, ArH), 4.46 (1H, d, C₅-H, J = 13 Hz), 2.44 (3H, s, NCH₃); m/e 271 (M⁺). 5·HBr·1/2 H₂O: mp 210-3°; $[\alpha]_D^{20}$ -38.2 (1.10, MeOH). Bromination of 5 with 1.1 mole of bromine in acetic acid afforded the bromoketone 6·HBr in 50% yield: mp 218-221°, $[\alpha]_D^{20}$ -43.2° (0.92, MeOH); ir ($\gamma_{\max}$, KBr, cm⁻¹) 3300 (OH), 1705 (C=O); nmr (δ , CD₃OD) 7.31 (1H, d, ArH, J = 9.5 Hz), 6.63 (1H, d, ArH, J = 9.5 Hz), 4.30 (1H, d, C₅-H, J = 14 Hz), 2.97 (3H, s, NCH₃); m/e 349 (M⁺), 351 (M⁺ + 2). This bromination pattern is consistent with previously noted brominations⁵ and with the spectral evidence. Bromination of 5 with 2.5 mole of bromine in acetic acid and reduction of the mixture of brominated materials over 10% Pd-C in acetic acid and sodium acetate⁶ gave ketone 4 in 82% yield. The material thus obtained was identical in all respects to an authentic sample of 3-deoxydihydromorphinone (4).² The morphinan structure of 5 was confirmed by Wolff-Kishner reduction affording (-)-4-hydroxy-N-methylmorphinan (7) in 75-85% yield: mp 213-214.5° (lit.³ 216-217°); $[\alpha]_D^{20}$ -35.4° (1.27, MeOH) [lit.³ -35° (1.07, MeOH)]; ir ($\gamma_{\max}$, CHCl₃, cm⁻¹) 3350, 3600 (OH); nmr (δ , CDCl₃) 6.95 (1H, dd, ArH, J = 8, 8 Hz), 6.65 (1H, d, ArH, J = 8 Hz), 6.42 (1H, d, ArH, J = 8 Hz), 3.48 (1H, d, C₅-H, J = 13 Hz), 2.39 (3H, s, NCH₃); m/e 257 (M⁺). 7·HCl·EtOH: mp 282-4° (dec) lit.³ 303-5°; $[\alpha]_D^{20}$ -17.4° (0.998, MeOH) [lit.³ $[\alpha]_D^{25}$ -17.94° (0.958, MeOH)], and identical in all respects with a comparison sample prepared by total synthesis.^{3,7}

Reduction of ketone 4 with sodium borohydride in methanol at room temperature afforded a mixture of two epimeric alcohols in a ratio of 3:2, which could be separated by preparative tlc on silica gel (CHCl₃:MeOH:NH₄OH = 90:10:1). The faster moving alcohol and also the major one was identical with 3-deoxydihydromorphine (2). The slower moving alcohol 3 was characterized as its hydrochloride salt, 3·HCl·1/2 H₂O: mp 310-2° (dec); $[\alpha]_D^{20}$ -159.7° (1.1433, CH₃OH); ir ($\gamma_{\max}$, KBr, cm⁻¹) 3350 (OH); nmr (δ , CD₃OD) 7.37 (1H, t, ArH, J = 8 Hz), 6.76 (2H, dd, ArH, J = 8, 8 Hz), 4.32 (1H, d, C₅-H, J=8 Hz), 3.88 (1H, m, CH-OH), 2.95 (3H, s, NCH₃).

A selective high yield reduction of ketone 4 to 3-deoxydihydromorphine (2) was accomplished with lithium tri-*sec*-butylborohydride in tetrahydrofuran. This retrosynthesis of 3-deoxydihydromorphine (2) from the 4-hydroxy 6-keto-N-methylmorphinan (5) marks the latter an interesting intermediate for a total synthesis of 3-deoxyopioides.

References:

1. This work was presented by F. H. at the 20th Annual Meeting of the American Society of Pharmacognosy at Purdue University, July 29, 1979.

2. J. Reden, M. F. Reich, K. C. Rice, A. E. Jacobson, A. Brossi, R. A. Streaty, and W. A. Klee, J. Med. Chem., 1979, 22, 256.
3. E. Mohacsi and W. Leimgruber, U.S. Patent 3,914,234, 1975.
4. R. L. Augustine, "Reduction", Marcel Dekker, Inc., N. Y. 1968, p. 136.
5. H. L. Holmes in Manske "The Alkaloids", Vol. II, p. 48-49, 1952. A detailed examination of morphinans with varied substituents in the aromatic ring by ^{13}C NMR is in progress.
6. D. D. Weller and H. Rapoport, J. Med. Chem., 1976, 19, 1171.
7. We would like to thank Dr. E. Mohacsi from Hoffmann-La Roche Inc., Nutley, New Jersey, for providing a comparison sample of compound 4.

Received, 6th October, 1979