


REACTIVITY OF 5-METHYL-2H-1,2,6-THIADIAZIN-3(6H)-ONE 1,1-DIOXIDE  
TOWARDS ALKYLATING AGENTS AND CARBENES

José Elguero\*, Carmen Ochoa, and Manfred Stud

Instituto de Química Médica, CSIC, Juan de la Cierva 3, Madrid-6, Spain

*Abstract :* The reaction between 5-methyl-2H-1,2,6-thiadiazin-3(6H)-one 1,1-dioxide 1 and a variety of alkylating agents (diazomethane, trimethyl phosphite, dimethyl sulfate, and methyl iodide) in different conditions has been studied. Five methyl derivatives have been isolated and characterized. The reaction between 1 and carbenes proceeds with difficulty and only in the case of the carbene obtained from ethyl diazoacetate has the addition product been isolated.

Compound 1, 5-methyl-2H-1,2,6-thiadiazin-3(6H)-one 1,1-dioxide, is an isosteric analog of 6-methyluracil in which a carbonyl group is replaced by a sulfone group.<sup>1</sup> This molecule has four different nucleophilic positions, namely N<sub>2</sub>, O<sub>3</sub>, C<sub>4</sub>, and N<sub>6</sub>, thus, four monosubstituted, six disubstituted, and three trisubstituted derivatives may be expected from alkylation reactions. However, only five isomers (2, 3, 4, 5, and 6) could be isolated.<sup>2</sup>



The relative amounts of the methylated isomers are strongly dependent on the experimental conditions (Table 1). Treatment of compound 1 with an excess of diazomethane in an ethereal solution gave three isomers which were separated by TLC (toluene as eluent). These same isomers were also obtained with trimethyl phosphite but in a slightly different ratio. Reaction of 1 with dimethyl sulfate at room temperature afforded different isomers depending on the reaction medium used. Thus, when working in 1N NaOH only compound 2 (white needles from water) precipitated after acidification. When NaHCO<sub>3</sub> was used a complex mixture of four isomers was obtained. These were separated by TLC (chloroform as eluent). Finally, when compound 1 was dissolved in acetone, and methyl iodide and potassium carbonate (1:1) were added, a mixture of four isomers containing the 3,4,6-trisubstituted 6 was obtained. The relative amounts of the isomers (Table 1) have been determined by <sup>1</sup>H NMR on the crude reaction mixtures.

Table 1. Methylation of compound 1

| Experimental conditions                         | Total yield | 2   | 3  | 4  | 5  | 6  |
|-------------------------------------------------|-------------|-----|----|----|----|----|
| Diazomethane                                    | 95%         | --  | 55 | 30 | 15 | -- |
| Trimethyl phosphite                             | 75%         | --  | 50 | 45 | 5  | -- |
| Dimethyl sulfate (NaOH)                         | 60%         | 100 | -- | -- | -- | -- |
| Dimethyl sulfate (NaHCO <sub>3</sub> )          | 70%         | 5   | 65 | 25 | 5  | -- |
| Methyl iodide (K <sub>2</sub> CO <sub>3</sub> ) | 95%         | --  | 45 | 15 | 30 | 10 |

All the compounds have been obtained in an analytical grade of purity except 6 which was contaminated by 4. Their physical properties are gathered in Table 2.

Table 2. Physical data of methylated derivatives of compound 1

| Compound | R <sub>f</sub> <sup>a</sup> | mp <sup>b</sup>    | UV <sup>c</sup> | IR <sup>d</sup> | <sup>1</sup> H NMR <sup>e</sup> |      |      |      |                   |
|----------|-----------------------------|--------------------|-----------------|-----------------|---------------------------------|------|------|------|-------------------|
|          |                             |                    |                 |                 | 2                               | 3    | 4    | 5    | 6                 |
| 2        | 0.11                        | 184°C              | 262             | 3100, 1675      | 3.11                            | ---- | 5.48 | 2.05 | 8.7 (broad)       |
| 3        | 0.45                        | oil                | 264             | 1675            | 3.28 <sup>g</sup>               | ---- | 5.64 | 2.19 | 3.34 <sup>g</sup> |
| 4        | 0.63                        | 114°C <sup>f</sup> | 294             | 1610            | ----                            | 3.90 | 5.46 | 2.25 | 3.42              |
| 5        | 0.67                        | oil                | 259             | 1670            | 3.27 <sup>g</sup>               | ---- | 1.96 | 2.15 | 3.17 <sup>g</sup> |
| 6        | 0.78                        | oil                | ---             | ----            | ----                            | 3.98 | 1.93 | 2.06 | 3.42              |

<sup>a</sup>Silica gel, eluent : chloroform ; <sup>b</sup>uncorrected ; <sup>c</sup> $\lambda_{\text{max}}$  (nm) in 95% ethanol ; <sup>d</sup>cm<sup>-1</sup>, pure compounds (oils) or nujol mull (solids) ; <sup>e</sup>solvent : CDCl<sub>3</sub> except for 2 (DMSO-d<sub>6</sub>) ; <sup>f</sup>crystallized from ethanol-water ; <sup>g</sup>see note 3.

The position of the methyl group was established according to the data of Table 2. Compound 2 is an N-monosubstituted derivative (N-Me at 3.11 ppm) ; the fact that the nitrogen atom is N<sub>2</sub> was deduced from the UV data since the corresponding 2-n-butyl and 6-n-butyl derivatives (prepared from n-butylsulfamide<sup>4</sup>) show their maxima, respectively, at 258 and 295 nm. Compound 3 is an N<sub>2</sub>,N<sub>6</sub>-di-substituted derivative (N-Me at 3.28 and 3.34 ppm). Compound 4 is an N<sub>2</sub>,O-disubstituted derivative (N-Me at 3.42 and O-Me at 3.90 ppm) with an electronic spectrum of a fully conjugated structure. Compound 5 is similar to 3 with a supplementary methyl group bonded to carbon C<sub>4</sub> (C-Me at 1.96 ppm). Finally, compound 6 closely resembles compound 4 ; its third methyl group is bonded to C<sub>4</sub> (C-Me at 1.93 ppm).


Some of these compounds have been studied by <sup>13</sup>C NMR spectroscopy (Table 3).

Table 3. Carbon-13 NMR spectra of 2H-1,2,6-thiadiazin-3(6H)-one 1,1-dioxides (ppm from TMS)

| Compound | Solvent             | C <sub>3</sub> | C <sub>4</sub> | C <sub>5</sub> | Me <sub>5</sub> | N <sub>2</sub> -Me | O-Me | N <sub>6</sub> -Me |
|----------|---------------------|----------------|----------------|----------------|-----------------|--------------------|------|--------------------|
| 1        | DMSO-d <sub>6</sub> | 162.3          | 89.6           | 156.4          | 19.5            | ----               | ---- | ----               |
| 2        | DMSO-d <sub>6</sub> | 162.4          | 98.7           | 151.0          | 19.8            | 25.6               | ---- | ----               |
| 3        | CDCl <sub>3</sub>   | 161.8          | 104.6          | 150.9          | 20.1            | 27.5               | ---- | 32.7               |
| 4        | CDCl <sub>3</sub>   | 167.0          | 92.1           | 157.4          | 20.7            | ----               | 54.7 | 30.8               |


The differences observed in the  $^{13}\text{C}$  chemical shifts of  $\text{C}_4$  and  $\text{C}_5$  when  $\underline{1}$  and  $\underline{2}$  are compared may arise from tautomeric modifications (presence of 3-hydroxy or 4H-tautomers). Otherwise, the spectra are consistent with the structures proposed.

The results obtained with the *n*-butyl derivatives  $\underline{3}$  and the fact that it was not possible to C-methylate compounds  $\underline{3}$  and  $\underline{4}$  (to obtain  $\underline{5}$  and  $\underline{6}$  respectively)<sup>5</sup> shows that the simplest tree which connects all the compounds is the following one (3-methyl derivatives are methoxy compounds) :





Pandit *et al.*<sup>6</sup> have studied the reaction of carbenes with uracil and uridine derivatives and isolated the adducts both with dihalogenocarbenes and with ethoxycarbonylcarbene. In a similar way and following the analogy between uracil and our thiadiazine 1,1-dioxides we have studied the addition of carbenes to the  $\text{C}_4\text{-C}_5$  double bond of compound  $\underline{3}$ .

When the thiadiazinone 1,1-dioxide  $\underline{3}$  was allowed to react with ethoxycarbonylcarbene (generated by copper catalyzed decomposition of ethyl diazoacetate<sup>7</sup>) in dimethoxyethane, in addition to the starting material the following adduct was isolated in 30% yield (TLC, hexane-ethyl acetate 4:1).



It is an analytically pure oily product, with no UV absorption in EtOH (showing the saturation of the  $\text{C}_4\text{-C}_5$  double bond) and with IR bands (pure sample) at 3040 (cyclopropane ring) and 1725 and 1720  $\text{cm}^{-1}$  ( $\text{C=O}$ ). In mass spectrometry the highest  $m/e$  peak appears at 231 (M-EtO). In  $^1\text{H}$  NMR (solvent :  $\text{CDCl}_3$ ) the value of the coupling constant between  $\text{H}_4$  and  $\text{H}_7$  (6.0 Hz) proves the *trans* configuration of both protons.<sup>6</sup> No *endo* isomer was observed. The  $^{13}\text{C}$  NMR spectrum (same solvent) is also in agreement with the proposed structure (signals marked with \* or † may be reversed).



Other experiments were carried out to prepare cyclopropane adducts from thiadiazinone 1,1-dioxide derivatives. The reactions between 5 and ethoxycarbonylcarbene, 3 and 5 and diazomethane (catalyzed with palladium acetate<sup>8</sup>), and 3 and dimethylsulfoxonium methylide<sup>9</sup> were unsuccessful: in all the cases, the starting materials were recovered unchanged.

**Acknowledgement.** This work was carried out with financial support from the "Comisión Asesora de Investigación Científica y Técnica (Presidencia del Gobierno)".

#### REFERENCES AND NOTES

1. R. Die, J. Díez, G. García-Muñoz, R. Madroñero, and M. Stud, *J. Heterocyclic Chem.*, 1972, 9, 973.
2. In a preceding paper (P. Goya, P. Martínez, C. Ochoa, and M. Stud, *J. Heterocyclic Chem.*, 1981, 18, 459) methylation (dimethyl sulfate, diazomethane) of 4-nitro, 4-cyano, and 4-ethoxycarbonyl-2H-1,2,6-thiadiazin-3(6H)-one 1,1-dioxide has been described. Depending on the thiadiazine and the alkylating agent, 2-methyl, 6-methyl, and 3-methoxy-6-methyl derivatives could be isolated.
3. Methylation ( $\text{ICH}_3/\text{K}_2\text{CO}_3/\text{acetone}$ ) of 5-methyl-2-n-butyl-2H-1,2,6-thiadiazin-3(6H)-one 1,1-dioxide<sup>4</sup> gave a mixture of the 5,6-dimethyl ( $\text{Me}_6$ : 3.36 ppm) and 4,5,6-trimethyl ( $\text{Me}_6$ : 3.19 ppm) derivatives. Methylation in the same conditions of 5-methyl-6-n-butyl-2H-1,2,6-thiadiazin-3(6H)-one 1,1-dioxide<sup>4</sup> gave a mixture of the 2,5-dimethyl ( $\text{Me}_2$ : 3.31 ppm) and 3-methoxy-5-methyl (the 6-butyl analog of 4) isomers. These chemical shifts have been used to assign the N-methyl signals of compounds 3 and 5 (Table 2).
4. J. Díez, G. García-Muñoz, R. Madroñero, and M. Stud, *J. Heterocyclic Chem.*, 1973, 10, 469.
5. The following experiments have been carried out: reaction of 3 and 4 with diazomethane and with ethyl bromoacetate ( $\text{K}_2\text{CO}_3/\text{acetone}$ ); in both cases, the starting materials were recovered unchanged.
6. H.P.M. Thiellier, G.J. Koomen, and U.K. Pandit, *Tetrahedron*, 1977, 33, 1493.
7. A.P. Marchand and N. McBrockway, *Chem. Rev.*, 1974, 74, 431.
8. A.J. Anciaux, A.J. Hubert, A.F. Noels, N. Petiniot, and P. Teyssié, *J. Org. Chem.*, 1980, 45, 695.
9. E.J. Corey and M. Chaykowsky, *J. Amer. Chem. Soc.*, 1962, 84, 867 and 3782.

Received, 5th August, 1981