


LIPOPHILIC AND AMPHIPHILIC CYCLODEXTRIN DERIVATIVES AS THE
NEW CLASS OF HOST COMPOUNDS

Waichiro Tagaki, Yuichi Irinatsu, Hisao Takahashi and Seizi Kozuka

Department of Applied Chemistry, Faculty of Engineering, Osaka
City University, Sugimoto 3, Sumiyoshi-ku, Osaka 558, Japan

We wish to report the host-guest interactions in some lipophilic and amphi-
philic cyclodextrin derivatives as the hosts. The host compounds were prepared
by specific modification of β -cyclodextrin as outlined below:



When R = C₁₂H₂₅, the hosts 4 and 5 were lipophilic and soluble in chloroform. It was observed that 4a (R'=H, R=C₁₂H₂₅) was able to extract p-nitrophenol from the aqueous layer into the chloroform layer in a chloroform-water two system, in contrast to the inability of a model amine (dodecylmethylamine) to do it. In chloroform, 4a was also observed to shift an equilibrium of p-nitrophenol dissociation toward the phenolate anion, presumably by inclusion. Furthermore, the protonated 4a ammonium ion was soluble in water as a surfactant and showed inclusion interactions with hydrophobic anions. The host 5a (R=C₁₂H₂₅) was insoluble in water, but could be solubilized in surfactant micelles in water, and a co-micelle of 5a-CTABr cationic surfactant showed an enhanced catalytic activity in the hydrolysis of p-nitrophenyl carboxylates as compared to those of unsubstituted β -CD and CTABr micelle. Further studies on these host compounds as well as on the related compounds are in progress.