

ASYMMETRIC SYNTHESIS OF β -LACTAM ANTIBIOTICS


A.Nishida, M.Shibasaki,* and S.Ikegami

Faculty of Pharmaceutical Sciences, Teikyo University, Sagamiko, Kanagawa 199-01, Japan

As a part of our synthetic studies toward the β -lactam antibiotics, we have pursued the total synthesis of (+)-thienamycin from (+)-4-phenylthio-2-azetidinone ((+)-2) prepared by utilizing the asymmetric introduction of thiophenol to ($\pm$)-4-phenylsulfonyl-2-azetidinone (($\pm$)-1) in the presence of cinchonidine. In this presentation, at first, the detail of the asymmetric synthesis of (+)-2 and its conversion to (+)-thienamycin are presented.

In the second, the synthesis of the key synthetic intermediate (10) for optically active sulfazecin (3) is presented. Namely, the protected azetidinone (5) derived from (+)-2 was converted to the amine (6) by azidation (2-naphthalenesulfonyl azide) and reduction (H_2S), which was transformed to the benzyloxycarbonyl derivative (7) by the standard method. The introduction of a methoxy group to 7 was carried out by reaction with t -BuOCl and methanolic LiOMe in CH_2Cl_2 . Since the desulfurization of 8 catalyzed by Raney-Nickel did not give any desired product, 8 was converted to the sulfone (9), which could be converted to the key intermediate (10) for sulfazecin by reductive removal of the sulfonyl group with $NaBH_4$ in THF.

The synthesis of the key intermediate (11) for the synthesis of SQ26776 (4) was also attempted by employing the enantiomeric isomer (-)-2 obtainable from the similar asymmetric introduction of thiophenol.

Ns=2-naphthalenesulfonyl, Z=-COOCH₂Ph

* Present address; Sagami Chemical Research Center, Nishi-Ohnuma, Sagamihara, Kanagawa 229, Japan