


ENANTIOSELECTIVE SYNTHESIS OF OPTICALLY ACTIVE β -LACTAMS FROM
D-GLYCERALDEHYDE ACETONIDE

Hiroshi Matsunaga, Tomoko Sakamaki, Hiroto Nagaoka, and Yasuji Yamada*
Tokyo College of Pharmacy, 1432-1 Horinouchi, Hachioji, Tokyo 192-03, Japan

D-Glyceraldehyde acetonide (1) easily prepared from D-mannitol is known to be a useful and inexpensive starting material for the various optically active natural products. Here we have accomplished the enantioselective synthesis of optically active β -lactams, (R)-N-benzyl-4-[(methoxycarbonyl)methyl]-2-azetidinone (4) and (S)-4-[(methoxycarbonyl)methyl]-2-azetidinone (5), from 1.

The 1,4-addition reaction of the α,β -unsaturated ester 2, prepared from 1 by Wittig reaction, with benzylamine at -50°C gave highly stereoselectively (3R)-amino ester 3 in 85% yield. Two ways for β -lactam formation (route A and route B) from the resulting β -amino ester 3 provided the enantioselective synthesis of 4 and 5 as shown in Scheme. The β -lactam 4 can be transformed into unnatural series of carba-



Scheme

penem antibiotics, and the β -lactam 5 is a versatile intermediate for the synthesis of natural (+)-thienamycin.¹

An alternative approach to the β -lactam 5 from 2, and the stereoselective Michael type addition of various amines to 2 will be also described.

1) Formal total synthesis of (+)-thienamycin from D-mannitol has thus been accomplished.