

SYNTHESIS AND STRUCTURE OF NOVEL HALOSULFURANES, 5-CHLORO- AND
5-BROMO-5,11-DIHYDRODIBENZO[b,e]THIEPINS

Kazuhiro Onogi,* Mikio Hori,** Tadashi Kataoka**

Hiroshi Shimizu,** and Masaru Kido***

*Tokyo Research Laboratories, Kowa Co., Ltd., 2-17-43,
Noguchi-cho, Higashimurayama 189, Japan

**Gifu College of Pharmacy, 6-1, Mitahora-higashi 5-chome,
Gifu 502, Japan

***Laboratories of Natural Product Chemistry, Otsuka Pharmaceutical
Co., Ltd., Kawauchi-cho, Tokushima 771-01, Japan

5-Chloro- (2a,b) or 5-bromo-5,11-epoxy-5,11-dihydrodibenzo[b,e]thiepins (3a,b) were synthesized by the reaction of dibenzothiepin-11-ols (1a,b) with N-chloro- (NCS) or N-bromosuccinimide (NBS), respectively. Chlorosulfuranes (2a,b) were more stable thermally and hydrolytically than bromosulfuranes (3a,b).

The covalency of the sulfur-halogen bond of the halosulfuranes (2a,b-3a,b) was established by the $^1\text{H-NMR}$ and MS (field desorption) spectra, and X-ray structure determination. The absorption of $\text{C}_4\text{-H}$ at δ 9.11-9.24 for 2a was shifted to downfield relative to that of the corresponding sulfonium salt 4a in the $^1\text{H-NMR}$ spectra. Other halosulfuranes (2b,3a,b) showed this distinctive downfield shifts (0.6-0.7 ppm lower than those of 4a,b). The MS (field desorption) spectra of the sulfuranes showed the molecular ion peaks. The crystal structure of 2a reveals approximate trigonal bipyramidal geometry about sulfur. The apical S-Cl bond length (2.749 Å) is ca 0.72 Å longer than the sum of the covalent radii (2.03 Å) and the apical S-O bond length (1.639 Å) is ca 0.06 Å shorter than the sum of the covalent radii (1.70 Å).

