


ISOLATION AND CHARACTERIZATION OF A THIADIPHOSPHIRANE

Masaaki Yoshifuji, Katsuhiko Shibayama, Kaori Ando, and Naoki Inamoto

Department of Chemistry, Faculty of Science, The University of Tokyo, Hongo, Tokyo 113, Japan

Phosphorus containing small ring compounds, especially three-membered ring compounds, are of current interest. By introducing an extremely bulky group into a molecule, we were successful in the preparation and characterization of *E*-2,3-bis(2,4,6-tri-*tert*-butylphenyl)-1,2,3-thiadiphosphirane (2) as a stable compound by the following two different methods.

a) The phosphonothioic dichloride (1) was allowed to react with magnesium in tetrahydrofuran to give 2 in 54% yield as a stable crystalline material (mp 131.5 - 132°C, δ_P -65.1 ppm). 2 was thermally and photochemically stable and insensitive to atmospheric moisture and oxygen for a long period of time.



b) The orange diphosphene (4, mp 175 - 176°C, δ_P 492.4 ppm), obtained from the phosphonous dichloride (5) with magnesium as the first isolated P=P compound, was allowed to react with elemental sulfur in triethylamine to give the yellow diphosphene sulfide (3, mp 151.5 - 152°C), which was irradiated or heated in toluene to give 2 almost quantitatively. The intermediate 3 was isolated as a stable compound and was characterized: δ_P 255.8 and 247.8 ppm, J = 629.9 Hz as an AB quartet. 3 was desulfurized with tris(dimethylamino)phosphine to give 4 almost quantitatively while 2 resisted even under severer reaction conditions.

The compounds 2, 3, and 4 were analyzed by X-ray crystallography, for which we are indebted to Dr. Ken Hirotsu and Prof. Taiichi Higuchi at Osaka City University, Japan.