

REISSERT TYPE'S REACTION BY USE OF PHOSPHITE


Yoshiki Hamada, Isao Takeuchi, and Yasuhiro Shibata

Faculty of Pharmacy, Meijo University

Yagoto-Urayama, Tempaku-cho, Tempaku-ku, Nagoya 468, Japan

We have reported the Reissert reaction to the naphthyridine and phenanthroline.¹⁾ Recently, Akiba and coworker²⁾ reported the synthesis of phosphonates by reaction of pyridine, quinoline, and isoquinoline. We extended the reaction to the other heterocyclic compound. Reaction of benzo[f]quinoline, phthalazine, and 1,X-naphthyridine (X=5,6,8) by use of trimethyl phosphite in place of potassium cyanide gave the corresponding Reissert type's compound and benzo[h]quinoline did not give it. Benzo[f]quinoline gave the isomer of 3-phosphonate (1) and 1-phosphonate (2).

These structure of phosphonate were confirmed by the comparison of ¹H-NMR spectral data of phosphonate (3) reported in the literature²⁾ and synthesized phosphonates (4, 5).



1) Y. Hamada and K. Shigemura, Yakugaku Zasshi, **99**, 982 (1979).

2) K. Akiba, Y. Negishi, and N. Inamoto, Synthesis, **1979**, 55.