

A NEW, SYNCHRONOUS CONSTRUCTION OF SOME FUSED PYRIMIDINES BY THE
REACTION OF 6-ARYLIDENEHYDRAZINO-1,3-DIMETHYLURACILS WITH
THIONYL CHLORIDE

Misuzu Ichiba, Sadao Nishigaki, and Keitaro Senga

Pharmaceutical Institute, School of Medicine, Keio University

35, Shinanomachi, Shinjuku-ku, Tokyo 160, Japan

We have previously reported that the reaction of 6-hydrazinouracils with thionyl chloride offers [1,2,3]thiadiazolo[4,5-*d*]pyrimidines and their mesoionic compounds, a new class of heterocycles [J. Org. Chem., **43**, 1677 (1978)]. In connection with these findings, we now report the reaction of 6-arylidenehydrazino-1,3-dimethyluracils with thionyl chloride, which results in a new, synchronous construction of some biologically intriguing fused pyrimidines.

Treatment of 6-arylidenehydrazino-1,3-dimethyluracil (I) with excess thionyl chloride in benzene afforded purine (II), pyrazolo[3,4-*d*]pyrimidine (III), thiazolo[4,5-*d*]pyrimidine (IV), [1,2,3]thiadiazolo[4,5-*d*]pyrimidine (V), and pyrimido[4,5-*e*][1,3,4]thiadiazine (VI), while the reaction of 6-(arylidene-1-methylhydrazino)-1,3-dimethyluracil (VII) with thionyl chloride gave IV, VI, and pyrazolo[3,4-*d*]pyrimidine (VIII). The mechanism for the formation of fused pyrimidines will also be discussed.

