


THE PSEUDO-GOMBERG REACTION OF 4-AMINOPYRIDINE N-OXIDE

S. Saeki, S. Kondo, T. Hayashi and M. Hamana*


Faculty of Pharmaceutical Sciences, Kyushu University

Maidashi 3-1-1, Higashi-ku, Fukuoka 812, Japan

As an extension of work on the introduction of carbon substituents into the 4-position of pyridine ring¹⁾, we examined the psuedo-Gomberg reaction of 4-amino-pyridine N-oxide and found that treatment with aromatic hydrocarbons and amyl nitrite in acetic acid afforded the corresponding 4-arylpyridine N-oxides (1 and 2) in generally good yields. Reactions with mono-substituted benzene gave 4-(o-substituted-phenyl)pyridine N-oxides as the main products, except for that with nitrobenzene in which the p-nitrophenyl derivatives was dominant. The 1-oxido-4-pyridyl radical in these reactions was shown to have a rather electrophilic character by caluculations of relative rates (c.g., $K_{\text{PhOMe}}/K_{\text{PhH}}=1.8$, $K_{\text{PhNO}_2}/K_{\text{PhH}}=0.11$). On the other hand, o-substituted products were obtained as the sole products in reaction of furan, thiophene and N-methanesulfonylpyrrole (3).



Whereas aminopyridines resist the reaction with 3, 2- and 3-aminopyridine N-oxides also readily reacted with 3. Thus, we succeeded in the simple synthesis of nornicotyrine in the overall yield of 50% as shown below.



1) M. Hamana et al., *Heterocycles*, 11, 371 (1978).

* Present address: Central Res. Lab., Chugai Pharm. Co. Ltd., Takada 3-41-8, Toshima-ku, Tokyo 171, Japan.