

OXIDATION REACTIONS OF BICYCLIC PYRROLE DERIVATIVES







Yoshio OTSUJI, Nobuhiko OHMURA, Saburo NAKANISHI, and Kazuhiko MIZUNO
Department of Applied Chemistry, College of Engineering, University
of Osaka Prefecture, Mozu-Umemachi, Sakai, Osaka 591, Japan

The NaIO_4 oxidation and the photosensitized oxygenation of bicyclic pyrrole derivatives were studied, and chemical properties of hydroperoxides obtained by the photooxygenation were explored.

The oxidation of 2-substituted bicyclic pyrrole derivatives 1a-g with NaIO_4 in aqueous methanol at -5°C gave 8-11 membered cyclic ketolactams 2a-g in good yields. In these reactions, the double bond involved in the ring fusion was regioselectively oxidized.

Irradiation of 1a-f in methanol at $-50 - 0^\circ\text{C}$ in the presence of methylene blue (MB) under a continuous stream of O_2 with a 300 W halogen lamp gave hydroperoxides 3a-f in quantitative yields. The mechanistic study suggested that $^1\text{O}_2$ is an active species for this photooxygenation. The selectivity in this photo-reaction was extremely high.

The reduction of 3a-f with PPh_3 in Et_2O or Me_2S in MeOH afforded alcohols 4a-f in quantitative yields. The pyrolysis of 4a-d at 180°C for 3 min under argon atmosphere gave spiro lactams 5a-d in high yields (80-90%) via a 1,5-sigmatropic rearrangement. However, the pyrolysis of 4e-f under similar conditions afforded bicyclic compounds with an olefinic double bond 6e-f in 30-40% yields. The reaction of 3a with FeSO_4 in a 2N H_2SO_4 aqueous solution or with $\text{Fe}_3(\text{CO})_{12}$ in benzene yielded spiro lactam 5a in 80-90% yields.



a; $n=2$, $\text{X}=\text{H}$

b; $n=2$, $\text{X}=\text{CH}_3$

c; $n=2$, $\text{X}=\text{Cl}$

d; $n=2$, $\text{X}=\text{Br}$

e; $n=3$, $\text{X}=\text{H}$

f; $n=4$, $\text{X}=\text{H}$

g; $n=1$, $\text{X}=\text{H}$