

SYNTHESIS OF THIAZOLINE, THIOPHENE, AND 1,3-DITHIOLANE
USING ETHYL 4-HALOACETOACETATE

Takuo Chiba, Hirotoshi Sato, and Tetsuzo Kato

Pharmaceutical Institute, Tohoku University,


Aobayama, Sendai 980, Japan

Synthesis of heterocycles using ethyl 4-bromo(and 4-chloro)acetoacetate (1 and 2) is described. Thus, the esters 1 and 2 reacted with *N*-substituted dithiocarbamates prepared from carbon disulfide and amines to give the corresponding 3-substituted 2-thioxo-4-thiazoline (3a - 3e).

Reaction of 1 (and 2) with the thioacetamides prepared from phenyl isothiocyanate and active methylene compounds gave 2-substituted thiazolidines (4a - 4c) and the thiophenes (5a and 5b).


Compound 1 was allowed to react with the 2-cyanoethene-1,1-dithiol to give 2-substituted 4-hydroxy-1,3-dithiolane-4-acetates (6a and 6b).

Reaction of 1 with the 2-cyano-1-methylthioethene-1-thiol yielded the 5-(methylthio)thiophenes (7a and 7b).



1: X = Br

2: X = Cl




3a: R = CH₂Ph

3b: R = Me

3c: R = Et

3d: R = iso-Pr


3e: R = Ph



4a: R = CN, R' = CO₂Et


4b: R = R' = CN

4c: R = R' = CO₂Et




5a: R = CN

5b: R = CONH₂



6a: R = CN

6b: R = CO₂Et



7a: R = CN

7b: R = CO₂Et