


AIR OXIDATION REACTION OF DIAZOALKENES
EFFICIENT OXYGEN-TRANSFER REACTION TO GIVE EPOXYKETONES

Tsutomu Miyashi, Masaki Kamata, Yoshinori Nishizawa, Toshio Mukai

Photochemical Research Laboratory and Department of Chemistry

Faculty of Science, Tohoku University, Sendai 980, Japan

We have reported that allyldiazomethane **1a** ($n=1$, $R_1=C_6H_5$, $R_2=H$) undergoes a nitrene-type 1,1-cycloaddition¹ to give *exo*-6-phenyl-1,2-diazabicyclo[3.1.0]hex-2-ene **2a**. In order to explore the generality of the 1,1-cycloaddition reaction of diazoalkenes, reactivities of homologous diazoalkenes, **1b** ($n=2$, $R_1=C_6H_5$, $R_2=H$), **1c** ($n=2$, $R_1=R_2=C_6H_5$), **1d** ($n=3$, $R_1=C_6H_5$, $R_2=H$) and **1e** ($n=4$, $R_1=C_6H_5$, $R_2=H$) were tested, generating from the sodium salts of the corresponding tosylhydrazones. It was found that diazoalkenes **1d** and **1e** did not undergo the 1,1-cycloaddition but the 1,3-dipolar addition to give **3d** ($n=3$) and **3e** ($n=4$) under inert atmosphere, whereas **1b** and **1c** were unexpectedly stable and remained unchanged for more than 2 months when kept under inert atmosphere. In contrast with these reactivities under inert atmosphere, diazoalkenes **1b** and **1c** were readily oxidized by molecular oxygen, giving oxygen-transfer products, epoxyketones **4b** ($R_1=C_6H_5$, $R_2=H$) and **4c** ($R_1=R_2=C_6H_5$) when **1b** and **1c** were exposed to air. It is of interest to note that singlet oxygen did not give such epoxyketones at all. A possible mechanism will be discussed.



References: 1) Y. Nishizawa, T. Miyashi, T. Mukai, *J. Am. Chem. Soc.*, **102**, 1176 (1980); T. Miyashi, Y. Fujii, Y. Nishizawa, T. Mukai, *ibid.*, **103**, 725 (1981); see also Ref.2

2) A. Padwa, H. Ku, *Tetrahedron Lett.*, 1009 (1980); A. Padwa, A. Rodoriguez, *ibid.*, 187 (1981)