

SYNTHESIS OF 2-PYRONE DERIVATIVES USING KETENETHIOACETAL(3)

Yoshinori Tominaga, Atsuyuki Ushiroguchi, Yoshiro Matsuda,
and Goro Kobayashi

Faculty of Pharmaceutical Sciences, Nagasaki University,
Bunkyo-machi 1-14, Nagasaki 852, Japan

Ketenethioacetals appropriately functionalized (cyano, ester carbonyl, ketone carbonyl, sulfonyl groups and etc.) are very usefull reagents for the synthesis of heterocyclic compounds. We now wish to report the synthesis of 2-pyrone derivatives using dimethyl bis(methylthio)methylenemalonate (1).

The reaction of compound 1 with the acetyl compounds (acetophenone, p-methoxyacetophenone, 3,4-dimethoxyacetophenone, 3,4-methylenedioxyacetophenone, p-bromoacetophenone, and 3-acetylpyridine) in the presence of powdered potassium hydroxide as a base in dimethyl sulfoxide gave the corresponding 6-substituted 3-methoxycarbonyl-4-methylthio-2-pyrone derivatives(3a, b, c, d, e, and f) in good yields. Similarly, p-methoxybenzalacetone and 3,4-methylenedioxybenzalacetone were allowed to react with 1 to yield the corresponding 6-substituted styryl-3-methoxycarbonyl-4-methylthio-2-pyrones (4a and b) in 15% and 17% yields, respectively.

Compounds 3 (a: 6-phenyl, b: 6-(p-methoxy)phenyl, c: 6-(3,4-dimethoxy)phenyl, d: 6-(3,4-methylenedioxy)phenyl) were treated with sodium methylate in methanol to give the corresponding 6-substituted phenyl-4-methoxypyrones (5a: 4-methoxyphenyl-coumalin, 5b, 5c, and 5d: 4-methoxyparacotoin) in good yields. Phenylcoumalin (6-phenyl-2-pyrone) (7a) and paracotoin (6-(3,4-methylenedioxy)phenyl-2-pyrone) (7b) were obtained by the desulfurisation of 4-methylthio-6-phenyl-2-pyrone(6a) and 6-(3,4-methylenedioxy)phenyl-4-methylthio-2-pyrone (6b) which were prepared by the decarboxylation with polyphosphoric acid in good yields.

