


INDIRECT ELECTROOXIDATION OF tert-AMINES WITH RuO₄

Sigeru Torii, Tsutomu Inokuchi, and Takao Yukawa
Department of Industrial Chemistry, School of Engineering, Okayama
University, Okayama, Japan 700

RuO₄ can be used as a versatile reagent for various oxidative transformations. Instead of usual methods using periodates or hypochlorites as co-oxidants, we developed electrochemical process for generation of RuO₄ from RuO₂ and attempted to oxidize cyclic and acyclic tert-amines 1 thereby. As shown in the schematic picture (Chart I), we found that RuO₄ generated in situ in aqueous NaCl can act as an efficient mediator in the electrolysis system for the conversion of 1 to amides 2.



Typically, electrooxidation of 1 (0.6 mmol) was carried out using RuO₂·2H₂O (0.02 mmol) in a acetone-saturated NaCl (2.5 ml-5.0 ml)-(Pt-Pt) system under a constant current of 10 mA/cm² (applied voltage: 2.0-.2.4 V; electricity: 7.0 F/mol), giving the desired 2 in 92% yield. It is interesting to note that only methylene group at the α-position are readily oxidized to carbonyl function and no reaction has been observed at the alkyl substituted α'-position of nitrogen. Cyclic amines are much more easily oxidized than linear ones.

The present oxidation procedure of amines can be used for selective functionalization of nitrogen containing compounds. For example, we demonstrate here transposition of α,β-enone 3 to reversed α,β- unsaturated amide 6. Thus, electrooxidation of 4 prepared by amination of 3 followed by reduction with NaBH₄ and protection of both amino and hydroxyl groups gave 5 in 77% yield whose dehydroacetoxylation with DBU afforded the desired 6, smoothly.

