


SYNTHESIS AND PROPERTY OF 5-(2,2-DIFLUOROVINYL)URACIL DERIVATIVES

Kosaku Hirota, Fumiaki Iwami, Yukio Kitade, Shigeo Senda, and
Yoshifumi Maki

Gifu College of Pharmacy, Mitahora-higashi, Gifu 502, Japan

There has been great interest in 5-substituted pyrimidine nucleosides as potential antiviral and anticancer agents. Recently, it was found that 5-bromovinyl-2'-deoxyuridine (BVDU) was highly efficacious against herpes simplex virus type 1. In view of structural similarity to the BVDU, we planned to synthesize 5-(2,2-difluorovinyl)-2'-deoxyuridine (1). The present paper describes the results of the preliminary investigation on the synthesis and property of 5-(2,2-difluorovinyl)uracil derivatives.



BVDU; X=Br, Y=H

(1); X=Y=F

The reaction of 5-formyl-1,3-dimethyluracil with Ph_3P and $\text{ClCF}_2\text{COONa}$ in dry acetonitrile under argon resulted in the formation of 5-(2,2-difluorovinyl)-1,3-dimethyluracil (2) in 86% yield. Analogously, the reaction of 2',3',5'-tri-O-benzoyl-5-formyluridine with Ph_3P and $\text{ClCF}_2\text{COONa}$ afforded the corresponding 5-(2,2-difluorovinyl)uridine derivative (6).

Reactivities of (2) toward nucleophiles were observed to be higher than those of 5-bromovinyl-1,3-dimethyluracil. For example, treatment of (2) with NaOR ($\text{R}=\text{Me}$, Et) and methylamine gave the corresponding esters (3a, 3b) and amide (4) in high yields, respectively. The reaction of (2) with thiophenol led to the smooth formation of the adduct (5).

