

SYNTHESIS OF THE CYCLOBUTANE DERIVATIVES BY PHOTOCHEMICAL
CYCLOADDITION OF 1,2- OR 1,4-BENZOPYRONES TO ACETYLENE

Takano Matsui and Mitsuru Nakayama

Faculty of Engineering, Miyazaki University, Miyazaki 880 and

Faculty of Science, Hiroshima University, Hiroshima 730

We previously reported that photochemical cycloaddition of α,β -unsaturated carbonyl compounds, such as 4a,5,6,7,8,8a-hexahydro-4a-methoxycarbonyl-7-methylenedioxy-4-methylcoumarin with acetylene produced the 3:4-fused cyclobutane ring system in 62% yield, which was led to construct the five-membered ring of trichothecane nucleus (J. D. White, T. Matsui, and J. A. Thomas, *J. Org. Chem.*, **46**, 3376 (1981)). We now report the photocycloaddition products from 7- or 6-methoxychromones (1 or 5) and 7-methoxy-4-methylcoumarin (6) with acetylene and functionalization of the products.

Irradiation of a solution of 1 in acetone with introduction of acetylene using a 500-W high pressure mercury lamp through a Pyrex filter for 5 h afforded dihydrocyclobuta[b]chromone (2, mp 96-97°C, 35% yield) and its valence isomer, bicyclobutane (3, mp 64-67°C, 10% yield). Catalytic hydrogenation of 2 over Pd-C in ethanol yielded the cyclobutane derivative (4) in 72% yield. In the isomerization between valence isomers 2 and 3, when a solution of 2 in acetone was irradiated, 3 was obtained in 30% yield. However, irradiation of 3 led to recovery of the starting material. Whereas the thermal reaction of 3 at 205-210°C for 10 min gave 2 (26% yield). In contrast to the photochemical activity of 1 with acetylene, neither 5 nor 6 was reactive toward acetylene. Consequently, 5 was almost recovered. On the other hand, 6 dimerized upon irradiation in methanol to give the head-to-tail dimers (7, 45% yield, mp 226-227°C and 8, 25% yield, mp 199-201°C).

