

SYNTHESIS OF SOME  $\beta$ -D-RIBOFURANOSYL-4,7-METHANOINDAZOLES AND  
 PYRAZOLO[1,5-*a*]AZEPINES

Shin-ichi NAGAI and Jinsaku SAKAKIBARA

Faculty of Pharmaceutical Sciences, Nagoya City University,

3-1 Tanabe-dori, Mizuho-ku, Nagoya, 467, Japan

We have prepared a series of optically active (4*S*,7*R*)-7,8,8-trimethyl-4,5,6,7-tetrahydro-4,7-methano-1*H*-indazoles and elucidated their appreciable pharmacological activities. Since it is considered that the introduction of  $\beta$ -D-ribofuranosyl group into the pyrazole ring of 4,7-methano-1*H*-indazole might bring about interesting biological activity change, we report the convenient synthesis of a new class of pyrazole N- and C- $\beta$ -D-ribofuranosides.

Condensation of 2,3-*O*-isopropylidene-D-ribofuranosylhydrazine with (1*R*,4*S*)-3-hydroxymethylenebornan-2-ones led to selective N-1 ribosylation of pyrazole ring to provide (4*S*,7*R*)-1-(2,3-*O*-isopropylidene- $\beta$ -D-ribofuranosyl)-4,7-methano-1*H*-indazoles (1*a*-*b*), and corresponding deprotected 4,7-methano-1*H*-indazole after treatment with acid. Structure determination including anomeric configuration assignment was discussed based on <sup>1</sup>H-NMR spectroscopy.

1,3-Dipolar cycloaddition of (1*R*,4*S*)-3-diazobornan-2-one with methyl 3-(2,3-*O*-isopropylidene-5-*O*-trityl- $\beta$ -D-ribofuranosyl)propiolate followed by [1,5]sigmatropic rearrangement was used as a key reaction step in a novel synthesis of pyrazole C-ribofuranoside; (4*S*,7*R*)-3-(2,3-*O*-isopropylidene-5-*O*-trityl- $\beta$ -D-ribofuranosyl)-8-oxo-4,7-methano-8*H*-pyrazolo[1,5-*a*]azepine (2). The chemical structure and the binding position of  $\beta$ -D-ribofuranosyl group were confirmed. The isopropylidene and trityl groups were easily removed to give (4*S*,7*R*)-8-oxo-3-( $\beta$ -D-ribofuranosyl)-4,7-methano-8*H*-pyrazolo[1,5-*a*]azepine.

