REACTIVITY OF OXYGENATED 5-DEAZAFLAVINS

Fumio Yoneda,[a] Yoshiharu Sakuma,[a] and Masayuki Takamoto[b]
Faculty of Pharmaceutical Sciences, Kumamoto University,[a] Oe-honmachi, Kumamoto 862, Japan and Central Research Division, Takeda Chemical Industries, Ltd.,[b] Juso-honmachi, Yodogawa-ku, Osaka 532, Japan

Recently, we found that 5-deazaflavin-dependent oxidation of alcohols and amines to carbonyl compounds is automatically recycled under weakly basic conditions.[1] Hereupon, 1,5-dihydro-5-deazaflavins (II) initially formed are reoxidized to the original 5-deazaflavins (I) by adventitious oxygen (air), while oxygen may be hydrogenated to hydrogen peroxide. It may be possible that the hydrogen peroxide thus formed will partially catch the 1,5-dihydro-5-deazaflavin (II) as well as the 5-deazaflavin (I) in the autorecycling system to give birth to the corresponding oxygenated 5-deazaflavins.

Thus we have carried out the oxidation of I and II with m-chloroperbenzoic acid (which could be considered as an equivalent of hydrogen peroxide) in chloroform to afford 4a,5-epoxy-5-deazaflavin[2,3] (III) and 5-hydro-4a-hydroxy-5-deazaflavin (IV) respectively. The reactivities of these oxygenated compounds toward several kinds of reagents were examined, which may serve as a basis for understanding reactivity of the 5-deazaflavin cofactor (e. g. coenzyme F_{420}) in biological system and also providing new revelant preparative procedure in synthetic heterocyclic chemistry.



1) F. Yoneda, et. al., J. Chem. Soc. Perkin Trans. 1, 1981, 1836.
2) D. Vargo and M. S. Jorns, J. Am. Chem. Soc., 1979, 101, 7623.
3) F. Yoneda, Y. Sakuma, Tetrahedron Letters, 1981, 3977.