

TYPICAL ELECTROPHILIC REACTIONS OF FUROPYRIDINES

Shunsaku Shiotani and Hiroyuki Morita

College of Liberal Arts, University of Toyama, Gofuku 3190, Japan

In order to reveal the reactivities of furopyridines, we have undertaken the bromination and the nitration of furo[2,3-b]-, (1), furo[3,2-b]-, (2), furo[2,3-c]-, (3) and furo[3,2-c]pyridine (4) whose chemical properties had been almost unknown.

Bromination of 1, 2, 3 and 4 with 3 moles of Br_2 in CCl_4 afforded the corresponding trans-2,3-dibromo-2,3-dihydro derivatives (5, 6, 7 and 8) in 95, 92, 20 and 52% yield, respectively, which were derived to 3-bromofuropyridines (9, 10, 11 and 12) by treatment with NaOH in MeOH- H_2O in almost quantitative yield.

Nitration of 1 with a mixture of fuming HNO_3 and H_2SO_4 afforded a mixture of cis-, (13) and trans-2-nitro-2,3-dihydrofuro[2,3-b]pyrid-3-yl nitrate (14) and 2-nitrofuro[2,3-b]pyridine (15). Both 13 and 14 were easily converted to 15 by treatment with NaHCO_3 solution. Compound 2 was nitrated to give a mixture of cis-, (16) and trans-2,3-dihydrofuro[3,2-b]pyridin-3-ol (17) and 2-nitro derivative (18). Compound 16 was transformed to 17 by refluxing on silica gel in AcOEt. The trans isomer 17 was dehydrated with Ac_2O to give 18. Reaction of 3 with a mixture of fuming HNO_3 and H_2SO_4 gave a nitrolic acid derivative 19 in 46% yield. Nitration of 4 gave a mixture of 3-(trinitromethyl)-pyridin-4-ol (20) (yield: 30%) and 2-nitrofuro[3,2-c]pyridine (21) (yield: 32%).

The structure of 19 and 20 were established by X-ray analysis. The differences of behavior observed in these reactions are discussed in connection with the results of the determination of basicities and the relative reactivities of H-D exchange reaction of these furopyridines.

