

REACTION OF 7-METHOXY-1-PHENYL-1H-1,2,3-TRIAZOLO[4,5-d]PYRIDAZINE 5-OXIDE WITH NUCLEOPHILES

Etsuo Oishi, Yoshikazu Asahina and Eisaku Hayashi

Shizuoka College of Pharmacy
2-2-1, Oshika, Shizuoka 422, Japan

As a part of series of studies on the chemistry of the condensed pyridazines, chemical properties of 7-methoxy-1-phenyl-1H-1,2,3-triazolo[4,5-d]pyridazine 5-oxide(II), obtained in the reaction of 7-methoxy-1-phenyl-1H-1,2,3-triazolo[4,5-d]pyridazine(I) with m-chloroperbenzoic acid, were studied.

II reacted with nucleophiles to give 4,7-disubstituted 1-phenyl-1H-1,2,3-triazolo[4,5-d]pyridazines or 4,5-disubstituted 1-phenyl-1H-1,2,3-triazoles.

Nucleophiles[Products] ; phosphoryl chloride[III], thionyl chloride, sulfuryl chloride[III and IV], tosyl chloride[IV, V and VI(C₁₈H₁₆ClN₃O₄S)], phenyl isocyanate[VII and a mixture of structure unknown products], sodium hydroxide[VIII], acetophenone in the presence of sodium methoxide[VIII and IX], methylmagnesium iodide[Xa], ethylmagnesium bromide[Xb], acetic anhydride[XI and XII], acetyl chloride[XIIIa], benzoyl chloride[XIIIb], benzoyl chloride and potassium cyanide(V, XIV(C₂₂H₁₈N₆O₄) and XV(C₁₈H₁₅NO₄)), dimethyl acetylenedicarboxylate[XVI and XVII], methyl propiolate[XVIII and XIX].

In this lecture, we would show the reaction mechanism for the formation of the products.

