


PREPARATION AND THERMOLYSIS OF 1,2,3,6-TETRAHYDRO-1,2-DIPHENYL-
1^λ,2^λ-DIPHOSPHORIN 1,2-DISULFIDE


Takayuki KAWASHIMA, Michio SHIMAMURA, and Naoki INAMOTO

Department of Chemistry, Faculty of Science, The University of
Tokyo, Hongo, Bunkyo-ku, Tokyo 113, Japan


The titled compound (1) was prepared in 53% yield by reaction of 1,2-diphenyl-1,2-dipotassiodiphosphane (2) with *cis*-1,4-dichloro-2-butene, followed by sulfurization. *Trans* and *cis* isomers were separated by dry column chromatography on silica gel in a ratio of 8:1.



Trans-1 was refluxed in dodecane for 5h with 2,3-diphenyl-1,3-butadiene (3) to give *trans*-1,2,3,6-tetrahydro-1,2,4,5-tetraphenyl-1^λ,2^λ-diphosphorin 1,2-disulfide (4) and 1,3,4-triphenyl-3-phospholene 1-sulfide (5) in 15 and 50% yields, respectively.



On the other hand, *cis*-1 was heated at 210°C for 1h to afford *trans*-1, *trans*-4, 5, and 3,6-dihydro-2,4,5-triphenyl-2H-1,2^λ-thiaphosphorin 2-sulfide (6) in 12, 11, 32%, and trace yields, respectively. Temperature dependency was observed in a ratio of 5 to 6.



The possible mechanism is discussed.