

ADVANCES IN THE CHEMISTRY OF THE FURANODITERPENOIDS FROM TEUCRIUM SPECIESFranco Piozzi^{*a}, Benjamin Rodriguez^b, and Giuseppe Savona^a^aInstitute of Organic Chemistry, Palermo University,
Archirafi 20, 90123 Palermo, Italy^bInstituto de Quimica Organica General, C.S.I.C.,
Juan de la Cierva 3, 28006 Madrid, Spain

Abstract - This review describes the major advances reported during the last five years on the chemistry of these diterpenoids.


Six years elapsed since the publication of a review¹ on the chemistry of the furanoclerodane diterpenoids occurring in several species of genus Teucrium (family Labiatae). During this time, some research groups were active in this field, and many more species were investigated. The present paper aims at giving an up-to-date review on the results obtained after June 1980. It must be focused that specimens of the same botanical species, collected in different countries, showed a quite dissimilar qualitative content of diterpenoids. When the existence of subspecies is not recognized, these differences will be ascribed tentatively to the occurrence of chemotypes. As to the structure, all the new diterpenoids here reported do have the ent-clerodane skeleton (synonymous, neo-clerodane), like the products previously¹ reported. The nomenclature ent-clerodane or neo-clerodane is used indifferently through this paper.

RECENT RESEARCHES

This report starts with the substances whose isolation had been announced in the previous review¹.

It had been reported¹ the occurrence of auropolin in T. polium subsp. aureum (Schreber) Arcangeli (= T. aureum Schreber), Spanish chemotype, but its structure was not ascertained. The elucidation of the structure as [25] showed² that the usual C-20 → C-12 lactone or hemiacetal bridge is missing, and substituted by a C-20 → C-7 hemiacetal bridge. The absolute stereostructure of auropolin, with C-12 in S configuration, was proved, except for its C-20 centre, by X-ray determination on the product of oxidation of [25], having the C-20 → C-7 lactone bridge.

The structure of teugin from T. fragile Boiss. had been given¹ provisionally: it



ent-clerodane skeleton
and numbering

was subsequently completed³ and confirmed, by placing the second hydroxy group on C-6 in axial β -orientation and recognizing the configuration of the first hydroxy group on C-2 as β axial. Therefore the structure of teugin is depicted by [77].

A related derivative, dihydroteugin [78], was found⁴ in T.chamaedrys L., Spanish chemotype. A product with the same structure was obtained by NaBH_4 reduction of teugin; the hydrogen atom on C-4 has β -orientation (pseudoaxial). Recently, dihydroteugin was ascertained⁵ to be identical

with teucrin B, isolated by Popa⁶ from the Moldavian chemotype of T.chamaedrys. The structure given by Popa⁶ and quoted in the previous review¹ was incorrect, because the hydroxy groups are not on C-1 and C-7 as claimed⁶.

The Spanish chemotype of T.chamaedrys contained several more diterpenoids of quite unusual structure. Chamaedroxide was found⁷ to have structure [88] with an oxetane bridge between C-4 and C-6, as confirmed inter alia by X-ray analysis. Also teucroxide [47] has an oxetane ring, connecting C-19 and C-4, as in montanin D [46]¹; the structure of teucroxide was elucidated⁸ by determination of the natural-abundance ^{13}C - ^{13}C coupling constants observed via double quantum coherence. It can be remembered that the occurrence of oxetane rings is very rare in natural products. The same Spanish chemotype of T.chamaedrys yielded^{4,9} six already known diterpenoids: the norclerodanes teuflin [93]¹, teuflidin [95]¹, teucrin A [97]¹, and the clerodanes teucrin E [72]¹, teugin [77]³ and teuchamaedryn B (see below), the latter identical with teucrin H2 [76] from T.hyrceanicum¹. Two novel norclerodanes, 6-epi-teucrin A and isoteuflidin were found^{9,10}, and their structures were assigned respectively as [98] and [96]. Isoteuflidin has therefore reversed the configurations at C-6 and C-10 of teuflidin. The occurrence of the two stereoisomeric couples teucrin A/6-epi-teucrin A and teuflidin/isoteuflidin in the same plant is remarkable.

An Italian chemotype of T.chamaedrys yielded¹⁰ six already known¹ diterpenoids: the norclerodanes teucvin [89], teucvidin [91], teuflin [93], teucrin A [97], and the clerodanes teucrin F and teucrin G. The last two products had been isolated⁶ by Popa from the Moldavian chemotype of T.chamaedrys and attributed⁶ incorrect structures¹ that were now¹⁰ amended. The products are represented by [74] and [75]: here again the hydroxy group is on C-6 and not on C-7 as claimed by Popa. Concerning the Moldavian chemotype, the existence and structures of five products were therefore confirmed: teucrin A [97], teucrin B (= dihydroteugin)[78], teucrin

E [72], teucrin F [74] and teucrin G [75]. Nothing more was published by Popa about the so-called teucrin C and teucrin D he claimed to have isolated: it is possible that they are identical with products extracted from the other chemotypes or from other species of Teucrium.

A Bulgarian chemotype of T.chamaedrys gave¹¹ the known¹ teucrin A [97] and teucrin E [72]. A third product named teuchamaedryn A but identical with the known¹ teuflin [93] was found, and correlated¹¹ with montanin B [102] (from T.montanum¹). Another substance¹¹, teuchamaedryn B, is identical with teucrin H2 [76] (from T.hyrceanicum¹). Recently¹² the same research group isolated also dihydroteugin [78] (= teucrin B) and 6 α -hydroxyteuscordin [79] (see below for T.scordium). A new product, teuchamaedryn C [64], is interesting¹² for the occurrence of a δ -lactone bridge C-20 $\rightarrow$ C-19 and of an hemiacetal system linking C-18 and C-6.

An always open problem is concerned with T.polium, because many subspecies are reported in the taxonomical literature; also the occurrence of different chemotypes is probable, as in the case of T.polium subsp. aureum¹.


T.polium (L.) subsp. capitatum Arcangeli (= T.capitatum Arcangeli) contains, besides the three previously reported¹ clerodanes picropolin [1], picropolinone [4] and 19-acetylgnaphalin [22], three new products more^{13,14}: capitatin [5], teucapitatin [7] and lolin [48]. Structures and absolute configurations of capitatin and lolin were firmly established also by X-ray diffraction.

Recently, the examination of a new sample of the same subsp. capitatum gave different results¹⁵. The occurrence of picropolin, picropolinone and 19-acetylgnaphalin was confirmed, but no traces of capitatin, teucapitatin and lolin were detected. On the contrary, the previously known teucjaponin B [19] (see below for T.japonicum) was isolated, together with three new compounds: 7-deacetylcapitatin [6], picropolinol [49] and 20-epi-isoseriocephalin [17].

The first product was easily correlated with capitatin, picropolin and picropolinone. NOE experiments and CD curve finally confirmed the 12S and neo-clerodane configurations. The second product differs from picropolin by the opening of the 4,18-epoxide ring to form a 4 α -OH, 4 β -CH₂OAc system. Indeed, treatment of picropolin with acetic acid yielded picropolinol [49]. The third compound is the C-20 epimer of isoseriocephalin [16] (see below for T.lanigerum), whose structure and absolute stereochemistry were firmly established. Careful NOE experiments proved the 20R and 12S configurations, while the CD curve confirmed the neo-clerodane stereochemistry.

From T.polium subsp. polium collected in Bulgaria, besides the five already¹ reported diterpenes, three new products were isolated^{16,17}. Teupolin III [50] has a bicyclic acetal system including a five-membered C-20 $\rightarrow$ C-12 and a six-membered C-20 $\rightarrow$ C-19 ring, as in teucrin P₁ from which it differs only by the fact that

the C-4 $\rightarrow$ C-18 epoxide ring is open. Teupolin IV [11] has the same structure of gnaphalin [21] but has a 7 β -OAc group more. Teupolin V [65] is interesting for the occurrence of a C-18 $\rightarrow$ C-6 ether bridge and of two tertiary hydroxyls on C-4 and C-6. The biogenesis of this last product from teucrin P₁ is proposed¹⁷ to occur as reported in the following schema; the transformation of teucrin P₁ [24] into teupolin V [65] was performed also *in vitro*¹⁷ by treatment with H₃PO₄. It can be remarked that the elimination of C-19 by retroaldolic cleavage¹ is prevented here by the occurrence of the C-19 $\rightarrow$ C-20 ether bridge.



Recently the structure of teupolin I, previously given by the Bulgarian group and reported in our review¹, had to be amended¹⁸ to [9] by means of ¹H, ¹³C and NOE studies: i.e., the configuration at C-12 is *R* and not *S* like in the most of the clerodanes from Teucrium: this was the first, but not the last case, of a "reversed" configuration at C-12.

At last, Papanov and Malakov¹⁹ quoted also the occurrence of montanin E [51] (see below for T. montanum) in their sample of T. polium.

The extraction of T. japonicum Houtt. yielded²⁰, besides the well known teucvin [89], two new products, the C-6 epimers teucjaponin A [18] and teucjaponin B [19]. By CrO₃ oxidation, [18] was converted into 19-acetylgaphalin [22], thus confirming the absolute stereochemistry. Teucjaponin B [19] is identical with the product obtained²¹ by NaBH₄ reduction of 19-acetylgaphalin [22] and is therefore different from montanin C (see below).

Other results are concerned with T. montanum. Whereas T. montanum subsp. montanum

collected in Northern Italy and in Sicily does not contain any diterpenes (see below), T. montanum subsp. skorpilii growing in Bulgaria had been already reported¹ to contain four clerodanes, montanin A [101], B [102], C and D. The controversy on the structure of montanin C was eventually solved: indeed, it was proved¹⁸, also by X-ray analysis, that its configuration at C-12 is R (like in teupolin I) and not S as previously claimed²²; so montanin C is the C-12 epimer of teucjaponin B and must be represented by [8]. The stereostructure of montanin D, not previously given²³, was assigned as [46] by NMR studies²².

Two further clerodanes were isolated¹⁹ from T. montanum subsp. skorpilii, montanin E and montanin F. The latter is identical with teucjaponin A [18]. Montanin E is a highly hydroxylated product (two primary, one secondary, one tertiary OH groups) and was attributed the structure [51].

T. scordium L. growing in Bulgaria was the object of several investigations during 1980-1984, and it was the source of five new clerodane diterpenoids: teuscordinon [82]²⁴, the two epimers 6 α -hydroxyteuscordin²⁵ [79] and 6 β -hydroxyteuscordin²⁶ [80], 6-ketoteuscordin²⁵ [84] in which the 3,4-double bond is saturated, and 2-keto-19-hydroxyteuscordin²⁷ [71]. It is unfortunate for the nomenclature that the identical termination "teuscordin" is used for derivatives of differently saturated rings, as [79] and [84], and even for different skeletons, as [79] and [71]. Also two known¹ products were isolated²⁷, teucrin E [72] and teucrin H4 [94]; recently¹⁹ also the occurrence of montanin E [51] was reported. A further product, having structure [77] and called²⁶ 2 β ,6 β -dihydroxyteuscordin, is however identical with teugin³.

The structure [84] here reported represents 6-ketoteuscordin correctly²² with the equatorial 8 α -CH₃ configuration, and not with the previous²⁵ uncorrect axial 8 β -CH₃ orientation. The stereochemistry at C-8 was amended²² as 8 α -CH₃ also in the case of teucrin H2 [76], isolated from T. hyrcanicum¹.

Several of the products hitherto reported show a 4,18-epoxide ring, or arise from it by hydrolytic opening, or have undergone a subsequent ring closure between C-18 and C-6 or C-19 through an oxygen atom. In the case of T. africanum Thunb. from South Africa, the epoxide ring was open by chloride ions, producing two unprecedented chlorinated clerodanes²⁸, tafricanin A [52] and B [53]. It was proved that both substances are true natural products and not extraction artifacts. The structure and absolute configuration of [52] were confirmed by X-ray diffraction. When the chlorohydrins [52] and [53] were heated in the presence of Amberlite IR-400 resin, they yielded the related epoxides; treatment of the epoxide from [52] with concentrated hydrochloric acid and chloroform regenerated the parent tafricanin A. Another rich source of products is T. flavum L. subsp. glaucum (Jordan and Fourr.) Ronniger, harvested in Sardinia; this subspecies is different from T. flavum subsp.

flavum collected in Sicily and previously proved¹ to contain teuflin [93] and teuflidin [95]. Subsp. glaucum contains, besides teuflin²⁹ [93], the new diterpenes 12-epi-teucvin¹⁸ [90], teuflavin²⁹ [33], and a novel glucoside named teuflavoside [103]²⁹.

The structure of 12-epi-teucvin [90] shows the unusual R configuration at C-12, like in teupolin I [9] and montanin C [8] (see before).

The structure [33] of teuflavin was confirmed by CrO₃ oxidation to the epoxyderivative obtained from tafricanin A [52] (see before). This correlation also proved the ent-clerodane absolute configuration. For the configuration at C-20, not given in this paper²⁹, see a subsequent communication³⁴.

Teuflavoside [103] is the first case of a clerodane diterpene involved in a glucoside structure. It contains two acetyl groups; the bisdeacetyl derivative gave D(+)-glucose and montanin B [102]¹ by enzymic hydrolysis with β -(D)-glucosidase. The first acetyl group occurs on the primary hydroxyl at C-18 of montanin B, the second on the hydroxyl at C-2' of glucose. Therefore teuflavoside is 18-acetylmontanin B 6-2'-O-acetyl- β -D-glucopyranoside.

A sample of T.marum L., also collected in Sardinia, yielded³⁰ a new clerodane, teumarin [34]. However, several years ago we had found³¹ that a sample of the same species harvested in Eastern Spain did not contain any diterpene. Teumarin has an axial 2 β -OH group; its absolute configuration was proved by the CD curve of its 6-keto derivative (negative Cotton effect like in 19-acetylgnaphalin¹).

A third species collected in Sardinia, T.massiliense L., gave five diterpenoids³². Two of them, teucjaponin A [18] and montanin C [8], were already known, but the other three are novel and interesting for several details of their structures. The first new product was proved to have the structure [37] of deacetylajugarin II: it lacks the 20,12-lactone system, carbon C-20 is a methyl, there is no oxygen function on C-12; moreover, the furan ring is oxidized to an α,β -unsaturated γ -lactone ring, like in the ajugarins; this last feature could suggest a taxonomic similarity of T.massiliense with the genus Ajuga. The absolute configuration of [37] was proved by transformation into the well known ajugarin I.

The second and third new diterpenes³², teumassilin [35] and 6,19-diacetylteumassilin [36], do have the furan group and a hydroxy function on C-12, but lack the 20,12-lactone system because carbon C-20 is still a methyl group. Also their absolute configurations were proved.

Three new diterpenoids were extracted³³ from T.pyrenaicum L., teupyrenone [32], teupyreinin [28] and teupyreinidin [29]. A common feature is the occurrence of an equatorial 3 β -OAc group; it is noteworthy that all their hydroxy groups are acetylated; differences are given by the presence of the γ -lactone in [28], of the hemiacetal in [29], of the acetal system C-20/C-12 and C-20/C-19 in [32].

The structures [28] and [29] here reported differ from the ones given in our first paper³³. Indeed, quite recently³⁴ our group proved that teupyrein has the unusual R configuration at C-12 instead of S configuration; moreover, the configuration of teupyreinidin at C-20 is R and not S.

In the same paper³⁴ we were able to assign the configuration at C-20 to teuflavin as [33], configuration that had not been given in the first²⁹ paper.

These results were obtained by NOE experiments. Indeed this paper reports³⁴ conclusive evidence on the potentiality of this technique for the assignment of the absolute configuration at C-12 in products having a C-20 $\rightarrow$ C-12 γ -lactone ring, and at C-20 in products having the C-20 $\rightarrow$ C-12 hemiacetal system.

In fact, irradiation of the protons of the secondary 8α -CH₃ group produces a 8-12 % NOE enhancement of H-12 when these protons are on the same side of the plane defined by the lactone ring (configuration 12R). This enhancement is missing when the configuration is 12S, but in this case a 2-6% effect for the furanic H-14 and, sometimes, a 1-5% enhancement for the furanic H-16 are observed.

Analogously, irradiation of 8α -CH₃ in ent-clerodanes having the C-20 $\rightarrow$ C-12 hemiacetal function affords a strong (10-15%) NOE enhancement of H-20 when this proton and 8α -CH₃ are on the same side of the plane defined by the hemiacetal ring; this effect is missing when H-20 and 8α -CH₃ are on opposite sides of the plane.

Also the irradiation of H-20 can be diagnostic for the configuration at C-12: a 5% NOE effect was observed³⁷ for H-12 in the case of 20-epi-teulanigin [42], (see below for T.lanigerum), where H-20 and H-12 are on the same side of the hemiacetal ring.

A recent reinvestigation³⁵ of T.pyrenaicum led to the isolation of two minor constituents, teupyrin A [26] and teupyrin B [27].

The first product lacks the usual γ -lactone or γ -lactol ring between C-20 and C-12; indeed, C-12 bears an acetoxyl group and C-20 forms a hemiacetal system with C-19. All NMR data were consistent with structure [26], and NOE experiments proved R configuration at C-20. Finally, X-ray analysis established the absolute stereochemistry of the neo-clerodane skeleton, of C-12 as S and of the 3β -OH group.

The second product is similar to teumassilin [35] but has a 3β -OH group more; this is another rare case in Teucrium of the existence of C-20 as a methyl group. The stereochemistry at C-12 is not established, while the ent-clerodane configuration is supposed, like all the other diterpenoids from Teucrium.

A species exceedingly rich in diterpenes, both from quantitative and qualitative points of view, is T.lanigerum Lag. (synonymous T.eriocephalum Wk. subsp. rubrifolium Coincy). In fact, the dry aerial parts contain up to 1.7% of a mixture of at least ten products. Two of them were identified as known¹ substances: the first is¹⁸ teupolin I [9], the second is³⁶ eriocephalin [13], occurring in high amount

(1.11% from dry material) and previously found¹ in T.eriocephalum. Two new products, quite related to eriocephalin, were attributed³⁶ the structures of 20-deacetyleriocephalin [14] and isoeriocephalin [16].

Amongst the minor constituents of T.lanigerum we obtained³⁷ 7,8-dehydroeriocephalin [15] and an unseparable mixture of natural diterpenes that could be resolved only after acetylation: five derivatives, not previously described as natural or synthetic substances, were isolated after careful chromatography. They are³⁷ teulanigeral [43], teulanigin [41], 20-epi-teulanigin [42], teulanigerin [66] and teulanigeridin [67].

Teulanigeral [43] is remarkable for the occurrence of C-20 as a free aldehyde group: therefore there is neither the γ -lactone nor the hemiacetal ring between C-20 and C-12; its absolute configuration at C-12 was not ascertained, but all the other products from T.lanigerum have *S*-configuration at C-12.

The configurations at C-20 of teulanigin [41] and 20-epi-teulanigin [42] were proved by NOE experiments, and the absolute neo-clerodane configuration relies on their CD curves (negative Cotton effect).

Teulanigerin [66] has two equatorial tertiary hydroxy groups on C-4 and C-6, and a C-18 $\rightarrow$ C-6 ether bridge, already observed¹⁷ in teupolin V [65]. This bridge occurs also in teulanigeridin [67]: the substance shows also a rare orthoacetate system with the 4 α ,6 α ,19-hydroxy groups. All the C-12 and C-20 configurations of [66] and [67] were proved by NOE experiments; the absolute configuration, although not ascertained, is believed to belong to the neo-clerodane series like the other compounds.

Since compounds [41], [42], [43], [66] and [67] were isolated only as acetyl derivatives and, before acetylation, the ¹H NMR spectrum of the mixture of diterpenes was devoid of acetoxyl signals, it is clear that they occur in T.lanigerum as the corresponding deacetylated compounds. Finally, it is important to note that the ¹H NMR spectrum of the natural mixture of diterpenoids showed the signal corresponding to the orthoacetate methyl group of teulanigeridin [67], that therefore is not an artifact.

Another species, widespread in Western Europe, is T.scorodonia L. Samples of the subsp. scorodonia, harvested in Northern Spain, yielded^{38,39} seven compounds, only one of them being already known¹, i.e. teupolin I [9], whereas the other six are new natural products.

Teuscorolide [99] is a 19-norclerodane with the γ -lactone ring between C-18 and C-6, and an unusual 4,6-diene system. It was correlated³⁸ with the well known¹ teucrin A by dehydration of the latter by acetic anhydride - sodium acetate treatment. A close derivative, 2 α -hydroxyteuscorolide [100], was also found³⁹ in this subspecies.

Teuscorodal [54] is a 3-dehydro derivative with C-18 as an aldehyde group. Its absolute configuration was proved³⁸ by application of Horeau's method. Its reduction product, teuscorodol [55], cooccurs in the subspecies³⁸, and both compounds were transformed the one into the other.

Teuscorodin [85] is the first diterpenoid from Teucrium having a hemiacetal group at C-18 bonded with C-19. Its structure and absolute configuration were proved³⁹ by oxidation to 6-ketoteuscordin [84] and by its CD curve (negative Cotton effect). Teuscorodonin [68] is a 3-dehydro derivative with a γ -lactone ring between C-18 and C-6, while C-19 exists as a carbinol group. Its structure was proved³⁹ by careful NMR investigation, the 12S-configuration by NOE experiments and the absolute neo-clerodane stereochemistry by its CD curve.

From some other species of Teucrium growing in the Iberian peninsula, several new products were isolated recently.

The reinvestigation⁴⁰ of T.gnaphalodes L'Hér. yielded, besides the four already¹ reported products, a new diterpenoid, teugnaphalodin [69]. This substance is interesting for the occurrence of a free hydroxy group on C-12, as in teumassilin [35] and teupyrin B [27], of the ether bridge C-18 $\rightarrow$ C-6, and of a δ -lactone ring C-20 $\rightarrow$ C-19. The absolute configuration at C-12 was proved to be 12S by application of Horeau's method.

The extraction of T.webbianum Boiss. gave⁴¹ the new 2 β -hydroxyteucvidin [92] together with the previously known¹ teuflidin [95] and teucrin A [97]. It can be observed that all the three compounds are 19-norclerodanes.

From T.salviastrum Schreber six new products, teusalvins A to F, and the previously known teucvidin¹ [91] and teucroxide⁸ [47] were isolated⁴². Teusalvin A [86] has a hemiacetal function C-18 $\rightarrow$ C-19, like teuscorodin [85], and was correlated with dihydroteugin [78] whose structure is known⁴; the configuration at C-12 was confirmed to be 12S by NOE experiments.

Also teusalvin B [87], a reduction derivative of teusalvin A, was correlated with dihydroteugin; the 12S-configuration and the neo-clerodane stereochemistry were determined by NOE experiments and by CD curve respectively.

Teusalvin C [57] is similar to teuscorodol [55]: apart the absence of the acetyl group on C-19, the most striking difference is the 12R-configuration, as proved by NOE enhancement of proton H-12 when irradiating the 8 α -CH₃ group. All the other teusalvins have on the contrary the 12S-configuration. So teusalvin C is the fifth neo-clerodane diterpenoid from Teucrium to have 12R-configuration, together with montanin C, teupolin I, 12-epi-teucvin and teupyreinin.

The isomeric teusalvin D [58] and teusalvin E [59] can be partially transformed the one into the other by treatment of their methanolic solution with a trace of sodium carbonate. The first product is a 20,12- γ -lactone, while the second is a

20,19- δ -lactone. Both products have 12*S*-configuration.

Finally, the structure of teusalvin F [60] was confirmed by X-ray diffraction, in full agreement with all NMR data. The unprecedented (in *Teucrium*) ether bridge can arise from teusalvin D by an intramolecular nucleophilic attack of the C-19 hydroxy group at the C-2 position.

The investigation of *T.lepicephalum* Pau led to identify⁴³ three new compounds: teulepicin [44], 19-acetylteulepicin [45] and teulepicephin [70]. The first two products gave NMR data quite similar to the ones reported for the pair¹ gnaphalin [21] and 19-acetylgnaphalin [22]. Indeed, the only difference is the presence of an additional 3 β -OH group in [44] and [45]. The *ent*-clerodane configuration and the 12*S*-stereochemistry were ascertained by CD and NOE measurements respectively. Furthermore, [44] and [45] were correlated mutually and with a derivative of tafricanin A [52]. Another reliable result for the structure of teulepicin is the easy transformation into a C₁₉ furanic derivative, as observed previously for gnaphalin¹ [21].

The NMR data of teulepicephin [70] suggest that it is the 3 β -OH homolog of teugnapthalodin [69]: however, the stereochemistry at C-12 and the absolute configuration were not ascertained.

19-Acetylteulepicin [45] was isolated⁴³ also from *T.buxifolium* Schreber together with the well known 19-acetylgnaphalin [22].

T.botrys L. is a species which grows all over Europe; two samples collected in Eastern Spain gave⁴⁴ different results. The first sample contained only the previously known 6 β -hydroxyteuscordin [80]; the second sample yielded five different diterpenoids. Three of them are already known: teucvidin [91], montanin D [46] and teuchamaedryn C [64]. The other two compounds are new, 19-deacetylteuscorodol [56] and teubotrin [61]. The first product was easily correlated with teuscorodol [55]; the 12*S*-configuration was confirmed by NOE experiments, so [56] is the C-12 epimer of teusalvin C [57] (see before for *T.salviastrum*). The second product, teubotrin [61], is a 20,19- δ -lactone: its C-12 stereochemistry and the absolute configuration were not proved.

Recently we investigated⁴⁵ *T.polium* L. subsp. *pilosum* Decsne, widely occurring in Middle East (synonima, *T.pilosum* Aschers.Schweinf., *T.sinaicum* Boiss.). A sample collected in the State of Qatar (Arabian Gulf) yielded the not previously described 19-acetylteupolin IV [12] as the sole component.

The extraction of other species and subspecies led to the isolation of known products only. *T.barbeyanum* Aschers from Cyrenaica (Libya) yielded⁴⁶ teucrin A [97], teucrin F [74] and teucrin G [75]. *T.lucidum* L. from Maritime Alps, Italy, gave⁴⁶ teucvidin [91], teuflin [93], teucrin F [74], teucrin G [75] and 6 α -hydroxyteuscordin [79]. *T.intricatum* Lange from Southern Spain contained⁴⁶ teucvin [89].

T. chartaginense Lange subsp. homotricum Font Quer yielded⁴⁶ eriocephalin [13] and 19-acetylnaphthalin [22]. T. scorodonia L. subsp. euganeum (Vis.) Arcangeli (= subsp. siculum Rafin Guss.) from Sicily gave⁴⁶ teuflin [93]. T. scorodonia L. subsp. scorodonia from Northern Italy contained⁴⁶ teuflin [93], teuscorolide [99] and teuscorodin [85]. From T. subspinosum Willd. four compounds were isolated³⁵: teucvin [89], teuflin [93], teucrin H2 [76] (identical with teuchamaedryn B) and 6 α -hydroxyteuscordin [79]. Picropolinone [4] and 19-acetylnaphthalin [22] were found in T. carolipau C. Vicioso ex Pau subsp. carolipau from Southern Spain⁴⁷. The sole diterpenoid isolated⁴⁸ from T. heterophyllum L'Hér. collected in Canary Islands was teucvidin [91].

As it appears from the present review, the genus Teucrium is a very rich source of neo-clerodane diterpenoids. However, several species were found to be devoid of such products, at least within the limits of our extractive and analytical methods. This is the case of T. lusitanicum Schreber⁴⁷ (from Southern Spain), T. scordioides Schreber (= T. scordium L. subsp. scordioides ex Schreber, Maire et Petitmougin)⁴⁶ T. subtrifidum Lag.⁴⁶, T. pseudochamaepytis L.⁴⁶, T. rotundifolium Schreber (= T. granatense Boiss. et Reuter)⁴⁶, all from Spain, T. apollinis Maire et Weiller⁴⁶, T. davaeanum Coss.⁴⁶, T. cyrenaicum (Maire et Weiller) Brullo et Furnari (= T. polium subsp. cyrenaicum Maire et Weiller)⁴⁶, all from Libya. Also T. flavum L. subsp. flavum collected in Tuscany contains no diterpenoids⁴⁶, whereas the same subspecies growing in Sicily gave¹ teuflin [93] and teuflidin [95]. At last T. montanum L. subsp. montanum of Northern Italy and Sicily contains no diterpenoids⁴⁶.

A recent paper by Bohlmann⁵⁰ reports on the examination of a sample of T. scordium L. collected in West Germany. Thirteen neoclerodanes were isolated, seven of them being already known: teucrin E [72], teugin [77], 6 α -hydroxyteuscordin [79], 6 β -hydroxyteuscordin [80], teuscordinon [82], dihydroteugin [78] (= teucrin B), teucroxide [47]. An eighth compound is 6-acetylteucjaponin B [20], not known as a natural product but previously prepared²¹ by acetylation of teucjaponin B [19]. Five products are new: 2,3-dehydroteucrin E [73], 2 β ,6 α -dihydroxyteuscordin [81], 2 β -hydroxyteuscordinon [83], 6,20-bisdeacetylteupyreinidin [31] and 6-deacetylteupyreinidin [30]. The assignment of the structures relies on exhaustive NMR analysis.

CHEMICAL AND TAXONOMIC REMARKS

Some structural conclusions can be reached at this point. All the diterpenes from the Teucrium genus belong without exception to the ent-clerodane group; all have the furan ring, except deacetylajugarin II in which the ring is oxidized to an α,β -unsaturated γ -lactone system; all have an oxygenated function on C-6 and an equatorial 8 α -CH₃ group. With the exception of teucvin [89] and teucvidin [91] found in two species of Euphorbiaceae¹, all the other products occur only in Teucrium.

In total, 88 C₂₀ and 15 C₁₉ products are known at the present. Different features distinguish groups of products, the most common being the 4,18-epoxide or the 20,12-lactone or the 20,12-hemiacetal system. A minority group lacks the 20,12 bridge system, thus having C-12 as -CHOH- or -CHOAc- or -CH₂-. Few products show C-20 as a methyl group. Only five compounds were proved to have the 12R-configuration, while the 12S stereochemistry is the most common.

The fact that different samples of the same species contain different compounds and that several species contain some identical products, leads us to suggest that the single diterpenes are not adequate for taxonomic purposes as chemical characters of the single species. However, somehow clearer qualitative differences seem to occur in the content of variously oxidized diterpenes, at least between the five sections (*Teucrium*, *Scordium*, *Scorodonia*, *Polium*, *Chamaedrys*) in which the species growing in Europe are divided⁵¹.

CONCLUSIONS

The genus *Teucrium* is still offering opportunities of chemical and taxonomic research. Techniques as ¹³C NMR, NOE, CD and X-ray diffraction are now routine tools. Modest antitumor and insect antifeedant activities have been observed on some pure products or on the raw extracts of some *Teucrium*. The challenge of a total synthesis has not yet been taken up, although significant goals have been scored in the case of the ajugarins and of some derivatives of the neo-clerodane skeleton. No biogenetic study has been reported till now on these diterpenes. A quite hypothetical pathway [A] → [B] → [C] → etc., starting from the *ent*-labdane [A] and linking some of the structures found in *Teucrium*, is reported below tentatively.








TABLE 1 (F = β -furyl)

- [1] picropolin R = H $C_{22}H_{26}O_8$
T.polium¹
T.polium subsp. capitatum¹⁵


- [2] 6-acetyl-picropolin R = Ac $C_{24}H_{28}O_9$
T.polium¹



- [3] isopicropolin $C_{22}H_{26}O_8$
T.polium¹




- [4] picropolinone $C_{22}H_{24}O_8$
T.polium subsp. capitatum^{1,15}
T.carolipau subsp. carolipau⁴⁷




- [5] capitatin R = Ac $C_{24}H_{26}O_9$
T.polium subsp. capitatum¹³

- [6] 7-deacetyl-capitatin R = H $C_{22}H_{26}O_8$
T.polium subsp. capitatum¹⁵




- [7] teucapitatin $C_{24}H_{30}O_9$
T. polium subsp. capitatum¹³




- [8] montanin C $C_{24}H_{30}O_8$ R = Ac
T. montanum subsp. skorpilii¹
T. massiliense³²

- [9] teupolin I $C_{22}H_{28}O_7$ R = H
T. polium subsp. polium¹
T. lanigerum¹⁸
T. scorodonia subsp. scorodonia³⁹




- [10] teupolin II $C_{22}H_{28}O_7$
T. polium subsp. polium¹




- [11] teupolin IV $C_{22}H_{26}O_8$ R = H
T. polium subsp. polium¹⁷

- [12] 19-acetyl-teupolin IV $C_{24}H_{28}O_9$ R = Ac
T. polium subsp. pilosum⁴⁵




- [13] eriocephalin $R = \text{Ac}$ $\text{C}_{24}\text{H}_{30}\text{O}_9$
T.eriocephalum¹
T.lanigerum³⁶
T.chartaginense subsp. homotrichum⁴⁶

- [14] 20-deacetyl-eriocephalin $R = \text{H}$ $\text{C}_{22}\text{H}_{28}\text{O}_8$
T.lanigerum³⁶




- [15] 7,8-dehydro-eriocephalin $\text{C}_{24}\text{H}_{28}\text{O}_9$
T.lanigerum³⁷




- [16] isoeriocephalin $R=\text{OAc}, R'=\text{H}$ $\text{C}_{24}\text{H}_{30}\text{O}_9$
T.lanigerum³⁶

- [17] 20-epi-isoeriocephalin $R=\text{H}, R'=\text{OAc}$ $\text{C}_{24}\text{H}_{30}\text{O}_9$
T.polium subsp. capitatum¹⁵




- [18] teucjaponin A $\text{C}_{22}\text{H}_{28}\text{O}_7$
T.japonicum²⁰
T.massiliense³²
T.montanum subsp. skorpilii¹⁹




- [19] teucjaponin B $R = OH$ $C_{22}H_{28}O_7$
T.japonicum²⁰
T.polium subsp. capitatum¹⁵

- [20] 6-acetyl-teucjaponin B $R = OAc$ $C_{24}H_{30}O_8$
T.scordium⁵⁰




- [21] gnaphalin $R = H$ $C_{20}H_{24}O_6$
T.gnaphalodes¹


- [22] 19-acetyl-gnaphalin $R = Ac$ $C_{22}H_{26}O_7$
T.gnaphalodes¹
T.spinosum¹
T.hyrceanicum¹
T.polium subsp. aureum¹
T.polium subsp. polium¹
T.polium subsp. capitatum¹
T.buxifolium⁴³
T.chartaginense subsp. homotrichum⁴⁶
T.carolipau subsp. carolipau⁴⁷




- [23] gnaphalidin $C_{24}H_{30}O_8$
T.gnaphalodes¹
T.polium subsp. aureum¹




- [24] teucrin P₁ $C_{20}H_{24}O_5$
T.polium subsp. aureum¹
T.polium¹
T.polium subsp. polium¹
T.gnaphalodes¹




[25] auropolin

 $C_{24}H_{30}O_9$ T. polium subsp. aureum²

[26] teupyrin A


 $C_{24}H_{30}O_9$ T. pyrenaicum³⁵

[27] teupyrin B

 $C_{22}H_{32}O_7$ T. pyrenaicum³⁵

[28] teupyreinin


 $C_{26}H_{32}O_{10}$ T. pyrenaicum^{33,34}




[29] teupyreinidin $R^1=R^2=Ac$ $C_{28}H_{36}O_{11}$
T.pyrenaicum^{33,34}

[30] 6-deacetyl-teupyreinidin $R^1=H, R^2=Ac$ $C_{26}H_{34}O_{10}$
T.scordium⁵⁰


[31] 6,20-bisdeacetyl-teupyreinid. $R^1=R^2=H$ $C_{24}H_{32}O_9$
T.scordium⁵⁰




[32] teupyrenone $C_{22}H_{26}O_7$
T.pyrenaicum³³



[33] teuflavin $C_{22}H_{28}O_8$
T.flavum subsp. glaucum^{29,34}




[34] teumarin $C_{22}H_{28}O_8$
T.marum³⁰




- [35] teumassilin $R = H$ $C_{20}H_{30}O_5$
T.massiliense³²

- [36] 6,19-diacetyl-teumassilin $R = Ac$ $C_{24}H_{34}O_7$
T.massiliense³²




- [37] deacetyl-ajugarin II $C_{20}H_{30}O_5$
T.massiliense³²




- [38] fruticolone $R = H$ $C_{22}H_{30}O_6$
T.fruticans¹

- [39] 8β-hydroxy-fruticolone $R = OH$ $C_{22}H_{30}O_7$
T.fruticans¹




- [40] isofruticolone $C_{22}H_{30}O_6$
T.fruticans¹




[41] teulanigin $R^1 = H, R^2 = OAc$ $C_{26}H_{32}O_{10}$
T.lanigerum³⁷

[42] 20-epi-teulanigin $R^1 = OAc, R^2 = H$ $C_{26}H_{32}O_{10}$
T.lanigerum³⁷




[43] teulanigeral $C_{26}H_{32}O_{10}$
T.lanigerum³⁷




[44] teulepicin $R = H$ $C_{20}H_{24}O_7$
T.lepicephalum⁴³


[45] 19-acetyl-teulepicin $R = Ac$ $C_{22}H_{26}O_8$
T.lepicephalum⁴³
T.buxifolium⁴³




[46] montanin D $C_{20}H_{26}O_6$
T.montanum subsp. skorpilii^{22,23}
T.botrys⁴⁴




[47] teucroxide
T.chamaedrys⁸
T.scordium⁵⁰

 $C_{20}H_{26}O_7$


[48] lolin
T.polium subsp. capitatum¹⁴


 $C_{24}H_{28}O_9$


[49] picropolinol
T.polium subsp. capitatum¹⁵

 $C_{20}H_{24}O_8$


[50] teupolin III
T.polium subsp. polium¹⁶

 $C_{20}H_{26}O_6$




[51] montanin E

$C_{20}H_{28}O_7$

T. montanum subsp. skorpilii¹⁹

T. polium subsp. polium¹⁹


T. scordium¹⁹



[52] tafricanin A

$C_{22}H_{25}O_8Cl$


T. africanum²⁸



[53] tafricanin B

$C_{24}H_{29}O_9Cl$


T. africanum²⁸




[54] teuscorodal

$C_{22}H_{26}O_7$


T. scorodonia subsp. scorodonia³⁸




[55] teuscorodol

 $C_{22}H_{28}O_7$ T. scorodonia subsp. scorodonia³⁸

[56] 19-deacetyl-teuscorodol


 $C_{20}H_{26}O_6$ T. botrys⁴⁴

[57] teusalvin C

 $C_{20}H_{26}O_6$ T. salviastrum⁴²


[58] teusalvin D

 $C_{20}H_{26}O_7$ T. salviastrum⁴²




[59] teusalvin E
T.salviastrum⁴²

C₂₀H₂₆O₇




[60] teusalvin F
T.salviastrum⁴²

C₂₀H₂₄O₅



[61] teubotrin
T.botrys⁴⁴

C₂₀H₂₆O₆



[62] teuspinin
T.spinosum¹


R = H

C₂₀H₂₄O₇


[63] 19-acetyl-teuspinin
T.spinosum¹

R = Ac


C₂₂H₂₆O₈




[64] teuchamaedryn C

 $C_{20}H_{26}O_6$ T.chamaedrys¹²T.botrys⁴⁴

[65] teupolin V


 $C_{20}H_{26}O_6$ T.polium subsp. polium¹⁷

[66] teulanigerin

 $C_{24}H_{32}O_9$ T.lanigerum³⁷

[67] teulanigeridin


 $C_{26}H_{32}O_{10}$ T.lanigerum³⁷



[68] teuscorodonin

$C_{20}H_{22}O_6$


T. scorodonia subsp. scorodonia



[69] teugnaphalodin

$C_{20}H_{26}O_7$


T. gnaphalodes⁴⁰



[70] teulepicephin

$C_{20}H_{26}O_8$


T. lepicephalum⁴³



[71] 2-keto-19-hydroxy-teuscordin

$C_{20}H_{22}O_7$


T. scordium²⁷



[72] teucrin E

 $C_{20}H_{24}O_6$


T.chamaedrys^{9,11,1}
T.scordium^{27,50}



[73] 2,3-dehydro-teucrin E

 $C_{20}H_{22}O_6$


T.scordium⁵⁰



[74] teucrin F

 $C_{20}H_{22}O_7$


T.chamaedrys^{10,1}
T.barbeyanum⁴⁶
T.lucidum⁴⁶



[75] teucrin G


 $C_{20}H_{22}O_8$

T.chamaedrys^{1,10}
T.barbeyanum⁴⁶
T.lucidum⁴⁶




- [76] teucrin H2
T.hyrceanicum^{1,22}
T.chamaedrys^{9,11}
T.subspinosum³⁵

C₂₀H₂₄O₆




- [77] teugin
T.fragile³
T.chamaedrys⁴
T.scordium^{26,50}

C₂₀H₂₂O₇




- [78] dihydroteugin (= teucrin B)
T.chamaedrys^{4,5,6,12}
T.scordium⁵⁰

C₂₀H₂₄O₇




- [79] 6α-hydroxy-teuscordin
T.scordium^{25,50}
T.chamaedrys¹²
T.subspinosum³⁵
T.lucidum⁴⁶


C₂₀H₂₂O₆



[80] 6β-hydroxy-teuscordin

 $C_{20}H_{22}O_6$ T.scordium^{26,50}T.botrys⁴⁴

[81] 2β,6α-dihydroxy-teuscordin

 $C_{20}H_{22}O_7$ T.scordium⁵⁰


[82] teuscordinon

R = H

 $C_{20}H_{20}O_6$ T.scordium^{24,50}


[83] 2β-hydroxy-teuscordinon

R = OH

 $C_{20}H_{20}O_7$ T.scordium⁵⁰

[84] 6-keto-teuscordin


 $C_{20}H_{22}O_6$ T.scordium^{22,25}




- [85] teuscorodin $R = H_2$ $C_{20}H_{24}O_6$
T.scorodonia subsp. scorodonia^{39,46}

- [86] teusalvin A $R = O$ $C_{20}H_{22}O_7$
T.salviastrum⁴²


- [87] teusalvin B $R = \alpha-H, \beta-OH$ $C_{20}H_{24}O_7$
T.salviastrum⁴²




- [88] chamaedroxide $C_{20}H_{22}O_7$
T.chamaedrys⁷



- [89] teucvin $C_{19}H_{20}O_5$
T.viscidum subsp. miquelianum¹
T.cubense¹
T.chamaedrys¹⁰
T.japonicum²⁰
T.subspinosum³⁵
T.intricatum⁴⁶




- [90] 12-epi-teucvin $C_{19}H_{20}O_5$
T.flavum subsp. glaucum¹⁸




- [91] teucvidin $R = H$ $C_{19}H_{20}O_5$
T.viscidum subsp. miquelianum¹
T.chamaedrys¹⁰
T.salviastrum⁴²
T.botrys⁴⁴
T.lucidum⁴⁶
T.heterophyllum⁴⁸

- [92] 2 β -hydroxy-teucvidin $R = OH$ $C_{19}H_{20}O_6$
T.webbianum⁴¹




- [93] teuflin $R = H$ $C_{19}H_{20}O_5$
T.flavum subsp. flavum¹, subsp. glaucum²⁹
T.viscidum subsp. miquelianum⁴⁹
T.subspinosum³⁵
T.lucidum⁴⁶
T.chamaedrys^{9,10,11}
T.scorodonia subsp. scorodonia⁴⁶, subsp. euganeum⁴⁶

- [94] teucrin H4 $R = OH$ $C_{19}H_{20}O_6$
T.hyrceanicum¹
T.scordium²⁷




- [95] teuflidin $C_{19}H_{20}O_6$
T.flavum subsp. flavum¹
T.hyrceanicum¹
T.chamaedrys⁹
T.webbianum⁴¹



- [96] isoteuflidin $R^1 = OH, R^2 = H$ $C_{19}H_{20}O_6$
T.chamaedrys¹⁰


- [97] teucrin A $R^1 = H, R^2 = OH$ $C_{19}H_{20}O_6$
T.chamaedrys^{1,4,10,11}
T.barbeyanum⁴⁶
T.webbianum⁴¹



[98] 6-epi-teucrin A

$C_{19}H_{20}O_6$

T. chamaedrys⁹



[99] teuscorolide

R = H

$C_{19}H_{18}O_5$


T. scorodonia subsp. scorodonia^{38,46}

[100] 2α-hydroxy-teuscorolide

R = OH

$C_{19}H_{18}O_6$


T. scorodonia subsp. scorodonia³⁹



[101] montanin A

$C_{19}H_{20}O_4$

T. montanum subsp. skorpilii¹



[102] montanin B $R^1 = H, R^2 = H$

$C_{19}H_{24}O_5$

T. montanum subsp. skorpilii¹

T. polium subsp. polium¹

[103] teuflavoside $R^1 = Ac, R^2 = 2\text{-acetylglucose}$ $C_{29}H_{38}O_{12}$

T. flavum subsp. glaucum²⁹

REFERENCES

1. F.Piozzi, Heterocycles, 1981, 15, 1489.
2. L.Eguren, A.Perales, J.Fayos, G.Savona, M.Paternostro, F.Piozzi and B.Rodriguez, J.Org.Chem., 1981, 46, 3364.
3. M.Bruno, G.Savona, C.Pascual and B.Rodriguez, Phytochemistry, 1981, 20, 2259.
4. G.Savona, M.C.Garcia-Alvarez and B.Rodriguez, Phytochemistry, 1982, 21, 721.
5. M.C.Rodriguez, J.Barluenga, C.Pascual, B.Rodriguez, G.Savona and F.Piozzi, Phytochemistry, 1984, 23, 2960.
6. A.M.Reinbold and D.P.Popa, Khim.Prirod.Soedin., 1974, 10, 589.
7. L.Eguren, A.Perales, J.Fayos, B.Rodriguez, G.Savona and F.Piozzi, J.Org.Chem., 1982, 47, 4157.
8. M.C.Garcia-Alvarez, G.Lukacs, A.Neszmelyi, F.Piozzi, B.Rodriguez and G.Savona, J.Org.Chem., 1983, 48, 5123.
9. F.Fernandez-Gadea, C.Pascual, B.Rodriguez and G.Savona, Phytochemistry, 1983, 22, 723.
10. M.C.Rodriguez, J.Barluenga, G.Savona, F.Piozzi, O.Servettaz and B.Rodriguez, Phytochemistry, 1984, 23, 1465.
11. G.Y.Papanov and P.Y.Malakov, Z.Naturforsch., 1980, 35B, 764.
12. P.Y.Malakov and G.Y.Papanov, Phytochemistry, 1985, 24, 301.
13. C.Marquez, R.M.Rabanal, S.Valverde, L.Eguren, A.Perales and J.Fayos, Tetrahedron Letters, 1980, 21, 5039.
14. C.Marquez, R.M.Rabanal, S.Valverde, L.Eguren, A.Perales and J.Fayos, Tetrahedron Letters, 1981, 22, 2823.
15. P.Fernandez, B.Rodriguez, G.Savona and F.Piozzi, Phytochemistry, 1986, 25, 181.
16. P.Y.Malakov, G.Y.Papanov and J.Ziesche, Phytochemistry, 1982, 21, 2597.
17. P.Y.Malakov and G.Y.Papanov, Phytochemistry, 1983, 22, 2791.
18. J.Fayos, F.Fernandez-Gadea, C.Pascual, A.Perales, F.Piozzi, M.Rico, B.Rodriguez and G.Savona, J.Org.Chem., 1984, 49, 1789.
19. G.Y.Papanov and P.Y.Malakov, Phytochemistry, 1983, 22, 2787.
20. T.Miyase, H.Kawasaki, T.Noro, A.Ueno, S.Fukushima and T.Takemoto, Chem.Pharm.Bull.(Tokyo), 1981, 29, 3561.
21. M.Martinez-Ripoll, J.Fayos, B.Rodriguez, M.C.Garcia-Alvarez, G.Savona, F.Piozzi, M.Paternostro and J.R.Hanson, J.Chem.Soc., Perkin Trans. I, 1981, 1186.
22. E.Gacs-Baitz, M.Kajtar, G.Y.Papanov and P.Y.Malakov, Heterocycles, 1982, 19, 539.
23. P.Y.Malakov, G.Y.Papanov, N.M.Mollov and S.L.Spassov, Z.Naturforsch., 1978, 33B, 1142.

24. G.Y.Papanov, P.Y.Malakov and F.Bohlmann, Phytochemistry, 1981, 20, 170.
25. G.Y.Papanov and P.Y.Malakov, Z.Naturforsch., 1981, 36B, 112.
26. G.Y.Papanov and P.Y.Malakov, Z.Naturforsch., 1982, 37B, 519.
27. G.Y.Papanov and P.Y.Malakov, Phytochemistry, 1985, 24, 297.
28. J.R.Hanson, D.E.A.Rivett, S.L.Ley and D.J.Williams, J.Chem.Soc., Perkin Trans. I, 1982, 1005.
29. G.Savona, F.Piozzi, O.Servettaz, B.Rodriguez, F.Fernandez-Gadea and M.Martin-Lomas, Phytochemistry, 1984, 23, 843.
30. G.Savona, F.Piozzi, O.Servettaz, F.Fernandez-Gadea and B.Rodriguez, Phytochemistry, 1984, 23, 611.
31. Unpublished results.
32. G.Savona, M.Bruno, F.Piozzi, O.Servettaz and B.Rodriguez, Phytochemistry, 1984, 23, 849.
33. M.C.Garcia-Alvarez, J.L.Marco, B.Rodriguez, G.Savona and F.Piozzi, Phytochemistry, 1982, 21, 2559.
34. C.Pascual, P.Fernandez, M.C.Garcia-Alvarez, J.L.Marco, F.Fernandez-Gadea, M. C. de la Torre, J.A.Hueso-Rodriguez, B.Rodriguez, M.Bruno, M.Paternostro, F. Piozzi and G.Savona, Phytochemistry, 1986, 25, 715.
35. P.Fernandez, B.Rodriguez, J.Villegas, A.Perales, G.Savona, F.Piozzi and M. Bruno, Phytochemistry, 1986, 25, 1405.
36. F.Fernandez-Gadea, B.Rodriguez, G.Savona and F.Piozzi, Phytochemistry, 1984, 23, 1113.
37. J.A.Hueso-Rodriguez, F.Fernandez-Gadea, C.Pascual, B.Rodriguez, G.Savona and F.Piozzi, Phytochemistry, 1986, 25, 175.
38. J.L.Marco, B.Rodriguez, G.Savona and F.Piozzi, Phytochemistry, 1982, 21, 2567.
39. J.L.Marco, B.Rodriguez, C.Pascual, G.Savona and F.Piozzi, Phytochemistry, 1983, 22, 727.
40. M.C. de la Torre, B.Rodriguez, G.Savona and F.Piozzi, Phytochemistry, 1986, 25, 171.
41. G.Savona, F.Piozzi, B.Rodriguez, C.Pascual and O.Servettaz, Phytochemistry, 1986, 25, in press.
42. M.C. de la Torre, C.Pascual, B.Rodriguez, F.Piozzi, G.Savona and A.Perales, Phytochemistry, 1986, 25, 1397.
43. G.Savona, F.Piozzi, O.Servettaz, B.Rodriguez, J.A.Hueso-Rodriguez and M.C. de la Torre, Phytochemistry, 1986, 25, in press.
44. M.C. de la Torre, F.Fernandez-Gadea, A.Michavila, B.Rodriguez, F.Piozzi and G.Savona, Phytochemistry, 1986, 25, in press.

45. M.C. de la Torre, F.Piozzi, A.M.Rizk, B.Rodriguez and G.Savona, Phytochemistry, 1986, 25, in press.
46. M.Bruno, F.Piozzi, B.Rodriguez, G.Savona and O.Servettaz, Phytochemistry, 1985, 24, 2597.
47. S.Valverde, unpublished results.
48. B.M.Fraga, A.G.Gonzalez and A.G.Ravelo, unpublished results.
49. M.Node, M.Sai and E.Fujita, Phytochemistry, 1981, 20, 757.
50. J.Jakupovic, R.N.Baruah, F.Bohlmann and W.Quack, Planta Medica, 1985, 341.
51. O.Servettaz, Giornale Bot.Ital., 1985, 119 Suppl.2, 201.

Received, 12th June, 1986