

HETEROCYCLES, Vol. 80, No. 1, 2010, pp. 99 - 102. © The Japan Institute of Heterocyclic Chemistry
 Received, 28th January, 2009, Accepted, 5th March, 2009, Published online, 9th March, 2009
 DOI: 10.3987/COM-09-S(S)7

EFFORTS TOWARD THE SYNTHESIS OF PSEUDOLARIC ACID A: INTRAMOLECULAR BROMOETHERIFICATION AS A MULTIPURPOSE SYNTHETIC TOOL[‡]


Jeremy D. Pettigrew and Leo A. Paquette*

Evans Chemical Laboratories, The Ohio State University, 100 West 18th Avenue,
 Columbus, Ohio, USA, 43210. e-mail: paquette.1@osu.edu

Abstract – Regioselective double bond cleavage within **5**, required for access to the trans-fused perhydroazulene framework of pseudolaric acid A (**1**), has been accomplished via a highly diastereoselective intramolecular bromoetherification process, the latter serving a dual role as a protecting group tactic.

INTRODUCTION

The oxygenated diterpene pseudolaric acid A (**1**),¹ initially discovered to possess antifungal and antifertility properties,² has more recently been shown to exhibit exciting cytotoxic potential.³ Our retrosynthetic conceptualization for construction of the uncommon trans-fused perhydroazulene framework of the compact tricyclic core of **1** relies on the regioselective oxidative cleavage of the more highly strained olefinic linkage in **2**, thereby revealing a direct means for installing the three contiguous tetrahedral carbon centers of **1** (Scheme 1). The precursor **2** was envisioned to originate from the previously described norbornenol **3**.⁴




Scheme 1. Retrosynthetic Analysis of Pseudolaric Acid A (**1**)

[‡] Submitted in celebration of the 80th birthday of Professor Emeritus Akira Suzuki.

RESULTS AND DISCUSSION


Addition of the reagent prepared from 4-bromo-2-methylbut-1-ene⁵ and lithium metal to **3** in THF afforded the corresponding bridged alcohols in satisfactory yield (79%) albeit with moderate selectivity (dr = 5:2 anti:syn) (Scheme 2). The TBS group was removed with TBAF to liberate the corresponding diols, at which point the two diastereomers could be readily separated by column chromatography. Chemoselective oxidation of the anti alcohol with the Dess-Martin periodinane led efficiently to **4**. Conversion of **4** into **5** was accomplished via the Wittig olefination involving the known ylide **6**⁶ followed by relay ring closing metathesis⁷ upon exposure of the resulting triene to the first-generation Grubbs catalyst in 69% overall yield. The selection of **6** instead of $\text{Ph}_3\text{P}=\text{CH}_2$, which would potentially lead to the same desired ruthenium carbene, was made in an effort to avoid ring opening polymerization metathesis of the norbornene moiety by reducing the steric congestion imparted by the bicycle during the initial ruthenium insertion. The correspondence in framework structure residing in **5** and **1** was now made apparent.



Scheme 2. *Reagents and reaction conditions:* (i) 4-bromo-2-methylbut-1-ene, Li, THF, rt, 6 h 79% dr = 5:2 anti:syn; (ii) TBAF, THF, 0 °C to rt, 5 h, 89%; (iii) Dess-Martin periodinane, NaHCO_3 , CH_2Cl_2 , 0 °C to rt, 7 h, 89%; (iv) **6**, Et_2O , -78 °C to rt, 5 h; (v) Grubbs' I, CH_2Cl_2 , reflux, 72 h, 69% over two steps.


The anticipated direct regioselective cleavage of the more strained double bond in **5** proved unrealizable, as the undesired regioselectivity was observed. This unexpected result was attributed to steric crowding by the 7-membered ring in the vicinity of the more strained olefinic center. Consequently, the decision was made to mask this site of unsaturation. The first ploy to be attempted involved the use of an epoxide ring as a protecting group. Thus, alcohol **5** was allowed to react with one equivalent of *m*-CPBA to afford **7** (Scheme 3). As expected, the double bond of the 7-membered ring underwent smooth reaction to afford the single diastereomer **7** which was attributed to the directing ability of the free hydroxyl group. The remaining double bond in **7** was next dihydroxylated with OsO_4 . Cleavage of the resultant diol was accomplished with $\text{Pb}(\text{OAc})_4$ to deliver **9**, which was found to be an unstable compound whose ready degradation is possible by retro aldol reactions similar to the observations made in Trost's reported synthesis of pseudolaric acid B.⁸ To circumvent these retro aldol reactions, the hydroxyl group resident in

7 was protected as the MOM ether, the double bond in **8** was dihydroxylated with OsO₄ and the resultant diol was cleaved with NaIO₄ to afford the stable aldehyde **10**. The attempted direct oxidation of this aldehyde to the corresponding ester with Oxone afforded the mixed acetals **11** or **12** resulting from backside opening of the epoxide by the hemiacetal reaction intermediates.⁹ Formation of the carboxylic acid was next attempted. However, similar backside attack operated to furnish **13**.



Scheme 3. *Reagents and reaction conditions:* (i) *m*-CPBA, NaHCO₃, CH₂Cl₂, 0 °C, 4 h, 79%; (ii) KH, MOMCl, THF, rt, 20 h, 66% (99% BRSM); (iii) OsO₄, NMO, THF, H₂O, 0 °C to rt, 42 h; (iv) NaIO₄, THF, H₂O, rt, 45 min, 48% over two steps *or* Pb(OAc)₄, PhH, rt, 10 min, 47% over two steps (see text); (v) ROH, Oxone, rt, 19-22 h, R = Me 49%, R = *i*-Pr 66%; (vi) 2-methyl-2-butene, NaClO₂, NaHPO₄·H₂O, *t*-BuOH, H₂O, rt, 90 min, 50%.

Our alternative strategy involved concomitant protection of the double bond and the hydroxyl moiety in **5** as a cyclic bromoether. To this end, alcohol **5** was reacted with NBS in *t*-BuOH in a darkened reaction flask to afford **14** as a single diastereomer in good yield (Scheme 4). The regiochemistry of this reaction was easily deduced from the HMQC NMR spectra, which showed that the ¹³C signal assigned to the ether carbon (δ 83.3 ppm) had no hydrogen atoms attached to it. Notably, the relative configurations inherent to this transformation product are seen to be entirely controlled by the stereochemistries of the starting material. The remaining double bond was dihydroxylated and the resulting diol was cleaved to afford the corresponding keto aldehyde. Oxidation of this aldehyde to the carboxylic acid, followed by *O*-methylation with diazomethane, afforded ester **15**. Subsequent deprotection of **15** with zinc-copper couple gave rise to **16**.



Scheme 4. *Reagents and reaction conditions:* (i) NBS, *t*-BuOH, dark, rt, 4 h, 82%; (ii) OsO₄, NMO, THF, H₂O, 0 °C to rt, 5½ h, 58%; (iii) Pb(OAc)₄, 0 °C to rt, 2 h, 62%; (iv) 2-methyl-2-butene, NaClO₂, NaHPO₄·H₂O, *t*-BuOH, H₂O, rt, 90 min; (v) CH₂N₂, Et₂O, rt, 2 h, 77% over two steps; (vi) Zn/Cu, TMSCl, PhH, dimethylacetamide, 0 °C, 4 h, 50%, (87% BRSM).

In conclusion, the implementation of a dual protecting group strategy based on the regio- and diastereoselective intramolecular formation of a bromo ether has allowed for the preparation of the trans-fused perhydroazulene framework of **1**. We are currently investigating the completion of its total synthesis and we hope to report our results in due course.

ACKNOWLEDGEMENTS

JDP would like to acknowledge NSERC for a postdoctoral fellowship.

REFERENCES

1. For synthetic efforts towards **1** see: Z. Geng, B. Chen, and P. Chiu, *Angew. Chem. Int. Ed.*, 2006, **45**, 6197; B. Chen, R. Y. Y. Ko, M. S. M. Yuen, K. F. Cheng, and P. Chiu, *J. Org. Chem.*, 2003, **68**, 4195; P. Chiu, B. Chen, and K. F. Cheng, *Org. Lett.*, 2001, **3**, 1721; L. Ou, Y. Hu, G. Song, and D. Bai, *Tetrahedron*, 1999, **55**, 13999; Y. Hu, L. Ou, and D. Bai, *Tetrahedron Lett.*, 1999, **40**, 545; P. Chiu, B. Chen, and K. F. Cheng, *Tetrahedron Lett.*, 1998, **39**, 9229; B. C. Pan, H. Y. Chang, G. L. Cai, and Y. S. Guo, *Pure Appl. Chem.*, 1989, **61**, 389.
2. (a) E. Li, A. M. Clark, and C. D. Hufford, *J. Nat. Prod.*, 1995, **58**, 57; B. N. Zhou, B. P. Ying, G. Q. Song, Z. X. Chen, J. Han, and Y. F. Yan, *Planta Med.*, 1983, **47**, 35.
3. (a) V. K. W. Wong, P. Chiu, S. S. M. Chung, L. M. C. Chow, Y. Z. Zhao, B. B. Yang, and B. C. B. Ko, *Clin. Cancer Res.*, 2005, **11**, 6002; M. S. Jardat, D. J. Noonan, B. Wu, M. A. Avery, and D. R. Feller, *Planta Med.*, 2002, **68**, 667; D. J. Pan, Z. L. Li, C. Q. Hu, K. Chen, J. J. Chang, and K. H. Lee, *Planta Med.*, 1990, **56**, 383; M. O. Hamburger, H. L. Shieh, B. N. Zhou, J. M. Pezzuto, and G. A. Cordell, *Magn. Reson. Chem.*, 1989, **27**, 1025.
4. L. A. Paquette and J. D. Pettigrew, *Synthesis*, 2009, 379.
5. W. F. Berkowitz and Y. Wu, *J. Org. Chem.*, 1997, **62**, 1536.
6. M. R. Pitts, J. R. Harrison, and C. J. Moody, *J. Chem. Soc., Perkin Trans. 1*, 2001, 955.
7. P. A. Roethle, I. T. Chen, and D. Trauner, *J. Am. Chem. Soc.*, 2007, **129**, 8960; D. J. Wallace, *Angew. Chem. Int. Ed.*, 2005, **44**, 1912; X. Wang, E. J. Bowman, B. J. Bowman, and J. A. Porco, Jr., *Angew. Chem. Int. Ed.*, 2004, **43**, 3601; T. R. Hoye, C. S. Jeffrey, M. A. Tennakoon, J. Wang, and H. Zhao, *J. Am. Chem. Soc.*, 2004, **126**, 10210.
8. B. M. Trost, J. Waser, and A. Meyer, *J. Am. Chem. Soc.*, 2008, **130**, 16424; B. M. Trost, J. Waser, and A. Meyer, *J. Am. Chem. Soc.*, 2007, **129**, 14556.
9. B. R. Travis, M. Sivakumar, G. O. Hollist, and B. Borhan, *Org. Lett.*, 2003, **5**, 1031.