

HETEROCYCLES, Vol. 80, No. 2, 2010, pp. 725 - 738. © The Japan Institute of Heterocyclic Chemistry
Received, 20th August, 2009, Accepted, 26th October, 2009, Published online, 28th October, 2009
DOI: 10.3987/REV-09-SR(S)6

RECENT DEVELOPMENT FOR FORMATION OF AROMATIC COMPOUNDS VIA METALLACYCLOPENTADIENES AS METAL-CONTAINING HETEROCYCLES

Lishan Zhou, Shi Li, Ken-ichiro Kanno, and Tamotsu Takahashi*

Catalysis Research Center, Hokkaido University, Kita-ku, Sapporo 001-0021, Japan

Abstract – Metallacyclopentadiene, which is recognized as a metal-containing heterocycle, is an important intermediate for the formation of aromatic derivatives. In this short review, we summarized the recent advance in selective aromatic compounds formation catalyzed or mediated by transition metal complexes via metallacyclopentadienes which are formed by intermolecular coupling of various alkynes including unsymmetrical alkynes, two different alkynes and three different alkynes. Some reaction mechanisms were also reviewed.


1. INTRODUCTION

Reactions of two alkynes with low valent transition metals generally provide metallacyclopentadienes **1** as intermediates for the formation of benzene derivatives. The metallacyclopentadiene **1** is a cyclic compound with a metal. Therefore, it can be recognized as a metal-containing heterocycle (Scheme 1). Stepwise reaction of the metallacyclopentadienes with the third alkyne affords benzene derivatives **2** and **3**.

Many reviews on benzene derivatives formation from alkynes using transition metals have been published.¹ This review focuses on the recent development of selective formation of benzene derivatives from alkynes using transition metals as stoichiometric reactions or catalytic reactions.

Generally speaking, these type reactions include two different modes: intramolecular coupling reaction and intermolecular coupling reaction (Scheme 1). The intramolecular cyclotrimerization of triynes to prepare benzene derivatives **2** via metallacyclopentadienes **1** have been established as powerful synthetic methods. However, intermolecular and partially intermolecular versions generally have encountered difficulties in the control of chemo- and regioselectivity. When unsymmetrical alkynes or different


alkynes were employed for the cyclotrimerization, in most cases, isomeric aromatic derivatives **3** were generally formed.² This severely limited the utility of this method. Therefore, the search for a highly chemo- and regioselective intermolecular [2 + 2 + 2] cycloaddition reaction of alkynes is greatly challenging. Recently, several groups have devoted substantial effort in this area. Various transition metal complexes have been used to overcome this problem. This review introduces some advance in intermolecular cyclotrimerization of unsymmetrical alkynes or different alkynes catalyzed or mediated by transition metal complexes.



Scheme 1

2. CATALYTIC HOMO-CYCLOTRIMERIZATION OF UNSYMMETRICAL ALKYNES



There are two possible isomeric products of trisubstituted benzenes in a catalytic homo-cyclotrimerization of an unsymmetrical alkyne. As shown in Scheme 2, the reaction of a terminal alkyne can afford the corresponding 1,3,5- and 1,2,4-trisubstituted benzenes **4** and **5**, respectively.



Scheme 2

According to generally accepted metallacyclopentadiene mechanism shown in Scheme 3, 1,2,4-isomer **5** is formed exclusively via symmetrical metallacyclopentadienes **6** or **8**, and to some degree via unsymmetrical **7**. On the other hand, the possibility for the formation of 1,3,5-isomer **4** is smaller than that for 1,2,4-isomer **5**, since **4** can be formed via metalacycle **7** only in competition with formation of **5**. This consideration would conclude that the key points to achieve high selectivity for the formation of one

of the isomers are (1) selective formation of the intermediary metallacyclopentadiene, and (2) in the case of metalacycle **7**, regioselective addition of the third alkyne. Especially, for selective formation of 1,3,5-isomer **4**, the second key point is very crucial.









Tables 1 and 2 summarized the results of catalytic homo-cyclotrimerization of phenylacetylene to afford the 1,3,5- or 1,2,4-triphenylbenzenes **9** or **10** as a major isomer, respectively (Scheme 4). A comparison of the degree and type of selectivity depending on the employed catalysts is discussed here.

When camphorimine palladium complex **11** was employed for cyclotrimerization of phenylacetylene, 1,3,5-triphenylbenzene **9** was obtained as an exclusive benzene product (Table 1, entry 1).³ Although the product yield was low, the unique regioselectivity is worth noting. The same catalyst showed much higher efficiency for unsymmetrical internal alkyne without loss of regioselectivity. The cyclotrimerization of 1-phenylpropyne gave 1,3,5-trimethyl-2,4,6-triphenylbenzene in 74% yield as a sole product.

The predominant formation of the 1,3,5-isomer **9** was also observed in the reactions with ruthenium dinuclear complex **12** (entry 2),⁴ half-sandwiched cobalt **13** (entry 3),⁵ cycloheptatrienyliron **14** (entry 4),⁶ and dinuclear niobium **15** (entry 5).⁷ Among them, niobium catalyst **15** showed excellent selectivity and excellent product yield compared with the others.

Table 1. Homocyclotrimerization of Phenylacetylene with Various Catalysts to Afford 1,3,5-Triphenylbenzene **9** as a Major Isomer





entry	catalyst	conditions	1,3,5-/%	1,2,4-/%	ref.
1		3 mol % CH ₂ Cl ₂ , rt, 20 h	20	-	3
2		2.5 mol % H ₂ O: MeOH = 9:1 75 °C, 19 h	57	34	4
3		5 mol % H ₂ O: EtOH = 4:1 25 °C, 8 h	56	18	5
4		0.7 mol % THF, rt, 46 h	49	26	6
5	 L = THT or Me ₂ S	17 mol % CH ₂ Cl ₂ , rt, 4 h	98	0	7




Scheme 5

According to the metalacycle mechanism mentioned above, 2,4-disubstituted metallacyclopentadiene **7** must be formed as a major intermediate in these cases. In the niobium-catalyzed reaction, a possible mechanism was proposed as shown in Scheme 5.⁷ Dinuclear complex **15** reacts with two molecules of phenylacetylene to afford chlorine-bridged dimer of metallacyclopentadiene **16**, which was observed spectroscopically. Further insertion of the second and third alkynes can afford the corresponding dimer of metallacyclopentadiene **17**, and that of metallacycloheptatriene **18** followed by reductive elimination to produce the benzene **9** along with regeneration of the catalyst **15**. It is worth noting that both of the alkyne insertion steps proceed with high regioselectivity. Since the related mononuclear niobium complex did not show any catalytic activity, such superior efficiency can be ascribed to the dinuclear structure.

Table 2. Homocyclotrimerization of Phenylacetylene with Various Catalysts to Afford 1, 2, 4-Triphenylbenzene as a Major Isomer

entry	catalyst	conditions	1,3,5-/%	1,2,4-/%	ref.
1	RS-Au-Pd	19 4 mol %, CDCl ₃ , 62 °C, 3 h	19	80	8
2		20 2 mol %, toluene, reflux, 1 h	-	96	9
3	CoBr ₂ , 2,2'-bpy, Zn, ZnI ₂	21 5 mol %, MeCN, rt, <1 h	3	96	10
4	 CoCl ₂ ·6H ₂ O, Zn	22 5 mol %, THF, rt, 1-24 h	-	91	11
5		23 5 mol %, CH ₂ Cl ₂ , rt	3	91	12
6	RhCl ₃ ·3H ₂ O, <i>i</i> -Pr ₂ NEt	24 8 mol %, toluene, reflux, 24 h	6	92	13
7		25 0.2 mol %, C ₆ D ₆ , 75 °C	<3	>92	14

Palladium catalyst supported on gold nanoparticle **19** gave the 1,2,4-isomer **10** as a major product (Table 2, entry 1).⁸ Using a cobalt carbonyl cluster $\text{Co}_3(\text{CO})_9(\mu^3\text{-CH})$ **20** as catalyst, 1,2,4-isomeric benzene **10** was obtained with excellent regioselectivity (entry 2).⁹ The trinuclear complex is stable and easily prepared from dicobalt octacarbonyl compound and trihalomethane. The simple cobalt catalyst system **21** was also found for the efficient and regioselective cyclotrimerization of alkynes (entry 3).¹⁰ In the case of phenylacetylene, 1,2,4-triphenylbenzene was obtained in 99% yield in 95:5 regioselectivity. The cobalt catalyst system **21** is applicable for cyclotrimerization of unsymmetrical internal alkynes to furnish the corresponding 1,2,4-substituted benzenes with good regioselectivity in high yields. In the cobalt catalyst system, when sterically demanding iminomethylpyridine was used as a ligand instead of bipyridine, the reaction system **22** became less sensitive to moisture without loss of efficiency (entry 4).¹¹ Rhodium catalysts also showed outstanding regioselectivity for the 1,2,4-isomer **10**. Catalytic activity of cationic rhodium(I) segphos complex **23** is exceedingly high and the reaction can be carried out under mild conditions (entry 5).¹² Although the reaction conditions are more drastic, just simple combination of rhodium(III) chloride and amine **24** produced efficient catalytic system (entry 6).¹³ Except for the late transition metal catalysts mentioned above, a titanium complex **25** having 1,2-alternate bridged *p*-*tert*-butylcalix[4]arene can also act as a highly regioselective catalyst (entry 7).¹⁴ The complex is pre-activated with sodium or magnesium metal. The reaction of activated **25** in the presence of an excess of trimethylsilylacetylene produced the corresponding titanium complex with η^6 -coordinated 1,2,4-tris(trimethylsilyl)benzene, which is also an efficient catalyst for cyclotrimerization of alkynes at room temperature.




Scheme 6

Cross-cyclotrimerization of trimethylsilylacetylene with 4-chlorostyrene with titanium complex **25** gave the corresponding cyclohexadiene **27** as a major product (scheme 6). The result suggests the formation of α,β' -disubstituted metalacycle **26** predominantly. This means that although the α,α' - or β,β' -disubstituted metalacycles can afford the corresponding 1,2,4-trisubstituted benzene exclusively in the homo-cyclotrimerization, their contribution is minor. Moreover, insertion of the third alkyne into


metalacycle **26** should occur in highly regioselective manner. Probably, the major reason of such high regioselectivity can be ascribed to sterically demanding nature of the calix[4]arene ligand.

3. CATALYTIC CROSS-CYCLOTRIMERIZATION OF TWO DIFFERENT ALKYNES

In trimerization reactions using two different alkynes, late transition metals, especially group 8-10 metals, were mainly used as catalysts. Mitsudo *et al.* developed a ruthenium-catalyzed synthesis of *o*-phthalates by intermolecular cycloaddition of two different alkynes.¹⁵ Excess amount of terminal alkynes reacted with DMAD in the presence of Cp^{*}RuCl(cod) catalyst to give an isomeric mixture of dialkylated *o*-phthalates in high yield. As shown in Scheme 7, ruthenacycle **28** which consists of two different alkynes was proposed as a plausible intermediate.




An efficient cobalt-catalyzed cyclotrimerization for synthesis of benzolactones **29** was reported (Scheme 8).¹⁶ In this reaction, alkynyl alcohols reacted with 2 equiv of acetylenic esters to form substituted benzolactones **29** in a highly chemo and regioselective fashion with excellent yields. Interestingly, when the catalyst was changed, the regioselectivity was reversed at the metalacycle formation step (Scheme 9). In the case of the cobalt catalyst, the corresponding metallacyclopentadiene **30** having two ester groups at α, α' -positions, whereas the β, β' -isomer **31** formed in that of the nickel catalyst and no transesterification occurred to afford benzyl alcohol **32** in a good yield.




Adoption of iridium-phosphine complex in alkyne cyclotrimerization was also reported.¹⁷ As shown in Scheme 10, chemoselectivity of the iridium catalyst can be controlled by changing the employed phosphine ligands. When dppe ligand was used for the catalysis, 2:1 coupling products benzene tetraester **34** was obtained in excellent yield and selectivity. On the other hand, when polyfluorinated diphosphine, 1,2-bis[bis(pentafluorophenyl)phosphino]ethane is used, 1:2 coupling product *o*-phthalate **36** was obtained with high selectivity. These results suggest the complete difference of the

intermediary metalacycles. In the case of dppe, benzene tetraester **34** was formed via iridacycle **33**, whereas *o*-phthalate **36** was obtained in the reaction with polyfluorinated dppe via iridacycle **35** as a predominant intermediate. The electron-deficient nature of the fluorinated ligand would make coordination of electron-rich alkyne **B** toward the iridium center much more favored than that of electron-deficient alkyne **A**.



Scheme 9




Scheme 10


Highly chemo- and regioselective intermolecular cyclotrimerization of two different alkyne moieties catalyzed by rhodium (I) complex was also reported (Scheme 11).^{12,18} In the presence of a cationic rhodium(I)/H₈-BINAP complex, excellent regioselectivity was observed. Two molecules of a wide variety of terminal alkynes cleanly reacted with one molecule of a dialkyl acetylenedicarboxylate to give 3,6-disubstituted dialkyl phthalates **37** in high yields and with high regioselectivity.

Based on the numerous experimental results, the origin of the high selectivity was explained as shown in Scheme 12. First, two different alkynes react with cationic rhodium(I) in 1:1 ratio to afford the

corresponding metallacyclopentadiene **38**. The origin of this selectivity is due to the steric reason. Coordination of another terminal alkyne to **38** affords complex **39**. In this step, the electronic nature of the α -carbon of the metalacycle and the coordinated alkyne determines the regioselectivity to afford 1,2,3,4-tetrasubstituted benzene.




Scheme 11



Scheme 12

4. CROSS-CYCLOTTRIMERIZATION OF THREE DIFFERENT ALKYNES

One of the major problems of selective benzene formation from three different alkynes is the control of combination of alkynes. It has been difficult to prepare cleanly the unsymmetrical metallacyclopentadienes using transition metals by the intermolecular coupling of two different alkynes. To solve this problem,¹⁹ Takahashi and co-workers studied the formation of unsymmetrical zirconacyclopentadienes. Symmetrical zirconacyclopentadienes were conveniently prepared from alkynes and Cp_2ZrBu_2 (Negishi reagent).²⁰ They found the use of ethylene gas was a prerequisite to prepare very clean unsymmetrical zirconacyclopentadienes.²¹ The first alkyne was treated with Negishi reagent under ethylene gas to give zirconacyclopentadienes **40**. Subsequent addition of a second alkyne to the species at 50 °C gave unsymmetrical zirconacyclopentadienes **41** selectively in high yields. Homo-coupling products of either the first alkyne or the second alkyne were negligible (Scheme 13).




Scheme 13

The use of unsymmetrical zirconacyclopentadienes resulted in the formation of unsymmetrical benzene derivatives **42**.²² Thus prepared benzene rings have alkyl, aryl, trimethylsilyl and ester groups. This was the first and practical example of the selective coupling of three different alkynes in high yields in one-pot.


They found zirconacyclopentadienes **41** reacted with alkynes with electron-donating groups as well as alkynes with electron-withdrawing groups to give benzene derivatives **42** in the presence of $\text{NiX}_2(\text{PPh}_3)_2$ ($\text{X} = \text{Cl}$ or Br) in good yields.²³ Using Zr/Cu and Zr/Ni-mediated synthetic methods, not only aromatic carbocycles but also aromatic heterocycles could be prepared.²⁴

Sato and co-workers reported a metalative Reppe type of acetylene cyclotrimerization.²⁵ In this reaction, three different unsymmetrical acetylenes and one molecule of titanium species were combined together in a highly controlled manner to give aryltitanium compounds as a single isomer. As shown in Scheme 14, a dialkoxytitanacyclopentadienes **43** were prepared from two different acetylenes and a divalent titanium alkoxide reagent $(\eta^2\text{-propene})\text{Ti}(\text{O-}i\text{-Pr})_2$. Addition of ethynyl para-tolyl sulfone as the third alkyne afforded aryl titanium intermediates **44**, which reacted with various electrophiles to produce a series of highly substituted benzenes **45**.



Scheme 14


Compared with the stoichiometric reactions mentioned above, the catalytic version of the selective three-component cycloaddition is highly crucial task. $\text{Cp}^*\text{RuCl}(\text{cod})$ -catalyzed system was applied to chemoselective $[2+2+2]$ cycloaddition of three different alkynes.²⁶ An internal alkyne, a terminal alkyne and dimethyl acetylenedicarboxylate was efficiently catalyzed by $\text{Cp}^*\text{RuCl}(\text{cod})$ to give substituted *o*-phthalates **46** and **48** (Scheme 15). It is critical to control the molar ratio of the three substrates to achieve high chemoselectivity.



Scheme 15


Using semi-intramolecular strategy, selective catalytic cyclotrimerization of three unsymmetrical alkynes

was realized (Scheme 16).²⁷ Alkynylboronates reacted with propargyl alcohol to form unsymmetrical diynes with a temporal C-B-O linkage. In the presence of Ru catalyst, cycloaddition of diynes with the third terminal alkynes afford arylboronates **50** via ruthenacycles **49** with excellent chemo- and regioselectivity. The crude arylboronates **50** underwent Suzuki–Miyaura cross-coupling with various aryl iodides in the presence of Pd-catalyst to afford a series of substituted biaryls **51** as single regioisomers. They also developed the Pd (II)-catalyzed carbonylation of the boraphthalides to establish the novel four-component coupling synthesis of substituted phthalides in good yields.²⁸



Scheme 16

The similar strategy was also reported with a cobalt complex.²⁹ Use of temporary silylated tethers **52** as starting materials allowed the transformation of intermolecular reaction to intramolecular ones. Silicon-tethered triynes **52** led to the corresponding cycloadducts **53** in good yields in the presence of 5 mol% of $[\text{CpCo}(\text{CO})_2]$. The opening of the silylated ether was easily achieved either with *n*-tetrabutylammonium fluoride or with methyllithium to deliver quantitatively the benzene derivatives **54** after the reaction (Scheme 17).



Scheme 17

5. SUMMARY

A remarkable effect of transition metals on the trimerization of alkynes was discovered in 1940's by Reppe. After that, this strategy has been developed enormously as the most powerful tool for construction benzene and the other aromatic ring skeletons. At the beginning of the history, two possible mechanisms were proposed for the $[2+2+2]$ cyclotrimerization. One is a cyclobutadiene

mechanism in which a cyclobutadiene-transition metal complex derived from two alkynes by [2+2] cycloaddition is an intermediate. The other is a metallacyclopentadiene mechanism, which is now commonly accepted from a large number of experimental results.

Recent developments of the aromatic ring construction methods by the [2+2+2] cycloaddition can be regarded as the consequence of the intermediacy of the metallacyclopentadiene. That provides an important opportunity for regioselective construction of substituted benzene rings from substituted alkynes. In fact, the controlled formation of the metallacyclopentadiene intermediate is one of the most important key points for selective construction of aromatic rings as we described above. Further development of this chemistry has still been desired because of growing importance of the aromatic compound in various areas in human society.

6. REFERENCES

1. K. P. C. Vollhardt, *Acc. Chem. Res.*, 1977, **10**, 1; K. P. C. Vollhardt, *Angew. Chem., Int. Ed. Engl.*, 1984, **23**, 539; N. E. Schore, *Chem. Rev.*, 1988, **88**, 1081; I. Ojima, M. Tzamarioudaki, Z. Li, and R. J. Donovan, *Chem. Rev.*, 1996, **96**, 635; H.-W. Fruhauf, *Chem. Rev.*, 1997, **97**, 523; S. Saito and Y. Yamamoto, *Chem. Rev.*, 2000, **100**, 2901; S. Kotha, E. Brahmachary, and K. Lahiri, *Eur. J. Org. Chem.*, 2005, 4741; V. Gandon, C. Aubert, and M. Malacria, *Chem. Commun.*, 2006, 2209; P. R. Chopadea and J. Louie, *Adv. Synth. Catal.*, 2006, **348**, 2307; B. R. Galan and T. Rovis, *Angew. Chem. Int. Ed.* 2009, **11**, 2830; N. Agenet, O. Buisine, F. Slowinski, V. Gandon, C. Auvert, and M. Malacria, *Org. React.*, 2007, **68**, 1; Y. Yamamoto, *Cur. Org. Chem.*, 2005, **9**, 503.
2. G. P. Colman, G. W. Kang, W. F. Little, and M. F. Sullivan, *Inorg. Chem.*, 1968, **7**, 1298; R. A. Ferreri and A. P. Wolf, *J. Phys. Chem.*, 1984, **88**, 2256; M. Lautens, W. Klute, and W. Tam, *Chem. Rev.*, 1996, **96**, 49.
3. M. F. N. N. Carvalho, F. M. T. Almeida, A. M. Galvão, and A. J. L. Pombeiro, *J. Organomet. Chem.*, 2003, **679**, 143.
4. V. Cadierno, S. E. García-Garrido, and J. Gimeno, *J. Am. Chem. Soc.*, 2006, **128**, 15094.
5. L. Yong and H. Butenschön, *Chem. Commun.*, 2002, 2852.
6. C. Breschi, L. Piparo, P. Pertici, A. M. Caporusso, and G. Vitulli, *J. Organomet. Chem.*, 2000, **607**, 57.
7. M. Kakeya, T. Fujihara, T. Kasaya, and A. Nagasawa, *Organometallics*, 2006, **25**, 4131.
8. Y.-Y. Lin, S.-C. Tsai, and S. J. Yu, *J. Org. Chem.*, 2008, **73**, 4920.
9. T. Sugihara, A. Wakabayashi, Y. Nagai, H. Takao, H. Imagawa, and M. Nishizawa, *Chem. Commun.*, 2002, 576.
10. G. Hilt, T. Vogler, W. Hess, and F. Galbiati, *Chem. Commun.*, 2005, 1474; G. Hilt, W. Hess, T.

- Vogler, and C. Hengst, *J. Organomet. Chem.*, 2005, **690**, 5170.
11. N. Saino, F. Amemiya, E. Tanabe, K. Kase, and S. Okamoto, *Org. Lett.*, 2006, **8**, 1439; A. Goswami, T. Ito, and S. Okamoto, *Adv. Synth. Catal.*, 2007, **349**, 2368.
 12. K. Tanaka, K. Toyoda, A. Wada, K. Shirasaka, and M. Hirano, *Chem. Eur. J.*, 2005, **11**, 1145.
 13. K. Yoshida, I. Morimoto, K. Mitsudo, and H. Tanaka, *Chem. Lett.*, 2007, **36**, 998.
 14. O. V. Ozerov, F. T. Ladipo, and B. O. Patrick, *J. Am. Chem. Soc.*, 1999, **121**, 7941; O. V. Ozerov, C. P. Brock, S. D. Carr, and F. T. Ladipo, *Organometallics*, 2000, **19**, 5016; O. V. Ozerov, B. O. Patrick, and F. T. Ladipo, *J. Am. Chem. Soc.*, 2000, **122**, 6423; F. T. Ladipo, V. Sarveswaran, J. V. Kingston, R. A. Huyck, S. Y. Bylikin, S. D. Carr, R. Watts, and S. Parkin, *J. Organomet. Chem.*, 2004, **689**, 502.
 15. Y. Ura, Y. Sato, M. Shiotsuki, T. Kondo, and T. Mitsudo, *J. Mol. Catal. A: Chem.*, 2004, **209**, 35.
 16. H.-T. Chang, M. Jeganmohan, and C.-H. Cheng, *Chem. Commun.*, 2005, 4955.
 17. R. Takeuchi and Y. Nakaya, *Org. Lett.*, 2003, **5**, 3659.
 18. K. Tanaka and K. Shirasaka, *Org. Lett.*, 2003, **5**, 4697.
 19. A pioneering work in this subject, see: Y. Wakatsuki, T. Kuramitsu, and H. Yamazaki, *Tetrahedron Lett.* 1974, **15**, 4549.
 20. E. Negishi, D. R. Swanson, F. E. Cederbaum, and T. Takahashi, *Tetrahedron Lett.*, 1986, **27**, 2829; E. Negishi, S. J. Holmes, J. Tour, J. A. Miller, F. E. Cederbaum, D. R. Swanson, and T. Takahashi, *J. Am. Chem. Soc.*, 1989, **111**, 3336.
 21. Z. Xi, R. Hara, and T. Takahashi, *J. Org. Chem.*, 1995, **60**, 4444; T. Takahashi, M. Kageyama, V. Denisov, R. Hara, and E. Negishi, *Tetrahedron Lett.*, 1993, **34**, 687; T. Takahashi, Z. Xi, C. J. Rousset, and N. Suzuki, *Chem. Lett.*, 1993, 1001.
 22. T. Takahashi, M. Kotoru, and Z. Xi, *J. Chem. Soc., Chem. Commun.*, 1995, 361; T. Takahashi, Z. Xi, A. Yamazaki, Y. Liu, K. Nakajima, and M. Kotoru, *J. Am. Chem. Soc.*, 1998, **120**, 1672; T. Takahashi, F.-Y. Tsai, and Y. Li, *Chem. Lett.*, 1999, 1173; Z. Xi and T. Takahashi, *Huaxue Xuebao*, 2000, **58**, 1177; T. Takahashi, *Pure Appl. Chem.*, 2001, **73**, 271.
 23. T. Takahashi, F.-Y. Tsai, Y. Li, K. Nakajima, and M. Kotoru, *J. Am. Chem. Soc.*, 1999, **121**, 11093; H. Wang, F.-Y. Tsai, and T. Takahashi, *Chem. Lett.*, 2000, 1410.
 24. L. Zhou, M. Yamanaka, K. Kanno, and T. Takahashi, *Heterocycles*, 2008, **76**, 923. And references cited therein.
 25. D. Suzuki, H. Urabe, and F. Sato, *J. Am. Chem. Soc.*, 2001, **123**, 7925; R. Tanaka, Y. Nakano, D. Suzuki, H. Urabe, and F. Sato, *J. Am. Chem. Soc.*, 2002, **124**, 9682.
 26. Y. Ura, Y. Sato, H. Tsujita, T. Kondo, M. Imachi, and T. Mitsudo, *J. Mol. Catal. A: Chem.*, 2005, **239**, 166.

27. Y. Yamamoto, J. Ishii, H. Nishiyama, and K. Itoh, *J. Am. Chem. Soc.*, 2004, **126**, 3712; Y. Yamamoto, J. Ishii, H. Nishiyama, and K. Itoh, *Tetrahedron*, 2005, **61**, 11501.
28. Y. Yamamoto, J. Ishii, H. Nishiyama, and K. Itoh, *J. Am. Chem. Soc.*, 2005, **127**, 9625.
29. G. Chouraqui, M. Petit, C. Aubert, and M. Malacria, *Org. Lett.*, 2004, **6**, 1519.
-



Lishan Zhou received his bachelor's degree from the Hebei University of Technology in 1996 and Ph. D degree from Tianjin University in 2003 (China). After he finished the postdoctorate work at Tsinghua Univ. (Prof. Xi Chanjuan) in 2005 he moved to Professor Takahashi's group in Hokkaido University as a postdoctoral fellow and now a special assistant professor (Sapporo, Japan). His current research focuses on the development of new synthetic methods mediated by zirconocene and development of polyacene derivatives as organic materials.



Shi Li received his bachelor's and master's degree from Zhengzhou University (China) in 2000 and 2003, respectively. Then he joined Professor Takahashi's group at Hokkaido University (Sapporo, Japan) as a Ph. D. student from 2004. After he received his Ph. D. degree in 2007, he continued his research work in the same group as a postdoctoral fellow. His current research interest is focused on zirconium-mediated acene synthesis and their application as organic semiconductors.



Ken-ichiro Kanno received his B. S. (1994), M. S. (1998), and Ph. D. (2001) from Tohoku University (Sendai, Japan) under the supervision of Prof. Mitsuo Kira. In 2001, he joined to Prof. Takahashi's group at Hokkaido University (Sapporo, Japan) as postdoctoral research fellow, and in 2003 he became assistant professor at the same group. His current research focused on development of organic synthesis with organometallic reagents and application of extended π -conjugated molecules toward organic semiconductors.



Tamotsu Takahashi received his bachelor's and Ph. D. degree from the University of Tokyo in 1978 and 1983, respectively. He became an assistant professor at the same university in 1983. From 1984 to 1986, he joined Prof. Negishi's group at Purdue University as a postdoctoral research fellow. In 1991 he was promoted to an associate professor at Institute for Molecular Science. In 1995, he moved to Hokkaido University as a full professor. He received the Progress Award in Synthetic Organic Chemistry, Japan in 1993 and Divisional Award (Organic Chemistry) of Chemical Society of Japan in 1998. He was Director of Catalysis Research Center, Hokkaido University (2002-2006), a member of "Science Council of Japan" (2006-2008) and program officer of Research Center for Science System, Japan Society for the Promotion and Science (JSPS) (2006-2009). He was also a leader of CREST (1998-2003) and SORST (2003-2008) projects of Japan Science & Technology Agency (JST). His research covers several areas that include highly selective organic synthesis, development of novel reactions using organometallic compounds in particular, zirconium and titanium compounds, carbon-carbon bond cleavage reaction, organic semiconductor materials such as substituted soluble acene derivatives, and organic thin-film solar cells.