

HETEROCYCLES, Vol. 82, No. 2, 2011, pp. 1657 - 1662. © The Japan Institute of Heterocyclic Chemistry  
Received, 16th June, 2010, Accepted, 16th August, 2010, Published online, 17th August, 2010  
DOI: 10.3987/COM-10-S(E)59

## THE PREPARATION OF KETENE DITHIOACETALS AND THIOPHENES FROM CHLOROPYRIDINES CONTAINING AN ACTIVE METHYLENE GROUP

Keith R. Sturrock,<sup>1\*</sup> David H. Bremner,<sup>1</sup> and Grant Wishart<sup>2</sup>

<sup>1</sup>University of Abertay Dundee, School of Contemporary Sciences, Bell Street, Dundee, DD1 1HG, Scotland, U.K. k.sturrock@abertay.ac.uk

<sup>2</sup>Schering Plough Res Inst, Dept Chem, Newhouse ML1 5SH, Lanark, Scotland, U.K.


**Abstract** – The base catalysed reaction of carbon disulphide with the active methylene groups of chloropyridines **4** and **7**, followed by alkylation with reagents which also contain active methylene groups, lead to ketene dithioacetals. Further reaction with base afforded highly substituted thiophenes.

### INTRODUCTION

Our work<sup>1-3</sup> on the synthesis of thienopyridines required us to prepare a series of  $\alpha$ -halogenated pyridines containing a methylene moiety activated by an electron withdrawing group. We earlier reported<sup>1,2</sup> that reaction of pyridine 3-acetonitrile N-oxide **1** with refluxing POCl<sub>3</sub>, using the method reported by Okuda and Robison<sup>4</sup> gave the desired chloropyridines **3** and **5** with **4** as a byproduct. Similarly<sup>2</sup> we obtained **6**, **7** and **8** from ethyl 3-pyridylacetate N-oxide **2** (**Equation 1**). The ratio<sup>1,2</sup> of isomers differs for each substituent at the 3-position (**Table 1**). It is well known<sup>5-12</sup> that polyfunctional ketene dithioacetals are available from reaction of carbon disulphide with functionalized carbionic species followed by alkylation of the heteroanions including cyano(3-pyridyl)ketene dimethyldithioacetal<sup>11</sup> from 3-cyanomethylpyridine. Furthermore, it has also been reported<sup>5-10</sup> that treatment of appropriately functionalized ketene dithioacetals with base allows entry into a range of highly substituted heterocycles. In this work, haloalkanes with electron withdrawing groups on the same carbon as the halogen were employed as alkylating agents in the base catalysed nucleophilic addition reaction of **4** and **7** with carbon disulphide. This gave ketene dithioacetals with the required functionality for cyclisation to form thiophenes.

**Table 1.** Product Ratio by Substituent at C-3

Substitution pattern	R = CH <sub>2</sub> CN (%)	R = CH <sub>2</sub> CO <sub>2</sub> Et (%)
2,3- ( <b>3</b> , <b>6</b> )	37	33
2,5- ( <b>4</b> , <b>7</b> )	21	22
3,4- ( <b>5</b> , <b>8</b> )	5	13

**Equation 1**

**1**, **3**, **4**, **5**; R = CH<sub>2</sub>CN and **2**, **6**, **7**, **8**; R = CH<sub>2</sub>CO<sub>2</sub>Et


## RESULTS AND DISCUSSION

When 2-chloro-5-cyanomethylpyridine **4** and carbon disulphide were treated with a slight excess of sodium hydride in DMSO under anhydrous conditions followed by addition of ethyl chloroacetate, ketene dithioacetal **9** was synthesised in 58% yield. The infrared spectrum showed the molecule to contain a nitrile group (2209 cm<sup>-1</sup>) and an aliphatic ester (1736 cm<sup>-1</sup>). The <sup>1</sup>H NMR corroborated the structure, showing signals for two ethyl thioacetate groups and a 2,5 disubstituted pyridine. Further treatment of **9** with sodium ethoxide in ethanol gave thiophene **10** (55%). The infrared spectrum for this compound showed a primary amine (3447 and 3343 cm<sup>-1</sup>) and two ester groups (1724, 1661 cm<sup>-1</sup>). The absorbance at 1724 cm<sup>-1</sup> can be interpreted as being due to the aliphatic ethyl ester. The second, at 1661 cm<sup>-1</sup> can be assigned to the ester directly attached to the thiophene ring, having a similar absorption wavenumber as an anthranilate ester. There will be some hydrogen bonding between the ester carbonyl and the amino hydrogens and this phenomenon is known<sup>13</sup> to move the absorbance to a lower wavenumber. It is thought that the mechanism for the formation of **9** proceeds by base catalysed abstraction of the methylene protons of **4** and subsequent formation, with carbon disulphide, of a ketene dithioacetal dianion. These anions are then alkylated with ethyl chloroacetate. The thiophene **10** is subsequently formed by a Thorpe<sup>14</sup> type cyclisation of the active methylene adjacent to the sulphur onto the nitrile. Ketene dithioacetal **11** was similarly prepared from **4** (54%) using chloroacetonitrile as the alkylating agent and **11** was readily cyclised to the corresponding thiophene **12** (51%) with sodium ethoxide again probably via a Thorpe type reaction.

This methodology was extended to ethyl 2-chloro-5-pyridylacetate **7**, to produce ketene dithioacetal **13**. The infrared spectrum showed a carbonyl stretching absorbance in the expected region for aliphatic esters ( $1736\text{ cm}^{-1}$ ). This was unexpected, as it also contains an  $\alpha,\beta$ -unsaturated ethyl ester group and a second peak should have been observed at a lower wavenumber, typically,<sup>13</sup>  $1730\text{-}1705\text{ cm}^{-1}$ . The  $^1\text{H}$  NMR however supported the structure, showing signals for two ethyl thioacetate groups and an ethyl ester group. Further reaction of **13** with sodium ethoxide in ethanol lead to the production of thiophene **14** in 33% yield. Again two carbonyl stretching absorbances were observed in the IR spectrum, an aliphatic ester ( $1748\text{ cm}^{-1}$ ), and a salicylate type ester at  $1666\text{ cm}^{-1}$ . It is presumed that formation is by a Dieckmann<sup>15</sup> cyclisation of one of the methylene groups onto the vinyl ester. Reaction of **7** with chloroacetonitrile under the same reaction conditions, gave **15** in 57% yield. The IR spectrum this time showed a carbonyl stretching absorbance typical<sup>13</sup> of an  $\alpha,\beta$ -unsaturated ester group ( $1720\text{ cm}^{-1}$ ). Further reaction of **15** with base afforded **16** in 46% yield, again by a Dieckmann cyclisation.

In conclusion, novel ketene dithioacetals can be conveniently prepared, leading to polyfunctional thiophenes (**Scheme 1**) and we believe this method can be applied in the synthesis of a variety similar highly substituted thiophenes.

Scheme 1



## EXPERIMENTAL

### 1.0. General

Melting points were determined using an Electrothermal melting point apparatus and are uncorrected. Infrared spectra were recorded on a Perkin-Elmer Paragon spectrophotometer.  $^1\text{H}$  NMR spectra were recorded on a Jeol PMX 60i spectrometer. Column chromatography was performed using pressurised short path columns with Kieselgel 60, particle size  $< 0.063\text{ mm}$  (Merck No. 7729) and reactions were

monitored with Merck DC-Alufoilien 60 F<sub>254</sub> (Merck No. 5554) which were visualised by ultraviolet irradiation.

### 1.1. Preparation of ketene dithioacetals.

#### 1.1.1. General procedure; 2-chloro-5-(1-cyano-2,2-di(ethyloxycarbonylmethylthio)-1-vinyl)pyridine (9) as an example.

2-Chloro-5-cyanomethylpyridine **4** (2.0 g, 13.1 mmol) and carbon disulphide (1.1 g, 14.4 mmol) were dissolved in DMSO (40 mL) in a 2-neck round bottom flask fitted with a gas inlet tube under dry nitrogen and sodium hydride (0.66 g, 27.5 mmol) was added in portions with constant stirring and the mixture was stirred for 2 h. Ethyl chloroacetate (3.38 g, 27.5 mmol) was added and stirring continued for another hour, then poured onto ice (100 g) and extracted with EtOAc (3 x 50 mL). The extracts were combined, dried with magnesium sulphate, filtered and the solvent evaporated to give a pale yellow oil (4.46 g) which was chromatographed on silica gel using petroleum ether/EtOAc to give **9** as a colourless oil (58%). Bp (decomp); Anal. Calcd for C<sub>16</sub>H<sub>17</sub>ClN<sub>2</sub>O<sub>4</sub>S<sub>2</sub>: C, 47.9; H, 4.3; N, 7.0%. Found: C, 48.0; H, 4.4; N, 7.1%; IR  $\nu_{\max}$  (film)/cm<sup>-1</sup> 2982, 2209, 1736; <sup>1</sup>H NMR  $\delta_{\text{H}}$  (CDCl<sub>3</sub>) 1.28 (3H, t, J=7.2 Hz, CH<sub>3</sub>), 1.32 (3H, t, J=7.2 Hz, CH<sub>3</sub>), 3.72 (2H, s, SCH<sub>2</sub>), 3.80 (2H, s, SCH<sub>2</sub>), 4.16 (2H, q, J=7.2 Hz, CO<sub>2</sub>CH<sub>2</sub>), 4.20 (2H, q, J=7.2 Hz, CO<sub>2</sub>CH<sub>2</sub>), 7.33 (1H, d, J=8.4 Hz, H-3), 7.85 (1H, dd, J=8.4 & 2.4 Hz, H-4), 8.52 (1H, d, J=2.4 Hz, H-6).

#### 1.1.2. 2-Chloro-5-(1-cyano-2,2-di(cyanomethylthio)-1-vinyl)pyridine (11)

Off white crystals (61%, EtOAc). Mp 101-102 °C; Anal. Calcd for C<sub>11</sub>H<sub>7</sub>ClN<sub>4</sub>S<sub>2</sub>: C, 47.0; H, 2.4; N, 18.3%. Found: C, 46.9; H, 2.3; N, 18.3%; IR  $\nu_{\max}$  (KBr)/cm<sup>-1</sup> 2978, 2234; <sup>1</sup>H NMR  $\delta_{\text{H}}$  (CDCl<sub>3</sub>) 4.48 (4H, s, SCH<sub>2</sub>), 7.38 (1H, d, J=8.4 Hz, H-3), 7.66 (1H, dd, J=8.4 & 2.4 Hz, H-4), 8.32 (1H, d, J=2.4 Hz, H-6).

#### 1.1.3. 2-Chloro-5-(1-ethyloxycarbonyl-2,2-di(ethyloxycarbonylmethylthio)-1-vinyl)pyridine (13)

Colourless oil (43%); bp (0.4) 190 °C; Anal. Calcd for C<sub>18</sub>H<sub>22</sub>ClNO<sub>6</sub>S<sub>2</sub>: C, 49.3; H, 4.9; N, 3.1%. Found: C, 49.6; H, 4.8; N, 3.3%; IR  $\nu_{\max}$  (film)/cm<sup>-1</sup> 2982, 2937, 1736; <sup>1</sup>H NMR  $\delta_{\text{H}}$  (CDCl<sub>3</sub>) 1.24 (3H, t, J=7.2 Hz, CH<sub>3</sub>), 1.28 (6H, t, J=7.2 Hz, CH<sub>3</sub>), 3.56 (2H, s, SCH<sub>2</sub>), 3.68 (2H, s, SCH<sub>2</sub>), 4.08 (4H, q, J=7.2 Hz, CO<sub>2</sub>CH<sub>2</sub>), 4.14 (2H, q, J=7.2 Hz, CO<sub>2</sub>CH<sub>2</sub>), 7.28 (1H, d, J=8.4 Hz, H-3), 7.72 (1H, dd, J=8.4 & 2.4 Hz, H-4), 8.34 (1H, d, J=2.4 Hz, H-6).

#### 1.1.4. 2-Chloro-5-(1-ethyloxycarbonyl-2,2-di(cyanomethylthio)-1-vinyl)pyridine (15)

Pale yellow needles (57%, EtOH). Mp 98-99 °C; Anal. Calcd for C<sub>14</sub>H<sub>12</sub>ClN<sub>3</sub>O<sub>2</sub>S<sub>2</sub>: C, 47.5; H, 3.4; N, 11.9%. Found: C, 47.4; H, 3.0; N, 11.7%; IR  $\nu_{\max}$  (KBr)/cm<sup>-1</sup> 2976, 2246, 1720; <sup>1</sup>H NMR  $\delta_{\text{H}}$  (CDCl<sub>3</sub>) 1.25

(3H, t, J=7.2 Hz, CH<sub>3</sub>), 3.56 (2H, s, SCH<sub>2</sub>), 3.68 (2H, s, SCH<sub>2</sub>), 4.24 (2H, q, J=7.2 Hz, CO<sub>2</sub>CH<sub>2</sub>), 7.24 (1H, d, J=8.4 Hz, H-3), 7.58 (1H, dd, J=8.4 & 2.4 Hz, H-4), 8.24 (1H, d, J=2.4 Hz, H-6).

## 1.2. Preparation of thiophenes

### 1.2.1. General procedure; **2-chloro-5-(2-ethyloxycarbonyl-5-ethyloxycarbonylmethylthio-3-aminothienyl)pyridine (10)** as an example.

Ketene dithioacetal **9** (0.5 g, 1.1 mmol) was dissolved in EtOH (20 mL) in a 50 mL round bottom flask and sodium ethoxide (0.16 g, 2.4 mmol) added. The mixture was stirred at room temperature for 24 h. The solvent was removed *in vacuo*, the residue flooded with water and extracted with EtOAc (3 x 25 mL). The combined extracts were dried with magnesium sulphate, filtered and the solvent removed to give thiophene **10** (55%) after recrystallisation from EtOAc. Mp 94-95 °C; Anal. Calcd for C<sub>16</sub>H<sub>17</sub>ClN<sub>2</sub>O<sub>4</sub>S<sub>2</sub>: C, 47.9; H, 4.3; N, 7.0%. Found: C, 48.1; H, 4.2; N, 6.9%; IR  $\nu_{\max}$ (KBr)/cm<sup>-1</sup> 3447, 3343, 2967, 1724, 1661; <sup>1</sup>H NMR  $\delta_{\text{H}}$ (CDCl<sub>3</sub>) 1.17 (3H, t, J=7.2 Hz, CH<sub>3</sub>), 1.26 (3H, t, J=7.2 Hz, CH<sub>3</sub>), 3.46 (2H, s, SCH<sub>2</sub>), 4.02 (2H, q, J=7.2 Hz, CO<sub>2</sub>CH<sub>2</sub>), 4.18 (2H, q, J=7.2 Hz, CO<sub>2</sub>CH<sub>2</sub>), 5.36 (2H, s, NH<sub>2</sub>), 7.30 (1H, d, J=8.4 Hz, H-3), 7.64 (1H, dd, J=8.4 & 2.4 Hz, H-4), 8.26 (1H, d, J=2.4 Hz, H-6).

### 1.2.2. **2-Chloro-5-(2-cyano-5-cyanomethylthio-3-aminothienyl)pyridine (12)**

Pale yellow crystals (51%, EtOAc). Mp 113-114 °C; Anal. Calcd for: C<sub>12</sub>H<sub>7</sub>ClN<sub>3</sub>S<sub>2</sub>: C, 49.2; H, 2.4; N, 14.4%. Found: C, 49.1; H, 2.4; N, 14.2%; IR  $\nu_{\max}$ (KBr)/cm<sup>-1</sup> 3443, 3339, 2982, 2210; <sup>1</sup>H NMR  $\delta_{\text{H}}$ (CDCl<sub>3</sub>) 4.38 (2H, s, SCH<sub>2</sub>), 5.42 (2H, s, NH<sub>2</sub>), 7.32 (1H, d, J=8.4 Hz, H-3), 7.68 (1H, dd, J=8.4 & 2.4 Hz, H-4), 8.36 (1H, d, J=2.4 Hz, H-6).

### 1.2.3. **2-Chloro-5-(2-ethyloxycarbonyl-5-ethyloxycarbonyl methylthio-3-hydroxythienyl)pyridine (14)**

Pale yellow crystals (33%, EtOAc). Mp 74-75 °C; Anal. Calcd for: C<sub>16</sub>H<sub>16</sub>ClNO<sub>5</sub>S<sub>2</sub>: C, 47.8; H, 4.0; N, 3.4%. Found: C, 47.8; H, 3.7; N, 3.3%; IR  $\nu_{\max}$ (KBr)/cm<sup>-1</sup> 3462, 2978, 2914, 1748, 1666; <sup>1</sup>H NMR  $\delta_{\text{H}}$ (CDCl<sub>3</sub>) 1.20 (3H, t, J=7.2 Hz, CH<sub>3</sub>), 1.32 (3H, t, J=7.2 Hz, CH<sub>3</sub>), 3.48 (2H, s, SCH<sub>2</sub>), 4.02 (2H, q, J=7.2 Hz, CO<sub>2</sub>CH<sub>2</sub>), 4.24 (2H, q, J=7.2 Hz, CO<sub>2</sub>CH<sub>2</sub>), 7.22 (1H, d, J=8.4 Hz, H-3), 7.66 (1H, dd, J=8.4 & 2.4 Hz, H-4), 8.34 (1H, d, J=2.4 Hz, H-6), 9.70 (1H, s, OH).

### 1.2.4. **2-Chloro-5-(2-cyano-5-cyanomethylthio-3-hydroxythienyl)pyridine (16)**

Pale yellow prisms (46%, EtOAc). Mp 101-102 °C; Anal. Calcd for C<sub>12</sub>H<sub>6</sub>ClN<sub>3</sub>OS<sub>2</sub>: C, 46.8; H, 1.9; N, 13.6%. Found: C, 46.9; H, 1.7; N, 13.5%; IR  $\nu_{\max}$ (KBr)/cm<sup>-1</sup> 3452, 2975, 2926, 2211; <sup>1</sup>H NMR  $\delta_{\text{H}}$

(CDCl<sub>3</sub>) 4.04 (2H, s, SCH<sub>2</sub>), 7.58 (1H, d, J=8.4 Hz, H-3), 7.86 (1H, dd, J=8.4 & 2.4 Hz, H-4), 8.42 (1H, d, J=2.4 Hz, H-6).

## ACKNOWLEDGEMENTS

We thank UAD for the studentships (KRS & GW), and Synthetic Chemicals Ltd for some financial support.

## REFERENCES (AND NOTES)

1. D. H. Bremner, A. D. Dunn, and K. A. Wilson, *Synthesis*, 1992, 528.
2. D. H. Bremner, A. D. Dunn, K. A. Wilson, K. R. Sturrock, and G. Wishart, *Synthesis*, 1997, 949.
3. D. H. Bremner, A. D. Dunn, K. A. Wilson, K. R. Sturrock, and G. Wishart, *Synthesis*, 1998, 1095.
4. S. Okuda and M. M. Robison, *J. Am. Chem. Soc.*, 1959, **81**, 740.
5. R. Gompper, E. Kutter, and W. Toepfl, *Liebigs Ann. Chem.*, 1962, **659**, 90.
6. R. Gompper and H. Schaefer, *Chem. Ber.*, 1967, **100**, 591.
7. M. Augustin, R. Schmidt, and W. Rudolf, *Z. Chem.*, 1977, **17**, 289.
8. K. Clarke, W. R. Fox, and R. M. Scrowston, *J. Chem. Soc., Perkin Trans. 1*, 1980, 1029.
9. M. Kolb, *Synthesis*, 1990, 171.
10. R. M. Mohareb, S. M. Sherif, A. Hashabi, and N. I. Abdel-Sayed, *Coll. Czech. Chem. Comm.*, 1995, **60**, 1578.
11. R. B. Katz, M. B. Mitchell, and P. G. Sammes, *Tetrahedron*, 1989, **45**, 1801.
12. D. Thomae, E. Perspicace, D. Henryon, Z. Xu, S. Schneider, S. Hesse, G. Kirsch, and P. Seck, *Tetrahedron*, 2009, **65**, 10453 (and references therein).
13. A. D. Cross and R. A. Jones, 'An Introduction to Practical Infrared Spectroscopy', Butterworths, London, 1969, pp. 45-100.
14. H. Baron, F. G. P. Remfry, and J. F. Thorpe, *J. Chem. Soc.*, 1904, **85**, 1726.
15. W. Dieckmann, *Ber.*, 1894, **27**, 102.